


# Field Book for Describing and Sampling Soils



Version 3.0

**National Soil Survey Center  
Natural Resources Conservation Service  
U.S. Department of Agriculture**

September 2012; Reprint 2021

# **Field Book for Describing and Sampling Soils**

## **Changes to 3<sup>rd</sup> Printing**

**Version 3.0  
Reprint 2021**



**National Soil Survey Center  
Natural Resources Conservation Service  
U.S. Department of Agriculture  
Lincoln, Nebraska**

# THIS DOCUMENT

The "Field Book for Describing and Sampling Soils, Version 3.0" was first printed in 2012. Due to high demand, it was reprinted in 2016 and again in 2021.

This document lists changes between the second (2016) and third (2021) printings. The changes are primarily errata but, in some cases, replace material that became outdated. The third printing most accurately represents the official material as of 2021.

The third printing is a corrected reprint, not an update.

## **Non-Discrimination Statement**

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](#) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

USDA is an equal opportunity provider, employer, and lender.

## PAGE 1–4

### Second Printing

See the “Geomorphic Description **Section**” for complete lists (p. 3–1).

### Third Printing

See the “Geomorphic Description **System**” for complete lists (p. 3–1).

## PAGE 1–11

### Second Printing

**Subaqueous** Drainage—Free water is above the soil surface.

### Third Printing

**Subaqueous** Drainage—Free water is above the soil surface.

## PAGES 1–11 AND 2–101

### Second Printing

The soils have a peraquic soil moisture regime (proposed 2010; Soil Survey Staff revision online at [soils.usda.gov/technical/manual/proposed\\_changes.html](https://soils.usda.gov/technical/manual/proposed_changes.html)).

### Third Printing

The soils have a peraquic soil moisture regime.

## PAGE 1–18

### Second Printing

Describe the nature of the unconsolidated material (regolith) in which the soil is formed.

### Third Printing

Describe the nature of the unconsolidated material (regolith) in which the soil is formed (**e.g., till**).

## Second Printing

Use the most precise term for the in situ material.

## Third Printing

Use the most precise term possible for the in situ material.

## Second Printing

<b>IGNEOUS—INTRUSIVE</b>			
anorthosite	ANO	pyroxenite	PYX
diabase	DIA	quartzite	QZT
diorite	DIO	quartz-diorite	QZD
gabbro	GAB	quartz-monzonite	QZM
granite	GRA	syenite	SYE
granitoid 2	GRT	syenodiorite	SYD
granodiorite	GRD	tonalite	TON
monzonite	MON	ultramafic rock 2	UMU
peridotite	PER		

## Third Printing

<b>IGNEOUS—INTRUSIVE</b>			
anorthosite	ANO	peridotite	PER
diabase	DIA	pyroxenite	PYX
diorite	DIO	quartz-diorite	QZD
gabbro	GAB	quartz-monzonite	QZM
granite	GRA	syenite	SYE
granitoid 2	GRT	syenodiorite	SYD
granodiorite	GRD	tonalite	TON
monzonite	MON	ultramafic rock 2	UMU

## Second Printing

d) (for submerged soil) the same as b) but refers to the water/soil contact and extends out from shore to the limit of emergent rooted plants;

## Third Printing

d) (for submerged soil) the same as b) but refers to the water/soil contact and extends out from the shore to the limit of rooted plants;

## Second Printing

Concentration or Ped and Void Surface Feature; e.g., carbonate mass, clay film, and organic

## Third Printing

Concentration or Ped and Void Surface Feature; e.g., carbonate mass, clay film, and organic coat

## Second Printing

(<2 mm)

## Third Printing

(>0.25 to <2mm)

## Second Printing

Lining pores (see graphic p. 2-34)

LPO

## Third Printing

Lining pores (see Coats/Films graphic p. 2-34)

LPO

## PAGE 2–39

### Second Printing

Extremely Bouldery	XBY	BYX	≥35% but <60% boulders
--------------------	-----	-----	------------------------

### Third Printing

Very Bouldery	VBY	BYV	≥35% but <60% boulders
Extremely Bouldery	XBY	BYX	≥60% but <90% boulders

## PAGE 2–41

### Second Printing

(COMPOSITIONAL) TEXTURE MODIFIERS 1, 2—Compositional adjectives.

### Third Printing

(COMPOSITIONAL) TEXTURE MODIFIERS 1, 2—Compositional adjectives (e.g., ashy silt loam)

### Second Printing

LIMNIC MATERIALS (used only with Histosols)

### Third Printing

LIMNIC MATERIALS

## PAGE 2–43

### Second Printing

Ice <sup>4, 5</sup> (permanent, subsurface)	ICE
Material <sup>6</sup>	MAT
Water <sup>5</sup> (permanent, subsurface)	W

### Third Printing

Ice <sup>5</sup> (permanent, subsurface)	ICE
Water <sup>4</sup> (permanent, subsurface)	W

## Second Printing

<sup>5</sup> Used for permanent (nonseasonal), massive, subsurface ice; e.g., a glacial layer; proposed in NASIS.

<sup>6</sup> “Material” is used only in combination with Compositional Texture Modifiers (p. 2–41); e.g. *woody material*; *medial material*. In NASIS, “Cemented Material” denotes any cemented soil material (i.e., *duripan*, *ortstein*, *petrocalcic*, *petroferric*, *petrogypsic*).

## Third Printing

<sup>5</sup> Used for permanent (nonseasonal), massive, subsurface ice.

## Second Printing

FLUIDITY <sup>1</sup>		Use a palmful of soil (squeeze in hand)
Nonfluid	NF	After full compression, no soil flows through the fingers.
Slightly Fluid	SF	After full compression is exerted, some soil flows through fingers; most remains in the palm.
Mod. Fluid	MF	After full pressure is exerted, most soil flows through fingers; some remains in the palm.
Very Fluid	VF	Under very gentle pressure, most soil flows through the fingers as a slightly viscous fluid; very little or no residue remains in the palm of the hand.

## Third Printing

FLUIDITY <sup>1</sup>		Use a palmful of soil (squeeze in hand)
Nonfluid	NF	After full compression, no soil flows through the fingers. <b>n value = 0</b>
Slightly Fluid	SF	After full compression is exerted, some soil flows through fingers; most remains in the palm. <b>n value &gt;0 to &lt;0.7</b>
Mod. Fluid	MF	After full pressure is exerted, most soil flows through fingers; some remains in the palm. <b>n value &gt;0.7 to &lt;1.0</b>
Very Fluid	VF	Under very gentle pressure, most soil flows through the fingers as a slightly viscous fluid; very little or no residue remains in the palm of the hand. <b>n value &gt;1.0</b>



## **Second Printing**

(Version 4.2—03/01/2012)

## **Third Printing**

(Version 5.0—08/14/2017)

### **PAGES 3-15 AND 3-32; LANDFORMS**

## **Second Printing**

ledge	LE
-------	----

## **Third Printing**

ledge (also Micro)	LE
--------------------	----

### **PAGE 3-31; LANDSCAPES**

## **Second Printing**

plateau	PT
---------	----

## **Third Printing**

plateau (also LF)	PT
-------------------	----

## IGNEOUS ROCKS CHART

<b>CRYSTALLINE TEXTURE</b>	<b>KEY MINERAL COMPOSITION</b>							
	<i>Acidic</i> (≈felsic) Potassium (K) Feldspar > 2/3 of total Feldspar content		<i>Intermediate</i> (---) Potassium (K) Feldspar and Plagioclase (Na, Ca) Feldspar in about equal proportions		<i>Basic</i> (≈mafic) Plagioclase (Na, Ca) Feldspar > 2/3 of total Feldspar content		<i>Ultrabasic</i> (≈ultramafic) Pyroxene and olivine	
<b>PEGMATITIC</b> (very coarse, uneven-sized crystal grains)	<u>Quartz</u> granite pegmatite	<u>No Quartz</u> syenite pegmatite	<u>Quartz</u> ← monzonite- pegmatite → <u>No Quartz</u>	<u>Quartz</u> ← monzonite- pegmatite → <u>No Quartz</u>	<u>Sodic (Na) Plagioclase</u> <u>Quartz</u>	<u>No Quartz</u> diorite- pegmatite	Calcic (Ca) Plagioclase gabbro pegmatite	peridotite (mostly olivine)
<b>PHANERITIC</b> (crystals visible and of nearly equal size)	granite	syenite	quartz monzonite	monzonite	quartz-diorite granodiorite	diorite	gabbro	pyroxenite (mostly pyroxene)
<b>PORPHYRITIC</b> (relatively few visible crystals within a fine- grained matrix)	granite porphyry	syenite porphyry	quartz monzonite porphyry	monzonite porphyry	quartz-diorite porphyry	diorite porphyry	diabase	} lava <sup>3</sup>
	rhyolite porphyry	trachyte porphyry	quartz- latite porphyry	latite porphyry	dacite porphyry	andesite porphyry	porphyry basalt	
<b>APHANITIC</b> (crystals visible only with magnification) micro <sup>1</sup> crypto <sup>2</sup>	rhyolite	trachyte	quartz latite	latite	dacite	andesite	basalt	
<b>GLASSY</b> (amorphous: no crystalline structure)	obsidian (and its varieties: perlite, pitchstone, pumice, and scoria) pyroclastics are shown on the Sedimentary and Volcaniclastic Rocks chart				<sup>1</sup> Microcrystalline—crystals visible with ordinary magnification (hand lens, simple microscope) <sup>2</sup> Cryptocrystalline—crystals only visible with SEM <sup>3</sup> Lava – generic name for extrusive flows of nonclastic, aphanitic rocks (rhyolite, andesite, and basalt)			

## IGNEOUS ROCKS CHART

<b>CRYSTALLINE TEXTURE</b>	<b>KEY MINERAL COMPOSITION</b>							
	Acidic (Felsic)		INTERMEDIATE				Basic (mafic)	Ultrabasic (ultramafic)
	Potassium (K) Feldspar >2/3 of Total Feldspar Content		Potassium (K) Feldspar and Plagioclase (Na, Ca) Feldspar in about equal proportions		Plagioclase (Na, Ca) Feldspar >2/3 of Total Feldspar Content			Pyroxene and Olivine
	Quartz	No Quartz	Quartz	No Quartz	Sodic (Na) Plagioclase	Calcic (Ca) Plagioclase	peridotite (mostly olivine)	
<b>PEGMATITIC</b> <sup>1</sup>	granite pegmatite	syenite pegmatite	← monzonite pegmatite →		diorite pegmatite	gabbro pegmatite		
<b>PHANERITIC</b> <sup>2</sup>	granite	syenite	quartz monzonite	monzonite	quartz-diorite granodiorite	diorite	gabbro	pyroxenite (mostly pyroxene)
<b>PORPHYRITIC</b> <sup>3</sup>	granite porphyry	syenite porphyry	quartz-monzonite porphyry	monzonite porphyry	quartz-diorite porphyry	diorite porphyry	diabase	} lava <sup>7</sup>
	rhyolite porphyry	trachyte porphyry	quartz-latite porphyry	latite porphyry	dacite porphyry	andesite porphyry	basalt porphyry	
<b>APHANITIC</b> <sup>4</sup> <b>micro</b> <sup>5</sup> <b>crypto</b> <sup>6</sup>	rhyolite	trachyte	quartz latite	latite	dacite	andesite	basalt	
<b>GLASSY</b> <sup>8</sup>	Obsidian (and its varieties: perlite, pitchstone, pumice, scoria) Pyroclastics are shown on the Sedimentary and Volcaniclastic Rocks chart.							
<sup>1</sup> Pegmatitic: Very coarse, uneven-sized crystal grains; 5 to >20 mm <sup>2</sup> Phaneritic: Crystals discernable by eye or 10X lens; 1-5 mm <sup>3</sup> Porphyritic: Larger crystals embedded within a fine-grained matrix <sup>4</sup> Aphanitic: Crystals not visible by eye or 10X lens; <1 mm <sup>5</sup> Microcrystalline crystals resolvable by optical microscope				<sup>6</sup> Cryptocrystalline crystals resolvable by electron microscope <sup>7</sup> Lava: Generic name for extrusive flows of non-clastic, aphanitic rocks (rhyolite, andesite, basalt) <sup>8</sup> Glassy: Noncrystalline or weakly crystalline				

## METAMORPHIC ROCKS CHART

<b>NONFOLIATED STRUCTURE</b>			<b>CRUDE ALIGNMENT</b>	<b>FOLIATED STRUCTURE (e.g., banded)</b>				
<b>CONTACT METAMORPHISM</b>			<b>MECHANICAL METAMORPHISM</b>	<b>REGIONAL METAMORPHISM</b>		<b>PLUTONIC METAMORPHISM</b>		
<i>Low Grade</i>	<i>Medium Grade</i>	<i>High Grade</i>	<i>Very Low Grade</i>	<i>Low Grade</i>	<i>Medium Grade</i>	<i>High Grade</i>	<i>Extreme Grade</i>	
	granofels hornfels marble metaquartzite serpentinite soapstone (talc)		crush breccia mylonite  <-- metaconglomerate -->  <----- metavolcanics ----->	slate	phyllite greenstone	schist amphibolite	gneiss granulite	migmatite
<p>* Not all rock types listed here can be definitively identified in the field (e.g., may require grain counts).</p> <p>** Not all rock types shown here are available on Bedrock - Kind choice list. They are included here for completeness and as aids to using geologic literature.</p>								

(Schoeneberger and Wysocki, 1998)

## METAMORPHIC ROCKS CHART

[Not all rock types listed here can be definitively identified in the field (e.g., may require grain counts). Not all rock types shown here are available on Bedrock - Kind choice list. They are included here for completeness and as aids to using geologic literature.]

<b>NONFOLIATED STRUCTURE</b>			<b>CRUDE ALIGNMENT</b>	<b>FOLIATED STRUCTURE (e.g., banded)</b>				
<b>CONTACT METAMORPHISM</b>			<b>FAULT ZONE METAMORPHISM</b>	<b>REGIONAL METAMORPHISM</b>		<b>PLUTONIC METAMORPHISM</b>		
<i>Low Grade</i>	<i>Medium Grade</i>	<i>High Grade</i>	<i>Low Grade</i>	<i>Low Grade</i>	<i>Medium Grade</i>	<i>High Grade</i>		
	granofels hornfels marble metaquartzite serpentinite soapstone (talc)		crush breccia mylonite	slate	phyllite greenstone	schist amphibolite	gneiss granulite	migmatite*
				<-----Metaconglomerate----->				
				<-----Metavolcanics----->				
				<-----Metasedimentary----->				
* Partial melting occurs.								

(Schoeneberger and Wysocki, 1998)

**SEDIMENTARY AND VOLCANICLASTIC ROCKS**

CLASTIC				NONCLASTIC		
Dominant Grain Size				Chemical	Biochemical	Organic
Very Fine	Fine	Medium	Coarse	Evaporates, Precipitates	Accretionates	Reduzates
<----- (Argillaceous) -----> < 0.002 mm	0.002 - 0.06 mm	(Arenaceous) 0.06 - 2.0 mm	(Rudaceous) >2.0 mm	anhydrite (CaSO <sub>4</sub> )  gypsum (CaSO <sub>4</sub> • 2H <sub>2</sub> O)  halite (NaCl)	<b>CARBONATE ROCKS</b> Limestones (ls) (>50% calcite)  <u>chemical types</u> caliche travertine tufa  <u>accretionary types</u> biostromal ls organic reef pelagic ls (chalk) <u>bio-clastic types</u> coquina oolithic ls lithographic ls  <u>altered types</u> dolomite (>50% calcite + dolomite) phosphatic limestone	black shale (organics and fine sediments)  bituminous ls bog iron ores  coal
<b>VOLCANICLASTICS</b> (includes Pyroclastics)						
<----- ignimbrite ----->		agglomerate (rounded frags)				
<----- tuff ----->		volcanic breccia (angular frags)				
<----- pumice (specific gravity <1.0; highly vesicular)----->						
<----- scoria (specific gravity >2.0; slightly or moderately vesicular) ----->						
				<b>OTHER NONCLASTIC ROCKS</b>		
				Siliceous rocks (SiO <sub>2</sub> dominated): chert (jasper, chalcedony, opal) diatomite rock phosphate iron-bearing rocks (Fe-SiO <sub>2</sub> dominated)		

## SEDIMENTARY AND VOLCANICLASTIC ROCKS

CLASTIC				NONCLASTIC		
Dominant Grain Size				Chemical	Biochemical	Organic
Very Fine	Fine	Medium	Coarse	Evaporates, Precipitates	Accretionates	Reduzates
<---- (Argillaceous) ----> < 0.002 mm	0.002 - 0.06 mm	(Arenaceous) 0.06 - 2.0 mm	(Rudaceous) >2.0 mm			
<----- argillite -----> (more indurated, less laminated and fissile)		Sandstones (ss):  arenite arkose (mainly feldspar) glaucconitic ss ("greensand") graywacke (dark, "dirty" ss) orthoquartzite (mainly quartz)	breccia (nonvolcanic, angular frags)	anhydrite (CaSO <sub>4</sub> )	<b>CARBONATE ROCKS</b> Limestones (ls) (>50% calcite)	
<----- shale -----> (laminated, fissile)			conglomerate (nonvolcanic, rounded frags)	gypsum (CaSO <sub>4</sub> • 2H <sub>2</sub> O)		chemical types caliche travertine tufa
<----- mudstone -----> (nonlaminated, nonfissile) (≈ equal clay and silt)				halite (NaCl)		bituminous ls bog iron ores
claystone (non-laminated, nonfissile)	siltstone (nonlaminated, nonfissile)					coal
VOLCANICLASTICS (includes Pyroclastics)						
<----- ignimbrite -----> (mainly pumice frags; consolidated pyroclastic flows)		agglomerate (rounded frags)				
<----- tuff -----> (consolidate volcanic ash, tephra)		volcanic breccia (angular frags)				
<----- pumice (specific gravity <1.0; highly vesicular)----->				dolomite (>50% CaMg(CO <sub>3</sub> ))		
<----- scoria (specific gravity >2.0; slightly or moderately vesicular) ----->				phosphatic limestone		
				OTHER NONCLASTIC ROCKS		
				Siliceous rocks (Silica dominated): chert (jasper, chalcedony, opal); diatomite Rock phosphate Iron-bearing rocks (Fe-SiO <sub>2</sub> dominated): jaspilite, specular hematite, magnetite		

(Schoeneberger and Wysocki, 2000)

## **Second Printing**

(Doerr et al., 2000)  
(Robichand and Miller, 1999)  
(Robichand et al., 2008)

## **Third Printing**

(Doerr et al., 2006)  
(Robichand and Miller, 1999)  
(Robichand et al., 2008)

## **Second Printing**

- 2) Use an eyedropper or plastic squeeze bottle to randomly place 5 drops of distilled water (approximately 5 mm in diameter) from a 1-cm height onto the prepared surface.

Modified from Robichand, 2008

## **Third Printing**

- 2) Use an eyedropper or plastic squeeze bottle to randomly place 5 drops of distilled water (each drop approximately 5 mm in diameter) from a 1-cm height onto the prepared surface.

Modified from Robichand, 2008

## **Second Printing**

It is advisable to subsample soil horizons about 50 cm thick.

## **Third Printing**

It is advisable to subsample soil horizons >50 cm thick.



# ACKNOWLEDGMENTS

The science and knowledge in this document are distilled from the collective experience of thousands of dedicated soil scientists during the more than 100 years of the National Cooperative Soil Survey (NCSS) program. A special thanks is due to these largely unknown stewards of the natural resources of this nation.

Special thanks and recognition are extended to those who contributed extensively to the preparation and production of this book: the soil scientists from the NRCS and NCSS cooperators who reviewed and improved it; Tammy Umholtz for document preparation and graphics; and the NRCS Soil Science Division for funding it.

Proper citation for this document is:

*Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. Field book for describing and sampling soils, Version 3.0. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.*

---

**Cover Photo:** A polygenetic Calcic Argiustoll with an A, Bt, Bk, 2BC, 2C horizon sequence. This soil formed in Peoria Loess that blankets the fluvial Ash Hollow Formation of the Ogallala Group. It occurs in an undulating area of the Cheyenne Tablelands in northern Banner County, Nebraska. The scale is in meters. (Photo by Doug Wysocki, NRCS, Lincoln, NE, June 2011.)

---

*Trade names are used solely to provide specific information. Mention of a trade name does not constitute a guarantee of the product by the U.S. Department of Agriculture nor does it imply endorsement by the Department or the Natural Resources Conservation Service over comparable products that are not named.*

*The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex (including gender identity and expression), marital status, familial status, parental status, religion, sexual orientation, political beliefs, genetic information, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to: USDA, Assistant Secretary for Civil Rights, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, S.W., Stop 9410, Washington, D.C. 20250-9410, or call toll-free at (866) 632-9992 (English) or (800) 877-8339 (TDD) or (866) 377-8642 (English Federal-relay) or (800) 845-6136 (Spanish Federal-relay). USDA is an equal opportunity provider and employer.*

# FOREWORD

**Purpose:** The following instructions, definitions, concepts, and codes are a field guide for making or reading soil descriptions and sampling soils as presently practiced in the USA. (Note: References cited in the Foreword are listed at the end of Chapter 1 [p. 1–31].)

**Background:** Soil description methodology was developed by soil scientists throughout the entire course of the soil survey. The USDA published small instruction booklets for field parties, including soil descriptions, in 1902–1904, 1906, and 1914. The first USDA guide for soil horizon identification and description was released in 1937 (Bureau of Chemistry and Soils, 1937). Dr. Roy Simonson and others later summarized and revised this information (Soil Survey Staff, 1951; Soil Survey Staff, 1962). Brief “color-book” inserts with shorthand notation were released by the Soil Conservation Service (Spartanburg, SC, 1961; Western Technical Center, Portland, OR, 1974). Previous Field Books were released in 1998 (Schoeneberger et al.) and 2002 (Schoeneberger et al.). This is an updated Field Book version that summarizes current knowledge, includes updates since 2002, and reflects changes in source documents.

**Standards:** This Field Book summarizes and updates current National Cooperative Soil Survey conventions for describing soils (Soil Survey Manual [Soil Survey Division Staff, 1993]; National Soil Survey Handbook [Soil Survey Staff, 2012d]; National Soil Information System (NASIS), release 6.2 [Soil Survey Staff, 2012c]; and NASIS Data Dictionary [Soil Survey Staff, 2012a]). Some content is an abbreviation of primary sources.

**Regarding Pedon PC and NASIS:** The Field Book is a current, practical soil description guide for the soil science community. It is not a guide on “How To Use Pedon PC or NASIS.” Differences and linkages between soil science conventions, Pedon PC, NASIS, and older systems are shown, where reasonable to do so, as an aid for interpreting and converting archived data.

Standard procedures and terms for describing soils have changed and increased in recent years (e.g., redoximorphic features). Coincident with these changes has been the development and use of computer databases to store soil descriptions and associated information. The nature of databases, for better or worse, requires consistent and “correct” use of terms.

**Sources:** This Field Book draws from several primary sources: The Soil Survey Manual (Soil Survey Division Staff, 1993) and the National Soil Survey Handbook (NSSH), Parts 618 and 629 (Soil

Survey Staff, 2012d). Other important sources are footnoted throughout to give appropriate credit and encourage in-depth information review. Other material is unique to this book.

**Brevity:** In a field book, brevity is efficiency. Despite this book's apparent length, the criteria, definitions, and concepts are condensed. We urge users to review the comprehensive information in original sources to avoid errors resulting from our brevity.

**Measurement Units:** For soil description, metric units are the scientific standard. Both NASIS and Pedon PC use metric units.

**Format:** The "Site Description" and "Profile Description" sections generally follow conventional profile description format and sequence (e.g., SCS-Form 232, December 1984). Some descriptors are arranged in a sequence more compatible with field description rather than data entry (e.g., **Horizon Boundary** is next to **Horizon Depth**, rather than at the end). The sequence followed differs somewhat from and does *not* supersede convention for writing formal soil descriptions in soil survey reports or Official Soil Series Descriptions (e.g., National Soil Survey Handbook, Part 614; Soil Survey Staff, 2012d).

**Codes:** Shorthand notation is listed in the *Code* column for some descriptors. Long-standing conventional codes are retained because of widespread recognition. Some recent codes have been changed to make them more logical. Some data elements have different codes in various systems (e.g., conventional [Conv.] vs. NASIS vs. Pedon PC), and several columns may be shown to facilitate conversions. If only one code column is shown, it can be assumed that the conventional, NASIS, and Pedon PC codes are all the same.

**Standard Terms vs. Creativity:** *Describe and record what you observe.* Choice lists in this document are a minimal set of descriptors. Use additional descriptors, notes, and sketches to record pertinent information and/or features if no data element or choice list entry exists. Record such information as free-hand notes under **Miscellaneous Field Notes**.

**Changes:** Soil science is an evolving field. Changes to this Field Book should and will occur. Please send comments or suggestions to the Director, National Soil Survey Center, USDA-NRCS; 100 Centennial Mall North, Rm. 152; Lincoln, NE 68508-3866.

# TABLE OF CONTENTS

<b>ACKNOWLEDGMENTS</b> . . . . .	<b>i</b>
<b>FOREWORD</b> . . . . .	<b>ii</b>
<b>SITE DESCRIPTION</b> . . . . .	<b>1-1</b>
<b>Describer Name(s)</b> . . . . .	<b>1-1</b>
<b>Date</b> . . . . .	<b>1-1</b>
<b>Climate</b> . . . . .	<b>1-1</b>
(Weather Conditions, Air Temperature, Soil Temperature [Soil Temperature, Soil Temperature Depth])	
<b>Location</b> . . . . .	<b>1-2</b>
(Latitude, Longitude, Geodetic Datum)	
<b>Topographic Quadrangle</b> . . . . .	<b>1-2</b>
<b>Soil Survey Site Identification Number (Site ID)</b> . . . . .	<b>1-2</b>
<b>County FIPS Code</b> . . . . .	<b>1-3</b>
<b>MLRA</b> . . . . .	<b>1-3</b>
<b>Transects</b> . . . . .	<b>1-3</b>
(Transect ID, Stop Number, Interval)	
<b>Series or Component Name</b> . . . . .	<b>1-4</b>
(Map Unit Symbol, Photo #)	
<b>Geomorphic Information</b> . . . . .	<b>1-4</b>
Physiographic Location . . . . .	1-4
(Physiographic Division, Physiographic Province, Physiographic Section, State Physiographic Area, Local Physiographic/Geographic Name)	
Geomorphic Description . . . . .	1-4
(Landscape, Landform, Microfeature, Anthropogenic Feature)	
Surface Morphometry . . . . .	1-5
(Elevation, Slope Aspect, Slope Gradient, Slope Complexity, Relative Slope Segment Position, Slope Shape, Hillslope - Profile Position, Geomorphic Components [Hills, Terraces and Stepped Landforms, Mountains, Flat Plains], Microrelief, Drainage Pattern)	
<b>Water Status</b> . . . . .	<b>1-11</b>
Drainage . . . . .	1-11
Flooding . . . . .	1-13
(Frequency, Duration, Months)	
Ponding . . . . .	1-14
(Frequency, Depth, Duration)	
(Soil) Water State . . . . .	1-14

<b>Land Cover</b> . . . . .	<b>1-16</b>
(Earth Cover - Kind)	
<b>Vegetation</b> . . . . .	<b>1-17</b>
(Plant Symbol, Plant Common Name, Plant Scientific Name, Vegetation Cover)	
<b>Parent Material</b> . . . . .	<b>1-18</b>
(Kind)	
<b>Bedrock</b> . . . . .	<b>1-22</b>
(Kind, Fracture Interval Class, Weathering Class, Depth [to Bedrock])	
<b>Lithostratigraphic Unit(s)</b> . . . . .	<b>1-25</b>
<b>Erosion</b> . . . . .	<b>1-25</b>
(Kind, Degree Class)	
<b>Surface Fragments</b> . . . . .	<b>1-26</b>
(Kind, Surface Fragment Class)	
<b>Diagnostic Horizons or Characteristics</b> . . . . .	<b>1-28</b>
(Kind, Depth, Soil Taxonomy Classification, Particle-Size Control Section)	
<b>Restriction</b> . . . . .	<b>1-30</b>
(Kind, Hardness)	
<b>References</b> . . . . .	<b>1-31</b>
<b>PROFILE/PEDON DESCRIPTION</b> . . . . .	<b>2-1</b>
<b>Observation Method</b> . . . . .	<b>2-1</b>
(Kind, Relative Size)	
<b>Horizon and Layer Designations</b> . . . . .	<b>2-2</b>
Master and Transitional Horizons and Layers . . . . .	2-2
Horizon Suffixes . . . . .	2-4
Other Horizon Modifiers . . . . .	2-5
(Numerical Prefixes, Numerical Suffixes, The Prime, The Caret)	
Horizon Depth . . . . .	2-6
Horizon Thickness . . . . .	2-6
Horizon Boundary . . . . .	2-6
(Distinctness, Topography)	
<b>Soil Color</b> . . . . .	<b>2-8</b>
Decision Flowchart for Describing Soil Colors . . . . .	2-8
(Soil) Matrix Color . . . . .	2-9
([Soil] Color, Moisture State, Location or Condition)	
<b>Redoximorphic Features—RMFs (Discussion)</b> . . . . .	<b>2-10</b>
<b>Redoximorphic Features</b> . . . . .	<b>2-12</b>
(Kind, Quantity, Size, Contrast, Color, Moisture State, Shape, Location, Hardness, Boundary)	
Tabular List for Determination of Color Contrast . . . . .	2-16

<b>Mottles</b> . . . . .	<b>2-18</b>
(Quantity, Size, Contrast, Color, Moisture State, Shape, Location)	
<b>Concentrations (Discussion)</b> . . . . .	<b>2-19</b>
<b>Concentrations</b> . . . . .	<b>2-20</b>
(Kind, Quantity [Percent of Area Covered], Size, Contrast, Color, Moisture State, Shape, Location, Hardness, Boundary)	
<b>Pedogenic Carbonate Stages (Discussion)</b> . . . . .	<b>2-28</b>
(Development, Multiple Stages, Description)	
Pedogenic Carbonate Development Stages -	
Fine Earth Matrix . . . . .	2-30
Pedogenic Carbonate Development Stages -	
Coarse Fragment Matrix . . . . .	2-31
<b>Ped and Void Surface Features</b> . . . . .	<b>2-32</b>
(Kind, Amount, Continuity, Distinctness, Location, Color)	
<b>Soil Texture</b> . . . . .	<b>2-36</b>
Texture Class . . . . .	2-37
(Soil) Textural Triangle (Fine Earth). . . . .	2-38
Texture Modifiers . . . . .	2-38
(Quantity and Size, Compositional, Terms Used in Lieu of Texture)	
Comparison of Particle Size Classes in Different Systems . . . . .	
	2-45
<b>Rock and Other Fragments</b> . . . . .	<b>2-46</b>
(Kind, Volume Percent, Size Classes and Descriptive Terms, Roundness, Hardness)	
<b>Artifacts (<i>Human-derived</i>)</b> . . . . .	<b>2-49</b>
(Kind, Quantity, Roundness, Shape, Cohesion, Penetrability, Persistence, Safety)	
<b>(Soil) Structure</b> . . . . .	<b>2-52</b>
(Type, Grade, Size)	
<b>Consistence</b> . . . . .	<b>2-62</b>
Rupture Resistance . . . . .	2-62
(Blocks, Peds, and Clods; Surface Crusts and Plates)	
Cementing Agents. . . . .	2-64
Manner of Failure . . . . .	2-65
Stickiness. . . . .	2-66
Plasticity . . . . .	2-66
Penetration Resistance. . . . .	2-67
Penetration Orientation . . . . .	2-68
Excavation Difficulty . . . . .	2-69

<b>Roots</b> .....	<b>2-70</b>
(Quantity, Size, Quantity [graphic], Location)	
<b>Pores (Discussion)</b> .....	<b>2-73</b>
<b>Pores</b> .....	<b>2-73</b>
(Quantity, Size, Shape, Vertical Continuity)	
<b>Cracks</b> .....	<b>2-75</b>
(Kind, Depth, Relative Frequency)	
<b>Soil Crusts (Discussion)</b> .....	<b>2-77</b>
<b>Soil Crusts</b> .....	<b>2-79</b>
(Kind)	
<b>Special Features</b> .....	<b>2-80</b>
(Kind, Area [%] Occupied)	
<b>Saturated Hydraulic Conductivity and Permeability (Discussion)</b> .....	<b>2-81</b>
<b>Saturated Hydraulic Conductivity (<math>K_{sat}</math>)</b> .....	<b>2-83</b>
<b>Permeability Classes</b> .....	<b>2-85</b>
<b>Chemical Response</b> .....	<b>2-85</b>
Reaction (pH) .....	2-85
pH Method .....	2-86
Effervescence .....	2-87
(Class, Location, Chemical Agent)	
Reduced Conditions .....	2-88
(Dipyridyl - Location)	
Salinity Class (Discussion) .....	2-88
Salinity Class .....	2-89
Sodium Adsorption Ratio (SAR) .....	2-89
<b>Odor</b> .....	<b>2-90</b>
(Kind, Intensity)	
<b>Miscellaneous Field Notes</b> .....	<b>2-90</b>
<b>Minimum Data Set (for a soil description)</b> .....	<b>2-90</b>
<b>Pedon Description Data Sheet</b> .....	<b>2-91</b>
<b>Pedon Description Example</b> .....	<b>2-91</b>
Pedon Description (Data Sheet - Blank) .....	2-93
Pedon Description (Data Sheet - Example) .....	2-95
<b>Subaqueous Soils (SAS) Description</b> .....	<b>2-97</b>
(Discussion, Description)	
Bathymetry .....	2-97
Site .....	2-98
Water Column Measurements .....	2-102
Soil Profile Measurements and Description .....	2-103
Salinity (of Subaqueous Soils) .....	2-106
Subaqueous Soils Profile Description Data Sheet .....	2-109
Subaqueous Soils Profile Description Example .....	2-110

<b>Vibracore Sampling for Subaqueous Soils.</b> . . . . .	<b>2-111</b>
Discussion . . . . .	2-111
Site Description . . . . .	2-111
Core Descriptions . . . . .	2-111
Vibracore Log Sheet . . . . .	2-113
Vibracore Log Sheet Example . . . . .	2-114
<b>References.</b> . . . . .	<b>2-115</b>
<b>GEOMORPHIC DESCRIPTION.</b> . . . . .	<b>3-1</b>
<b>Part I: Physiographic Location</b> . . . . .	<b>3-2</b>
<b>Part II: Geomorphic Description (Outline)</b> . . . . .	<b>3-10</b>
<b>Part II: Geomorphic Description</b> . . . . .	<b>3-11</b>
Comprehensive Lists . . . . .	3-11
Geomorphic Environments and Other Groupings . . . . .	3-21
<b>Part III: Surface Morphometry</b> . . . . .	<b>3-39</b>
<b>References.</b> . . . . .	<b>3-46</b>
<b>SOIL TAXONOMY</b> . . . . .	<b>4-1</b>
<b>Introduction</b> . . . . .	<b>4-1</b>
<b>Horizon and Layer Designations</b> . . . . .	<b>4-1</b>
Master and Transitional Horizons or Layers. . . . .	4-1
Horizon Suffixes . . . . .	4-3
Horizon & Layer Designations Conversion Charts . . . . .	4-6
(Soil) Textural Triangle: Family Particle-Size Classes . . . . .	4-10
Combined Textural Triangles: Fine Earth Texture Classes and Family Particle-Size Classes. . . . .	4-11
Soil Moisture Regimes . . . . .	4-11
Soil Temperature Regimes and Classes . . . . .	4-13
<b>References.</b> . . . . .	<b>4-14</b>
<b>GEOLOGY</b> . . . . .	<b>5-1</b>
<b>Introduction</b> . . . . .	<b>5-1</b>
Bedrock [Kind] . . . . .	5-1
<b>Rock Charts</b> . . . . .	<b>5-4</b>
Igneous Rocks Chart . . . . .	5-5
Metamorphic Rocks Chart. . . . .	5-6
Sedimentary and Volcaniclastic Rocks . . . . .	5-7
Mass Movement (Mass Wasting) Types for Soil Survey. . . . .	5-8
North American Geologic Time Scale . . . . .	5-9
<b>Till Terms.</b> . . . . .	<b>5-10</b>
<b>Pyroclastic Terms</b> . . . . .	<b>5-11</b>
<b>Hierarchical Rank of Lithostratigraphic Units</b> . . . . .	<b>5-12</b>
<b>References.</b> . . . . .	<b>5-13</b>



<b>LOCATION</b> .....	<b>6-1</b>
<b>GPS Location</b> .....	<b>6-1</b>
<b>Public Land Survey</b> .....	<b>6-2</b>
Townships and Ranges .....	6-3
Sections .....	6-3
Section Subdivisions .....	6-4
<b>Universal Transverse Mercator (UTM) Rectangular Coordinate System</b> .....	<b>6-5</b>
<b>State Plane Coordinate System</b> .....	<b>6-7</b>
<b>References</b> .....	<b>6-7</b>
<b>MISCELLANEOUS</b> .....	<b>7-1</b>
<b>Percent of Area Covered</b> .....	<b>7-1</b>
<b>K<sub>sat</sub> Class Estimate</b> .....	<b>7-10</b>
<b>Soil Water Repellency (Discussion)</b> .....	<b>7-14</b>
<b>Soil Water Repellency</b> .....	<b>7-15</b>
<b>Measurement Equivalents and Conversions</b> .....	<b>7-16</b>
Metric to English .....	7-16
English to Metric .....	7-17
Common Conversion Factors .....	7-18
<b>Guide to Map Scales and Minimum Size Delineations</b> ..	<b>7-21</b>
<b>Common Soil Map Symbols (Traditional)</b> .....	<b>7-22</b>
<b>References</b> .....	<b>7-28</b>
<b>SOIL SAMPLING</b> .....	<b>8-1</b>
<b>Introduction</b> .....	<b>8-1</b>
<b>Types of Sampling</b> .....	<b>8-1</b>
(Horizon Sampling, Incremental Sampling, Fixed- Depth Sampling)	
<b>Sampling Techniques</b> .....	<b>8-3</b>
<b>Soil Sample Kinds</b> .....	<b>8-3</b>
(Characterization Samples, Reference Samples)	
<b>Field Equipment Checklist</b> .....	<b>8-4</b>
<b>Examples of Common Soil-Sampling Equipment</b> .....	<b>8-5</b>
<b>References</b> .....	<b>8-6</b>
<b>INDEX</b> .....	<b>9-1</b>



# SITE DESCRIPTION

*P.J. Schoeneberger, D.A. Wysocki, and E.C. Benham, NRCS, Lincoln, NE*

## DESCRIBER NAME(S)

---

**NAME** (or initials)—Record the observer(s) making the description; e.g., *Erling E. Gamble* or *EEG*.

## DATE

---

**MONTH/DAY/YEAR**—Record the observation date. Use numeric notation (MM/DD/YYYY); e.g., *05/21/2012* (for May 21, 2012).

## CLIMATE

---

Document the prevailing weather conditions at time of observation (a site condition that affects some field methods; e.g.,  $K_{sat}$ ). Record the major **Weather Conditions** and **Air Temperature**; e.g., *Rain, 27 °C*.

Weather Conditions	Code
sunny/clear	SU
partly cloudy	PC
overcast	OV
rain	RA
sleet	SL
snow	SN

**AIR TEMPERATURE**—Ambient air temperature at chest height (Celsius or Fahrenheit); e.g., *27 °C*.

**SOIL TEMPERATURE**—Record the ambient **Soil Temperature** and **Depth** at which it is determined; e.g., *22 °C, 50 cm*. (**NOTE:** Soil taxonomy generally requires a 50 cm depth.) Soil temperature should only be determined from a freshly excavated surface that reflects the ambient soil conditions. Avoid surfaces equilibrated with air temperatures.

**Soil Temperature**—Record soil temperature (in °C or °F).

**Soil Temperature Depth**—Record depth at which the ambient soil temperature is measured; e.g., *50 cm*.

## LOCATION

---

Record precisely the point or site location (e.g., coordinates). Latitude and longitude as measured with a Global Positioning System (GPS) is the preferred descriptor. Report lat. and long. as degrees, minutes, seconds, and decimal seconds with direction, or as degrees and decimal degrees with direction. For example:

**LATITUDE**— $46^{\circ} 10' 19.38'' N$ . or  $46^{\circ}.17205 N$

**LONGITUDE**— $95^{\circ} 23' 47.16'' W$ . or  $95^{\circ}.39643 W$

**GEODETTIC DATUM (Horizontal\_datum\_name in NASIS)**—A geodetic datum must accompany latitude and longitude. A geodetic datum is a model that defines the earth's shape and size and serves as a latitude, longitude reference. Geodetic datum is a selectable GPS parameter. The preferred datum is the **World Geodetic System 1984 (WGS-84)**. See "Location Section" for the complete geodetic datum list (p. 6–1).

Topographic maps display latitude and longitude and the geodetic datum employed (e.g., *NAD 27*, *NAD 83*). **NOTE:** NASIS requires latitude and longitude but allows other coordinate or location descriptors (e.g., *UTM*, *State Plane Coordinates*, *Public Land Survey*, *Metes and Bounds*). See "Location Section" (p. 6–1) for details.

## TOPOGRAPHIC QUADRANGLE

---

Record the topographic map name (USGS quadrangle) that covers the observation site. Include scale (or "series") and year printed; e.g., *Pollard Creek-NW; TX; 1:24,000; 1972*.

## SOIL SURVEY SITE IDENTIFICATION NUMBER (SITE ID)

---

An identification number must be assigned if samples are collected (called **User\_Pedon\_ID** in NASIS). For the Kellogg Soil Survey Laboratory (Soil Survey Staff, 2011), this identifier consists of five required and one optional item.

Example: S2004WA27009

- 1) S indicates a sampled pedon. ("S" is omitted for pedons described but *not* sampled.)
- 2) 2004=calendar year sampled. Use 4-digit format; e.g., 2012.
- 3) WA=two-character (alphabetic) Federal Information Processing Standards (FIPS 6-4) code for the state where sampled. For non-U.S. sites, use the Country Code from ISO

3166-1 (International Organization for Standards, 2012b); e.g., *CA* for Canada.

- 4) 027=3-digit (numeric) FIPS code for county where sampled. For non-U.S. sites, use the appropriate two- or three-letter Administrative Subdivision code from ISO 3166-2 (International Organization for Standards, 2012b) preceded by a 0 (zero) for two-letter codes; e.g., *0SK* for Saskatchewan.
- 5) 009=consecutive pedon number for calendar year for county. This should be a 3-digit number. Use 0s (zeros) as placeholders when necessary; e.g., 9 becomes *009*.
- 6) (Optional) A one-character "satellite" code can be used, if needed, to indicate a relationship between a primary pedon and satellite sample points; e.g., *A* in *S2004WA027009A*.

**NOTE:** Do not use spaces, dashes, or hypens (for database reasons). Use uppercase letters. A complete example is *S2011OK061005A*. A sampled soil characterization pedon collected in 2011 (*2011*) from Oklahoma (*OK*), Haskell County (*061*); this is a satellite pedon (*A*) of the fifth pedon (*005*) sampled in that county during 2011.

## **COUNTY FIPS CODE**

---

The Federal Information Processing Standards (FIPS) code is a 3-digit number for a county within a state in the U.S. (National Institute of Standards and Technology, 1990). Record the FIPS code for the county where the pedon or site occurs; e.g., *061* (Haskell County, OK). For non-U.S. sites, use the appropriate two- or three-character Country Code (International Organization for Standards-Country Codes ISO 3166-1; 2012a or current date).

## **MLRA**

---

This 1- to 3-digit number, often including one alpha character, identifies the Major Land Resource Area (NRCS, 2006); e.g., *58C* (Northern Rolling High Plains, Northeastern Part).

## **TRANSECTS**

---

If a soil description is one of multiple transect points, record transect information; e.g., **Transect ID**, **Stop Number**, **Interval**, **GPS Coordinates**. NASIS also accommodates **Transect Kind** (random point [*-R*], regular interval [*-I*]), **Transect Section Method** (biased [*-B*], random [*-R*]), **Delineation Size** (acres), **Transect Direction** (azimuth heading, e.g., 180°).

**TRANSECT ID**—A 4- or 5-digit number that identifies the transect; e.g., 0010 (the tenth transect within the survey area).

**STOP NUMBER**—If the sample/pedon is part of a transect, enter the 2-digit stop number along the transect; e.g., 07. (**NOTE:** NASIS allows up to 13 characters.)

**INTERVAL**—Record distance between observation points, compass bearing, and GPS coordinates, or draw a route map in the **Field Notes** (“User Defined Section”).

## ***SERIES OR COMPONENT NAME***

---

Assign the appropriate Soil Series or Map Unit Component name at time of description (e.g., *Cecil*). If unknown, enter *SND* for “Series Not Designated.” (In NASIS, “SND” is not used; assign an appropriate soil taxonomy class; e.g., *Udorthents*.) **NOTE:** A field-assigned series name may change after additional data collection and lab analyses.

**MAP UNIT SYMBOL**—Record the soil map unit symbol (if known) for the sample site.

**PHOTO #**—If aerial imagery is used, record the photograph number that covers the sample site.

## ***GEOMORPHIC INFORMATION***

---

See the “Geomorphic Description System” for complete lists (p. 3–1). Codes follow each listed choice. Conventionally, the entire name (e.g., *mountains*) is recorded.

### **PART 1: PHYSIOGRAPHIC LOCATION**

**Physiographic Division**—e.g., *Interior Plains* or *IN*

**Physiographic Province**—e.g., *Central Lowland* or *CL*

**Physiographic Section**—e.g., *Wisconsin Driftless Section* or *WDS*

**State Physiographic Area (Opt.)**—e.g., *Wisconsin Dells*

**Local Physiographic/Geographic Name (Opt.)**—e.g., *Bob’s Ridge*

### **PART 2: GEOMORPHIC DESCRIPTION**

**Landscape**—e.g., *Foothills* or *FH*

**Landform**—e.g., *Ridge* or *RI*


**Microfeature**—e.g., *Mound* or *MO*

**Anthropogenic Feature**—e.g., *sanitary landfill* or *SL*

### PART 3: SURFACE MORPHOMETRY


**Elevation**—The height of a point on the earth’s surface relative to Mean Tide Level (MTL), formerly Mean Sea Level (MSL). Record units; e.g., *106 m* or *348 ft*. Recommended methods: interpolation from topographic map contours; altimeter reading tied to a known elevation datum. **NOTE:** An elevation value from a GPS can be recorded. Since the GPS elevation value typically is less certain than the latitude and longitude values, a correction for quantifiable errors is important (e.g., WAAS, or averaging many elevation values at a point by collecting a track log at the point and averaging the elevation values). The latitude and longitude coordinates can be used to extract an elevation value from a DEM, if available. Note that all parts of a DEM cell return the same elevation value, so a higher resolution DEM is important for accuracy, especially if the point is on a steep slope.

**Slope Aspect**—The compass direction (in degrees and accounting for declination) that a slope faces, viewed downslope; e.g., *225°*.



**Slope Gradient**—The ground surface inclination with respect to the horizontal plane; commonly called “slope.” Make observations downslope to avoid errors from clinometer types; e.g., *18%*.


**Slope Complexity**—Describe the relative ground surface uniformity (smooth linear or curvilinear=*simple* or *S*) or irregularity (*complex* or *C*) downslope through the site; e.g., *simple* or *S*.



**Relative Slope Segment Position** (called **geomorph\_slope\_segment** in NASIS)—If useful to subdivide long slopes, describe relative slope location of the area or point of interest.

Relative Slope Segment Position	Code	Criteria
lower third	LT	on lower third of slope
middle third	MT	on middle third of slope
upper third	UT	on upper third of slope


**Slope Shape**—Slope shape is described in two directions: up and down slope (perpendicular to the elevation contour) and across slope (along the elevation contour); e.g., *Linear, Convex* or *LV*.






**Hillslope-Profile Position** (commonly called Hillslope Position)—Two-dimensional geomorphic descriptors that are segments (i.e., slope position) along a line that runs up and down slope; e.g., *backslope* or *BS*. This is best applied to points, not areas (e.g., map units).

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS




**Geomorphic Component**—Three-dimensional geomorphic descriptors for landforms, landform portions, or microfeatures that are applied to areas. Unique 3D descriptors are defined for Hills, Terraces and Stepped Landforms, Mountains, and Flat Plains; e.g., *Hills-nose slope*, or *NS*.


Hills	NASIS Code
interfluve	IF
crest	CT
head slope	HS
nose slope	NS
side slope	SS
free face	FF
base slope	BS




Terraces and Stepped Landforms	Code
riser	RI
tread	TR



<b>Mountains</b>	<b>Code</b>
mountaintop	MT
mountainflank	MF
upper third—mountainflank	UT
center third—mountainflank	CT
lower third—mountainflank	LT
free face	FF
mountainbase	MB



Flat Plains	Code
dip	DP
rise	RI
talf	TF



**Microrelief**—Small, relative elevation differences between adjacent areas on the earth’s surface; e.g., *microhigh* or *MH* or *microlow* or *ML*.

Microrelief	Code
microhigh	MH
microlow	ML
microslope	MS

**Drainage Pattern**—The interconnected system of drainage channels on the land surface; also called drainage network. (See graphics, p. 3–45.) Can be recorded as a Text Note.

Drainage Pattern	Code
annular	AN
artificial	AR
centripetal	CE
dendritic	DN
deranged	DR
karst	KA
parallel	PA
pinnate	PI
radial	RA

<b>Drainage Pattern</b>	<b>Code</b>
rectangular	RE
thermokarst	TH
trellis	TR

## **WATER STATUS**

---

**DRAINAGE**—An estimate of the natural drainage class (i.e., the prevailing wetness conditions) of a soil; e.g., *somewhat poorly drained* or *SP*.

<b>Drainage Class</b>	<b>Conv. Code</b>
Subaqueous Drainage	SA
Very Poorly Drained	VP
Poorly Drained	PD
Somewhat Poorly Drained	SP
Moderately Well Drained	MW
Well Drained	WD
Somewhat Excessively Drained	SE
Excessively Drained	ED

The following definitions are from the traditional, national criteria for natural soil drainage classes (Soil Survey Division Staff, 1993). Specific regional definitions and criteria exist. (Contact an NRCS State office for specific local criteria.)

**Subaqueous Drainage**—Free water is above the soil surface. The occurrence of internal free water is permanent, and there is a positive water potential at the soil surface for more than 21 hours each day. The soils have a peraquic soil moisture regime.

**Very Poorly Drained**—Water is at or near the soil surface during much of the growing season. Internal free water is *very shallow* and *persistent* or *permanent*. Unless the soil is artificially drained, most mesophytic crops cannot be grown. Commonly, the soil occupies a depression or is level. If rainfall is persistent or high, the soil can be sloping.

**Poorly Drained**—The soil is wet at shallow depths periodically during the growing season or remains wet for long periods. Internal free water is *shallow* or *very shallow* and *common* or *persistent*. Unless the soil is artificially drained, most mesophytic crops cannot be grown. The soil, however, is not

continuously wet directly below plow depth. The water table is commonly the result of a *low* or *very low* saturated hydraulic conductivity class or persistent rainfall or a combination of these factors.

**Somewhat Poorly Drained**—The soil is wet at a shallow depth for significant periods during the growing season. Internal free water is commonly *shallow* or *moderately deep* and *transitory* or *permanent*. Unless the soil is artificially drained, the growth of most mesophytic plants is markedly restricted. The soil commonly has a *low* or *very low* saturated hydraulic conductivity class or a high water table, receives water from lateral flow or persistent rainfall, or is affected by some combination of these factors.

**Moderately Well Drained**—Water is removed from the soil somewhat slowly during some periods of the year. Internal free water commonly is *moderately deep* and may be *transitory* or *permanent*. The soil is wet for only a short time within the rooting depth during the growing season but is wet long enough that most mesophytic crops are affected. The soil commonly has a *moderately low* or lower saturated hydraulic conductivity class within 1 meter of the surface, periodically receives high rainfall, or both.

**Well Drained**—Water is removed from the soil readily but *not* rapidly. Internal free water commonly is *deep* or *very deep*; annual duration is not specified. Water is available to plants in humid regions during much of the growing season. Wetness does not inhibit growth of roots for significant periods during most growing seasons.

**Somewhat Excessively Drained**—Water is removed from the soil rapidly. Internal free water commonly is *very deep* or *very rare*. The soils are commonly coarse textured and have *high* saturated hydraulic conductivity or are *very shallow*.

**Excessively Drained**—Water is removed from the soil very rapidly. Internal free water commonly is *very deep* or *very rare*. The soils are commonly coarse textured and have *very high* saturated hydraulic conductivity or are *very shallow*.

**FLOODING**—Estimate the **Frequency**, **Duration**, and **Months** that flooding is expected; e.g., *rare, brief, Jan.-March*.

**Frequency**—Estimate how often, typically, flooding occurs.

Frequency Class	Code	Criteria: estimated average number of flood events per time span <sup>1</sup>
None	NO	No reasonable chance (e.g., <1 time in 500 years)
Very Rare	VR	≥1 time in 500 years, but <1 time in 100 years
Rare	RA	1 to 5 times in 100 years
Occasional <sup>2</sup>	OC	>5 to 50 times in 100 years
Frequent <sup>2, 3</sup>	FR	>50 times in 100 years
Very Frequent <sup>3, 4</sup>	VF	>50% of all months in year

<sup>1</sup> Flooding Frequency is an estimate of the **current condition**, whether natural or human influenced (such as by dams or artificial levees).

<sup>2</sup> Historically, *Occasional* and *Frequent* classes could be combined and called *Common*; not recommended.

<sup>3</sup> *Very Frequent* class takes precedence over *Frequent*, if applicable.

<sup>4</sup> The *Very Frequent* class is intended for tidal flooding.

**Duration**—Estimate how long an area typically is flooded during a single flood event.

Duration Class	Code		Criteria: estimated average duration per flood event
	Conv.	NASIS	
Extremely Brief	EB	EB	0.1 to < 4 hours
Very Brief	VB	VB	4 to < 48 hours
Brief	BR	B	2 to < 7 days
Long	LO	L	7 to < 30 days
Very Long	VL	VL	≥ 30 days

**Months**—Estimate the beginning and ending month(s) in a year that flooding generally occurs; e.g., *Dec.-Feb*.

**PONDING**—Estimate or monitor the **Frequency**, **Depth**, and **Duration** of standing water. A complete example is: *occasional, 50 cm, brief, Feb.-Apr.*

**Frequency**—Estimate how often, typically, ponding occurs.

Frequency Class	Code	Criteria: estimated, average # of ponding events per time span
None	NO	<1 time in 100 years
Rare	RA	1 to 5 times in 100 years
Occasional	OC	>5 to 50 times in 100 years
Frequent	FR	>50 times in 100 years

**Depth**—Estimate the average, representative depth of ponded water at the observation site and specify units; e.g., *1 ft* or *30 cm*.

**Duration**—Estimate how long, typically, the ponding lasts.

Duration Class	Code		Criteria: estimated, average time per ponding event
	Conv.	NASIS	
Very Brief	VB	VB	<2 days
Brief	BR	B	2 to <7 days
Long	LO	L	7 to <30 days
Very Long	VL	VL	≥30 days

**(SOIL) WATER STATE** (called **Observed Soil Moisture Status** in NASIS.)—Estimate the water state of the soil at the time of observation; e.g., *wet, nonsatiated*. Soil temperature must be above 0 °C. (Does not apply to frozen soil.)



Water State Class	Code		Criteria: tension	Traditional Criteria: tension and field
	Conv.	NASIS		
Dry <sup>1</sup>	D	D	>1500 kPa	>15 bars of tension <sup>2</sup> (=1500 kPa)
Moist <sup>1</sup>	M	M <sup>4</sup>	≤1500 kPa to >1.0 kPa (or >0.5 kPa) <sup>3</sup>	Former Usage: >1/3 to 15 bars of tension (33 to 1500 kPa) (field capacity to wilting point)
Wet	W		≤1.0 kPa (or <0.5 kPa) <sup>3</sup>	0-1/3 bar tension (<33 kPa) (field capacity or wetter)
Wet: Non-satiated <sup>5</sup>	WN		>0.00 and ≤1.0 kPa (or <0.5 kPa) <sup>3</sup>	No Free Water: Water films are visible; sand grains and peds glisten, but no free water is present
Wet: Satiated <sup>5</sup>	WS	W	≤0.00 kPa	Free Water: Free water easily visible

<sup>1</sup> Additional subclasses of water state can be recognized for *Dry* and *Moist* classes, if desired (Soil Survey Division Staff, 1993, p. 91).

<sup>2</sup> Convention assumes 15 bars of tension as the wilting point for most annual agricultural row crops. *Caution:* Various perennials, shrubs, trees, and other native vegetation have a wilting point of as much as 66 bars tension (=6600 kPa) or more.

<sup>3</sup> Use the 1 kPa limit for all textures, *except* those coarser than loamy fine sand (which use 0.5 kPa limit; Soil Survey Division Staff, 1993, p. 90).

<sup>4</sup> NASIS uses the same three class names (Dry, Moist, Wet) but lumps the “wet: nonsatiated” subclass with the Moist class.

<sup>5</sup> **Satiation vs. Saturation:** Satiation implies minor amounts of entrapped air in the smallest pores. True saturation implies no entrapped air. In *Soil Taxonomy*, “Saturation is ... zero or positive pressure in the soil ...” (Soil Survey Staff, 2010). Satiation, for practical purposes, is ≈ saturation. Temporal monitoring of a water table by piezometer or other accepted methods may be needed to verify saturation. Related terms used for classifying soils (i.e., soil taxonomy): *Endosaturation* is saturation in all

layers to >200 cm (80 inches). *Episaturation* requires saturated layers that overlie unsaturated layers within the upper 2 m (80 inches). *Anthric saturation*, a variant of episaturation, is saturation due to management-induced flooding (e.g., for rice or cranberry production).

## LAND COVER

**LAND COVER** (called **EARTH COVER - KIND** in NASIS)—Record the dominant land cover at the site; e.g., *intermixed hardwoods* and *conifers*.

Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
<b>ARTIFICIAL COVER (A)—Nonvegetative cover; due to human activity.</b>			
rural transportation - roads, railroads	RU	urban and built-up - cities, farmsteads, industry	UR
<b>BARREN LAND (B)—&lt;5% vegetative cover naturally or from construction.</b>			
culturally induced - saline seeps, mines, quarries, and oil-waste areas	CI	other barren - salt flats, mudflats, slickspots, badlands	OB
permanent snow or ice	PS	rock	RK
		sand or gravel	SG
<b>CROP COVER (C)—includes entire cropping cycle (land prep, crop, or crop residue) for annual or perennial herbaceous plants.</b>			
close-grown crop - wheat, rice, oats, and rye; small grains	CG	row crop - corn, cotton, soybeans, tomatoes, and other truck crops, tulips	RC
<b>GRASS/HERBACEOUS COVER (G)—&gt;50% grass, grasslike (sedges/rushes), or forb cover, mosses, lichens, ferns; nonwoody.</b>			
hayland - alfalfa, fescue, bromegrass, timothy	HL	rangeland, savanna - 10 to 20% tree cover	RS
marshland - grasses and grasslike plants	ML	rangeland, shrubby - 20 to 50% shrub cover	RH

<b>Kind <sup>1</sup></b>	<b>Code</b>	<b>Kind <sup>1</sup></b>	<b>Code</b>
pastureland, tame - fescues, bromegrass, timothy, lespedeza	PL	rangeland, tundra	RT
rangeland, grassland; <10% trees, <20% shrubs; rangeland used for hayland	RG	other grass and herbaceous cover	OH
<b>SHRUB COVER (S)—&gt;50% shrub or vine canopy cover.</b>			
crop shrubs - filberts, blueberry, ornamental nursery stock	CS	native shrubs - shrub live oak, mesquite, sagebrush, creosote bush; rangeland >50% shrub cover	NS
crop vines - grapes, blackberries, raspberries	CV	other shrub cover	OS
<b>TREE COVER (T)—&gt;25% canopy cover by woody plants, natural or planted.</b>			
conifers - spruce, pine, fir	CO	swamp - trees, shrubs	SW
crop, trees - nuts, fruit, nursery, Christmas trees	CR	tropical - mangrove and royal palms	TR
hardwoods - oak, hickory, elm, aspen	HW	other tree cover	OC
intermixed hardwoods and conifers - oak-pine mix	IM		
<b>WATER (W)—water at the soil surface; includes seasonally frozen water.</b>			

<sup>1</sup> Land Cover Kinds are presented at two levels of detail: Bolded table subheadings are the "NASIS - Level 1" choices (NSSH, Part 622.16; Soil Survey Staff, 2012d). Individual choices under the subheadings are the "NASIS - Level 2" choices.

## **VEGETATION**

**PLANT SYMBOL**—Record the codes (scientific plant name abbreviations) for the major plant species found at the site (NRCS, 2012); e.g., ANGE (*Andropogon gerardii* or *big bluestem*). (**NOTE:** The combination of plant symbol and common name are the primary plant data element in NASIS.)

**PLANT COMMON NAME**—Record the common names of the major plant species found at the site (NRCS, 2012); e.g., *cottonwood*, *big bluestem*. This item may be recorded as a secondary data element to augment the **Plant Symbol**. **CAUTION:** Multiple common names exist for some plants; not all common names for a given plant are in the national PLANTS database.

**PLANT SCIENTIFIC NAME**—Record the scientific plant name along with or in lieu of common names; e.g., *Acer rubrum* (red maple). (**NOTE:** Although used in the past, scientific names of plants [NRCS, 2012] are not presently recorded by the NRCS.) (**NOTE:** NASIS codes for common plant names are derived from the scientific names.)

**VEGETATION COVER**—Estimate the percent of the ground covered by each plant species recorded at the site.

## **PARENT MATERIAL**

Describe the nature of the unconsolidated material (regolith) in which the soil is formed (e.g., till). If the soil is derived directly from the underlying bedrock (e.g., granite), identify the **Parent Material** as either *grus*, *saprolite*, or *residuum* and then record the appropriate **Bedrock - Kind** choice. (**NOTE:** NASIS uses “Component Parent Material Origin” to convey the source from which a Parent Material is derived, predominantly **Bedrock - Kind**.) Multiple parent materials, if present, should be denoted; e.g., *loess*, *over colluvium*, *over residuum*. Use numerical prefixes in the **Horizon** designations to denote different parent materials (lithologic discontinuities); e.g., *A*, *BE*, *2Bt*, *2BC*, *3C*; *Peoria Loess*, or *Calvert Formation*.

**KIND**—e.g., *saprolite*, *loess*, *colluvium*.

Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
<b>EOLIAN DEPOSITS (nonvolcanic)</b>			
eolian deposit	EOD	loess, calcareous	CLO
eolian sands	EOS	loess, noncalcareous	NLO
loess	LOE	parna	PAR
<b>GLACIAL and PERIGLACIAL DEPOSITS</b>			
cryoturbate	CRY	till, ablation	ATI
drift	GDR	till, basal	BTI
glaciofluvial deposit	GFD	till, flow	FTI
glaciolacustrine deposit	GLD	till, lodgment	LTI
glaciomarine deposit	GMD	till, melt-out	MTI

Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
outwash	OTW	till, subglacial	GTI
solifluction deposit	SOD	till, supraglacial	UTI
supraglacial debris-flow	SGF	till, supraglacial meltout	PTI
till	TIL		
<b>IN-PLACE DEPOSITS (nontransported)</b>			
bauxite	BAU	residuum <sup>2</sup>	RES
grus <sup>2</sup>	GRU	saprolite <sup>2</sup>	SAP
<b>MASS MOVEMENT DEPOSITS <sup>3</sup> (See Mass Movement Types table)</b>			
MISCELLANEOUS MASS MOVEMENT DEPOSITS			
colluvium	COL	slump block	SLB
scree	SCR	talus	TAL
MASS MOVEMENT DEPOSIT (Unspecified Landslide)			MMD
COMPLEX LANDSLIDE DEPOSITS			CLD
FALL DEPOSITS			FAD
debris fall deposit	DLD	soil fall deposit (=earth fall)	SFD
rock fall deposit	RFD		
FLOW DEPOSITS			FLD
earthflow deposit	EFD	debris avalanche deposit	DAD
creep deposit	CRP	debris flow deposit	DFD
mudflow deposit	MFD	debris slide deposit	DSD
sand flow deposit	SAD	lahar	LAH
solifluction deposit	SOD	rockfall avalanche deposit	RAD
SLIDE DEPOSITS			SD
debris slide deposit <sup>4</sup>			OSD
Rotational Slide deposit	RLD	Translational Slide deposit	TSD
rotational debris slide deposit	RDD	translational debris slide deposits	TDD
rotational earth slide deposit	RED	translational earth slide deposit	TED

<b>Kind <sup>1</sup></b>	<b>Code</b>	<b>Kind <sup>1</sup></b>	<b>Code</b>
rotational rock slide deposit	RRD	translational rock slide deposit	TRD
		block glide deposit	BGD
SPREAD DEPOSITS (=lateral spread)			LSD
debris spread deposit	DPD	rock spread deposit	RSD
earth spread deposit	EPD		
TOPPLE DEPOSITS			TOD
debris topple deposit	DTD	rock topple deposit	RTD
earth topple (=soil topple)	ETD		
<b>MISCELLANEOUS DEPOSITS</b>			
diamicton	DIM	limonite	LIM
gypsite	GYP		
<b>ORGANIC DEPOSITS <sup>5</sup></b>			
coprogenic materials	COM	organic materials	ORM
diatomaceous earth	DIE	organic, grassy materials	OGM
marl	MAR	organic, herbaceous materials	OHM
marl, coastal	CMA	organic, mossy materials	OMM
marl, freshwater	FWM	organic, woody materials	OWM
<b>VOLCANIC DEPOSITS (unconsolidated; eolian and mass movement)</b>			
ash, volcanic (<2 mm)	ASH	cinders (2-64 mm)	CIN
ash, acidic	ASA	lahar deposit (volcaniclastic mudflow)	LAH
ash, andesitic	ASN	lapilli (2-64 mm, >2.0 sg) <sup>6</sup>	LAP
ash, basaltic	ASB	pumice (<1.0 sg) <sup>6</sup>	PUM
ash, basic	ASC	pyroclastic flow	PYF
ash flow (pyroclastic)	ASF	pyroclastic surge	PYS
bombs, volcanic (>64 mm)	BOM	scoria (>2.0 sg) <sup>6</sup>	SCO

Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
		tephra (all ejecta)	TEP
<b>WATERLAID (or TRANSPORTED) DEPOSITS</b>			
alluvium	ALL	lagoonal deposits	LGD
backswamp deposit	BSD	marine deposit	MAD
beach sand	BES	marl	MAR
coprogenic materials	COM	marl, coastal	CMA
diatomaceous earth	DIE	marl, freshwater	FWM
estuarine deposit	ESD	overbank deposit	OBD
fluviomarine deposit	FMD	pedisediment	PED
greensands	GRS	slope alluvium	SAL
lacustrine deposit	LAD	valley side alluvium	VSA
<b>ANTHROPOGENIC DEPOSITS</b>			
coal extraction mine spoil	CES	metal ore extraction mine spoil	MES
dredge spoils	DGD	mine spoil or earthy fill	MSE
human-transported materials	HTM		

<sup>1</sup> Parent material definitions are found in the "Glossary of Landform and Geologic Terms," NSSH, Part 629 (Soil Survey Staff, 2012b), or the *Glossary of Geology* (Neuendorf et al., 2005).

<sup>2</sup> Use the most precise term possible for the in situ material. Residuum is the most generic term.

<sup>3</sup> Cruden and Varnes, 1996.

<sup>4</sup> *Debris slide* is a more general, encompassing term that may be further subdivided into *rotational debris slide* or *translational debris slide*.

<sup>5</sup> These generic terms refer to the dominant origin of the organic materials or deposits from which the organic soil has formed (i.e., parent material) (Soil Survey Division Staff, 1993). These terms partially overlap with those recognized in soil taxonomy (terms that refer primarily to what the organic material presently is); see the "Diagnostic Horizons or Characteristics" table.

<sup>6</sup> sg=specific gravity=the ratio of a material's density to that of water (weight in air/[weight in air - weight in water]).

## **BEDROCK**

Describe the nature of the continuous hard rock underlying the soil. Specify the **Kind**, **Fracture Interval**, **Hardness**, and **Weathering Class**.

**KIND**—e.g., *limestone*.

<b>Kind <sup>1</sup></b>	<b>Code</b>	<b>Kind <sup>1</sup></b>	<b>Code</b>
<b>IGNEOUS—INTRUSIVE</b>			
anorthosite	ANO	peridotite	PER
diabase	DIA	pyroxenite	PYX
diorite	DIO	quartz-diorite	QZD
gabbro	GAB	quartz-monzonite	QZM
granite	GRA	syenite	SYE
granitoid <sup>2</sup>	GRT	syenodiorite	SYD
granodiorite	GRD	tonalite	TON
monzonite	MON	ultramafic rock <sup>2</sup>	UMU
<b>IGNEOUS—EXTRUSIVE</b>			
aa lava	AAL	pahoehoe lava	PAH
andesite	AND	pillow lava	PIL
basalt	BAS	pumice (flow, coherent)	PUM
block lava	BLL	rhyolite	RHY
dacite	DAC	scoria (coherent mass)	SCO
latite	LAT	tachylite	TAC
obsidian	OBS	trachyte	TRA
<b>IGNEOUS—PYROCLASTIC</b>			
ignimbrite	IGN	tuff, welded	TFW
pyroclastics (consolidated)	PYR	tuff breccia	TBR
pyroclastic flow	PYF	volcanic breccia	VBR
pyroclastic surge	PYS	volcanic breccia, acidic	AVB
tuff	TUF	volcanic breccia, basic	BVB
tuff, acidic	ATU	volcanic sandstone	VST
tuff, basic	BTU		



Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
<b>METAMORPHIC</b>			
amphibolite	AMP	metavolcanics	MVO
gneiss	GNE	mica	MIC
gneiss, biotite	BTG	mica schist	MSH
gneiss, granodiorite	GDG	migmatite	MIG
gneiss, hornblende	HBG	mylonite	MYL
gneiss, migmatitic	MMG	phyllite	PHY
gneiss, muscovite-biotite	MBG	schist	SCH
granofels	GRF	schist, biotite	BTS
granulite	GRL	schist, graphitic	GRS
greenstone	GRE	schist, muscovite	MVS
hornfels	HOR	schist, sericite	SCS
marble	MAR	serpentinite	SER
meta-conglomerate	MCN	siltite	SIT
metaquartzite	MQT	slate	SLA
metasedimentary rocks <sup>2</sup>	MSR	slate, sulfidic	SFS
metasiltstone	MSI	soapstone (talc)	SPS
<b>SEDIMENTARY—CLASTICS</b>			
arenite	ARE	mudstone	MUD
argillite	ARG	ortho-quartzite	OQT
arkose	ARK	porcellanite	POR
breccia, nonvolcanic (angular fragments)	NBR	sandstone	SST
breccia, nonvolcanic, acidic	ANB	sandstone, calcareous	CSS
breccia, nonvolcanic, basic	BNB	shale	SHA
claystone	CST	shale, acid	ASH
conglomerate (rounded fragments)	CON	shale, calcareous	CSH
conglomerate, calcareous	CCN	shale, clayey	YSH
fanglomerate	FCN	siltstone	SIS
glaucinitic sandstone	GLS	siltstone, calcareous	CSI
graywacke	GRY		

Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
<b>SEDIMENTARY—EVAPORITES, ORGANICS, AND PRECIPITATES</b>			
bauxite	BAU	limestone, coral	COR
chalk	CHA	limestone, phosphatic	PLS
lignite	LIG	limonite	LIM
chert	CHE	novaculite	NOV
coal	COA	rock anhydrite	RAN
diatomite	DIA	rock gypsum	GYP
dolomite (dolostone)	DOL	rock halite	RHL
limestone	LST	travertine	TRV
limestone, arenaceous	ALS	tripoli	TRP
limestone, argillaceous	RLS	tufa	TUA
limestone, cherty	CLS		
<b>INTERBEDDED</b> (alternating layers of different sedimentary lithologies)			
limestone-sandstone-shale	LSS	sandstone-shale	SSH
limestone-sandstone	LSA	sandstone-siltstone	SSI
limestone-shale	LSH	shale-siltstone	SHS
limestone-siltstone	LSI		

<sup>1</sup> Definitions for kinds of bedrock are found in the "Glossary of Landform and Geologic Terms," NSSH, Part 629 (Soil Survey Staff, 2012b), or in the *Glossary of Geology* (Neuendorf et al., 2005).

<sup>2</sup> Generic term; use only with regional or reconnaissance surveys (order 3, 4).

**FRACTURE INTERVAL CLASS** (called **Bedrock\_fracture\_interval** in NASIS)—Describe the dominant (average) horizontal spacing between vertical joints (geogenic cracks or seams) in the bedrock layer.

Average Distance Between Fractures	Code
< 10 cm	1
10 to < 45 cm	2

Average Distance Between Fractures	Code
45 to < 100 cm	3
100 to < 200 cm	4
≥ 200 cm	5

**WEATHERING CLASS** (called **Bedrock\_weathering** in NASIS)—The subjective extent to which bedrock has weathered as compared to its presumed nonweathered state. Record in Notes, if used.

Class	Code
Slight	SL
Moderate	MO
Strong	ST

**DEPTH (TO BEDROCK)**—Record the depth (cm) from the ground surface to the contact with coherent (continuous) bedrock.

### **LITHOSTRATIGRAPHIC UNIT(S)**

---

Record the lithostratigraphic unit(s) of the unconsolidated material (regolith) and the bedrock in which the soil is formed or from which it is derived. (This is a text field in NASIS.) For example, *Peoria Loess over pre-Illinoian till over Dakota Formation*. (See discussion, p. 5–12.)

### **EROSION**

---

Estimate the dominant kind and magnitude of accelerated erosion at the site. Specify the **Kind** and **Degree**.

**KIND** (called **erosion\_accelerated\_kind** in NASIS)—

Kind	Code	Criteria <sup>1</sup>
wind	I	Deflation by wind
water:	—	Removal by running water
sheet	S	Even soil loss, no channels
rill	R	Small channels <sup>2</sup>
gully	G	Big channels <sup>3</sup>
tunnel	T	Subsurface voids within soil that enlarge by running water (i.e., piping)

<sup>1</sup> Soil Survey Division Staff, 1993, p. 82.

<sup>2</sup> Small runoff channels that can be obliterated by conventional tillage.

<sup>3</sup> Large runoff channels that cannot be obliterated by conventional tillage.

#### **DEGREE CLASS** (called **erosion\_class** in NASIS)—

<b>Class</b> <sup>1</sup>	<b>Code</b>	<b>Criteria:</b> Estimated % loss of the original, combined A + E horizons or the estimated loss of the upper 20 cm (if original, combined A + E horizons were <20 cm thick). <sup>2</sup>
None	0	0 %
1	1	> 0 up to 25%
2	2	25 up to 75%
3	3	75 up to 100%
4	4	> 75 % and total removal of A

<sup>1</sup> In NASIS, the choices include the preceding word "Class" (e.g., *Class 1*).

<sup>2</sup> Soil Survey Division Staff, 1993, pp. 86–89.

#### **SURFACE FRAGMENTS**

---

Record the amount of surface fragment cover (either as a class or as a numerical percent), as determined by either a "point count" or "line-intercept" method. In NASIS, additional details can be recorded: **Surface Fragment Kind** (called **surface\_frag\_kind** in NASIS), **Surface Fragment Class** (relative quantity), **Mean Distance Between Fragments** (edge to edge), **Shape** (FL-flat or NF-nonflat), **Size**, **Roundness** (use classes and criteria found in "Rock Fragment – Roundness Table"), and **Rock Fragment - Rupture Resistance**.

**KIND**—Document the types of coarse fragments present (same options as "Rock & Other Fragments - Kind").

<b>Kind</b>	<b>Code</b>	<b>Kind</b>	<b>Code</b>
Includes all choices in <b>Bedrock—Kind</b> (except <i>Interbedded</i> ), plus:			
calcrete (caliche) <sup>1</sup>	CA	metamorphic rocks <sup>2</sup>	MMR
carbonate concretions	CAC	mixed rocks <sup>3</sup>	MXR

<b>Kind</b>	<b>Code</b>	<b>Kind</b>	<b>Code</b>
carbonate nodules	CAN	ortstein fragments	ORF
carbonate rocks <sup>2</sup>	CAR	petrocalcic fragments	PEF
charcoal	CH	petroferric fragments	TCF
cinders	CI	petrogypsic fragments	PGF
durinodes	DNN	plinthite nodules	PLN
duripan fragments	DUF	quartz	QUA
foliated metamorphic rocks <sup>2</sup>	FMR	quartzite	QZT
gibbsite concretions	GBC	scoria	SCO
gibbsite nodules	GBN	sedimentary rocks <sup>2</sup>	SED
igneous rocks <sup>2</sup>	IGR	shell fragments	SHF
iron-manganese concretions	FMC	silica concretions	SIC
iron-manganese nodules	FMN	volcanic bombs	VB
ironstone nodules	FSN	volcanic rocks <sup>2</sup>	VOL
lapilli	LA	wood	WO

<sup>1</sup> Fragments strongly cemented by carbonate; may include fragments derived from petrocalcic horizons.

<sup>2</sup> Generic rock names may be appropriate for identifying fragments (e.g., a cobble) but are too general and should *not* be used to name Bedrock—Kind.

<sup>3</sup> Numerous unspecified fragment lithologies are present, as in till or alluvium; not for use with residuum.

<b>Surface Fragment Class <sup>1</sup></b>	<b>Code</b>		<b>Criteria: percentage of surface covered</b>
	<b>Conv. <sup>2</sup></b>	<b>NASIS</b>	
Stony or Bouldery	Class 1	%	0.01 to <0.1
Very Stony, or Very Bouldery	Class 2	%	0.1 to <3
Extremely Stony or Extremely Bouldery	Class 3	%	3 to <15
Rubblly	Class 4	%	15 to <50
Very Rubblly	Class 5	%	≥50 <sup>3</sup>

- <sup>1</sup> This is also used to record large wood fragments (e.g., tree trunks) on organic soils, if the fragments are a management concern and appear to be relatively permanent.
- <sup>2</sup> Historically called *Surface Stoniness* classes (now *Surface Fragment* classes). Use as a map unit phase modifier is restricted to stone-sized fragments or larger (>250 mm; Soil Survey Staff, 1951).
- <sup>3</sup> If the percentage of surface fragments is >80%, the fragments are considered to be a distinct, separate horizon (NSSH, Amendment #4; 1998).

## ***DIAGNOSTIC HORIZONS or CHARACTERISTICS***

Identify the **Kind** and **Upper** and **Lower Depths** of occurrence of soil taxonomic diagnostic horizons and characteristics (Soil Survey Staff, 2010); e.g., *mollic epipedon*; 0-45 cm. Multiple features per horizon can be recorded. (Called **Pedon Diagnostic Features** in NASIS.) Record **Kind**, **Thickness**, **Representative Value (RV)**. **High Value** and **Low Value** can also be recorded.

**KIND**—(see definitions in current *Keys to Soil Taxonomy*)

<b>Kind</b>	<b>Code</b>	<b>Kind</b>	<b>Code</b>
<b><i>EPIPEDONS (Diagnostic Surface Horizons)</i></b>			
anthropic	AN	mollic	MO
folic	FO	ochric	OC
histic	HI	plaggen	PL
melanic	ME	umbric	UM
<b><i>DIAGNOSTIC SUBSURFACE HORIZONS</i></b>			
agric	AG	natric	NA
albic	AL	ortstein	OR
argillic	AR	oxic	OX
calcic	CA	petrocalcic	PE
cambic	CM	petrogypsic	PG
duripan	DU	placic	PA
fragipan	FR	salic	SA
glossic	GL	sombric	SO
gypsic	GY	spodic	SP
kandic	KA	sulfuric	SU

Kind	Code	Kind	Code
<b>DIAGNOSTIC CHARACTERISTICS—MINERAL SOILS</b>			
abrupt textural change	AC	gypsum accumulations	GA
albic materials	AM	lamella/lamellae	LA
albic materials, interfingering of	AI	lithic contact <sup>1</sup>	LC
andic soil properties	AP	lithologic discontinuity	LD
anhydrous conditions	AH	paralithic contact <sup>1</sup>	PC
aquic conditions <sup>1</sup>	AQ	paralithic materials <sup>1</sup>	PM
artifacts	ART	permafrost <sup>1</sup>	PF
cryoturbation <sup>1</sup>	CR	petroferric contact	TC
densic contact <sup>1</sup>	DC	plinthite	PI
densic materials <sup>1</sup>	DM	resistant minerals	RM
durinodes	DN	slickensides	SS
fragic soil properties	FP	spodic materials	SPM
free carbonates	FC	sulfidic materials <sup>1</sup>	SM
gelic materials <sup>1</sup>	GM	weatherable minerals	WM
glacic layer <sup>1</sup>	GL		
<b>DIAGNOSTIC CHARACTERISTICS—ORGANIC SOILS</b> (also see <sup>1s</sup> above)			
fibric soil materials	FM	limnic materials:	LM
hemic soil materials	HM	coprogenous earth	CO
humilluvic material	UM	diatomaceous earth	DI
sapric soil materials	RM	marl	MA
<b>MISCELLANEOUS HORIZON FEATURES/CONDITIONS</b>			
anthric saturation <sup>1</sup>	AS	redox depletions with chroma 2 or less <sup>1</sup>	RD
endosaturation <sup>1</sup>	EN	reduced matrix <sup>1</sup>	RX
episaturation <sup>1</sup>	ED	salt accumulations	ST
fibers	FI	secondary carbonates <sub>2</sub>	SC
n-value >0.7	NV	strongly contrasting particle-size class	SR
redox concentrations <sup>1</sup>	RC	volcanic glass	VG

<sup>1</sup> Diagnostic properties, materials, or conditions that can occur in either mineral or organic soils.

<sup>2</sup> "Secondary carbonates" replaces "soft, powdery lime." **NOTE:** Gilgai is no longer a diagnostic feature in soil taxonomy.

**DEPTH**—Document the zone of occurrence for a diagnostic horizon or property, as observed, by recording the **Top Depth** and **Bottom Depth** and specifying units; e.g., 22-39 cm.

**SOIL TAXONOMY CLASSIFICATION**—After completely describing the soil, classify the pedon as thoroughly as possible (to the lowest level). See most recent version of *Soil Taxonomy* and *Keys to Soil Taxonomy* for complete choice lists; e.g., *fine, mixed, active, mesic Typic Haplohumults*.

**PARTICLE-SIZE CONTROL SECTION**—Record the **Upper** and **Lower Depth** of the zone used as the basis for identifying the particle-size control section; e.g., 30-80 cm (used to classify in soil taxonomy).

## **RESTRICTION**

---

**RESTRICTION - KIND**—Identify any root-limiting/restrictive layers within the soil profile. Also record the **Upper** and **Lower Depth** of occurrence.

<b>Kind</b>	<b>Code</b>	<b>Kind</b>	<b>Code</b>
abrupt textural change	AC	paralithic bedrock	BPL
cemented horizon	CH	permafrost	PF
densic material	DM	petrocalcic	PE
densic bedrock	BD	petroferric	TC
duripan	DU	petrogypsic	PG
fragipan	FR	placic	PA
human-manufactured materials	HF	plinthite	PI
lithic bedrock	BL	salic	SA
natric	NA	strongly contrasting textural stratification	SR
ortstein	OR	sulfuric	SU

**RESTRICTION - HARDNESS**—Estimate the hardness of a root-restrictive layer. (Use *Rupture Resistance - Cementation Classes*.)



## REFERENCES

---

Bureau of Chemistry and Soils. 1937. Soil survey manual. USDA Misc. Publ. 274. Washington, DC.

Cruden, D.M., and D.J. Varnes. 1996. Landslide types and processes. *In* A.K. Turner and R.L. Schuster (ed.) *Landslides: Investigation and mitigation*. Spec. Rep. 247, Transportation Research Board, National Research Council. National Academy Press, Washington, DC.

International Organization for Standards. 2012a (or current date). ([http://www.iso.org/iso/country\\_codes/iso\\_3166\\_code\\_lists.htm](http://www.iso.org/iso/country_codes/iso_3166_code_lists.htm)) Note: Free versions at [http://en.wikipedia.org/wiki/ISO\\_3166-1](http://en.wikipedia.org/wiki/ISO_3166-1) for country codes.

International Organization for Standards. 2012b (or current date). ([http://www.iso.org/iso/country\\_codes/iso\\_3166-2\\_database.htm](http://www.iso.org/iso/country_codes/iso_3166-2_database.htm)) Note: Free versions at [http://en.wikipedia.org/wiki/ISO\\_3166-1\\_alpha-2](http://en.wikipedia.org/wiki/ISO_3166-1_alpha-2) for administrative subdivisions within country.

National Institute of Standards and Technology. 1990. Counties and equivalent entities of the United States, its possessions and associated areas. U.S. Dep. Commerce, Federal Information Processing Standards Publ. (FIPS PUB 6-4).

Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. USDA, Agric. Handb. 296. U.S. Gov. Print. Office, Washington, DC.

Natural Resources Conservation Service. 2012 (or most current date) [Online]. Complete PLANTS checklist. (The national PLANTS database.) USDA, National Plant Data Center, Baton Rouge, LA (<http://plants.usda.gov>).

Neuendorf, K., J.P. Mehl, and J.A. Jackson. 2005. *Glossary of geology*, 5th ed. Am. Geol. Inst., Alexandria, VA.

Public Building Service. Sept. 1996. *Worldwide geographic location codes*. U.S. Gen. Serv. Admin., Washington, DC.

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and W.D. Broderson. 1998. *Field book for describing and sampling soils* (ver. 1.1). USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and W.D. Broderson (ed.) 2002. *Field book for describing and sampling soils* (ver. 2.0). USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Soil Conservation Service. 1981. *Land resource regions and major land resource areas of the United States*. USDA, Agric. Handb. 296. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Division Staff. 1993. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1951. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1962. Identification and nomenclature of soil horizons. Supplement to Agric. Handb. 18, Soil Survey Manual (replacing pages 173–188). USDA, SCS. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1983. National soil survey handbook, part 603, p. 45. USDA, SCS. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 2010. Keys to soil taxonomy, 11th ed. USDA, NRCS. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 2011. Soil survey laboratory information manual. Soil Surv. Invest. Rep. 45, ver. 2.0. USDA, NRCS, National Soil Survey Center, Lincoln, NE ([ftp://ftp-fc.sc.egov.usda.gov/NSSC/Lab\\_Info\\_Manual/SSIR\\_45.pdf](ftp://ftp-fc.sc.egov.usda.gov/NSSC/Lab_Info_Manual/SSIR_45.pdf)).

Soil Survey Staff. 2012a. Data dictionary. *In* National soil information system (NASIS), release 6.2 [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 2012b. Glossary of landform and geologic terms. Part 629, National soil survey handbook (NSSH) [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 2012c. National soil information system (NASIS), release 6.2 [Online]. USDA, NRCS, Lincoln, NE.

Soil Survey Staff. 2012d. National soil survey handbook (NSSH) [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE (<http://soils.usda.gov/technical/handbook/>).

Wysocki, D.A., P.J. Schoeneberger, D. Hirmas, and H.E. LaGarry. 2012. Geomorphology of soil landscapes. *In* P.M. Huang et al. (ed.) Handbook of soil science: Properties and processes, 2nd ed. CRC Press, Taylor and Francis Group, LLC, Boca Raton, FL. ISBN: 978-1-4398-0305-9.

Wysocki, D.A., P.J. Schoeneberger, and H.E. LaGarry. 2000. Geomorphology of soil landscapes. *In* M.E. Sumner (ed.) Handbook of soil science. CRC Press, LLC, Boca Raton, FL. ISBN: 0-8493-3136-6.

# PROFILE/PEDON DESCRIPTION

D.A. Wysocki, P.J. Schoeneberger, J.V. Chiaretti, and E.C. Benham, NRCS,  
Lincoln, NE

## OBSERVATION METHOD

For each layer, record the observation method by which the primary observations are made. (Common sampling devices are included in the "Soil Sampling" section.) Describe **Kind** and **Relative Size**; e.g., *bucket auger, 3"*; *trench, 2 x 4 m*.

**KIND** (called **Observation\_Method** in NASIS)—

Kind <sup>1</sup>	Code	Criteria
<b>"Disturbed" Samples</b>		
bucket auger	BA	Open, closed, sand, mud buckets (5-12 cm diam.)
dutch or mud auger	DA	An open, strap-sided bucket (5-10 cm diam.) with a sharpened outer edge and a screw tip with a partial twist
screw auger	SA	External thread hand augers, power (flight) auger (2-30 cm diam.)
<b>"Undisturbed" Samples</b>		
Macaulay sampler	MC	A half-cylinder, "gouge" sampler with a hinged door that's pushed in and partially rotated to obtain a sample of soft sediments (e.g., organics)
push tube	PT	Handheld or hydraulic, hollow stem (2-10 cm diam.)
shovel "slice" <sup>2</sup>	SS	Undisturbed block extracted with a shovel (sharpshooter: 20 x 40 cm)
vibracore tube	VT	A hollow tube (4-8 cm diam.) vibrated into wet sand, silt, or organics
<b>WALL/FLOOR—"Undisturbed" Area or Exposure</b>		
small pit	SP	Hand or machine dug (<1 m x 2 m)
trench	TR	Hand or machine dug (>1 m x 2 m)
beveled cut	BC	Roadcuts graded to <60% slope
cut	CU	Roadcut, streambank, medium borrow pit wall >60% slope (>4 m, <33 m)
large open pit or quarry	LP	Large borrow pit or quarry with large or irregular banks (>33 m)

Kind <sup>1</sup>	Code	Criteria
<b>Other Observations</b>		
dive	DV	A visual onsite assessment performed underwater
video	VO	Electronically recorded photo or sequential digital images of a subaqueous setting/site

<sup>1</sup> Refer to **Examples of Common Soil-Sampling Equipment** (p. 8–5) and **Bucket Auger Types** (p. 8–6) for examples of field equipment.

<sup>2</sup> Field method used for hydric soil investigations.

**RELATIVE SIZE** (of exposure) (called **Relative\_Exposure\_UOM** in NASIS)—Record the approximate size (scale) of the exposure observed. Use cm for “Drill Cores” and m for “Wall/Floor” observations; e.g., *bucket auger, 3 cm; trench wall, 3 m.* (**NOTE:** Common size range for each method is indicated in the “Criteria” column of the “Observation Method – Kind” table. These dimensions are approximate; not intended to be precise.)

Relative Size of Exposure Observed	Code	Criteria
centimeters	cm	
meters	m	

## **HORIZON AND LAYER DESIGNATIONS**

---

Use capital letters to identify master horizons; e.g., *A, B*. Use suffixes (lowercase letters) to denote additional horizon characteristics or features; e.g., *Ap, Btk*. (For more detailed criteria, see the “Soil Taxonomy” section [p. 4–1]; for complete definitions, see *Keys to Soil Taxonomy* [Soil Survey Staff, 2010].) Label a horizon (assign horizon designation) only *after* all morphology is recorded.

### **MASTER AND TRANSITIONAL HORIZONS AND LAYERS <sup>1</sup>—**

Identify the master horizons of the soil profile.

Horizon	Criteria (expanded details listed in “Soil Taxonomy” section)
<b>O</b>	Organic soil materials (not limnic).
<b>A</b>	Mineral; organic matter (humus) accumulation, loss of Fe, Al, clay.

<b>Horizon</b>	<b>Criteria</b> (expanded details listed in "Soil Taxonomy" section)
<b>AB</b> or <b>AE</b> or <b>AC</b>	Dominantly A horizon characteristics but also contains some B, E, or C horizon attributes.
<b>A/B</b> or <b>A/E</b> or <b>A/C</b>	Discrete, intermingled bodies of A and B, E, or C material; majority is A material.
<b>E</b>	Mineral; some loss of Fe, Al, clay, or organic matter.
<b>EA</b> or <b>EB</b> or <b>EC</b>	Dominantly E horizon characteristics but also contains some A, B, or C horizon attributes.
<b>E/A</b> or <b>E/B</b>	Discrete, intermingled bodies of E and A or B horizon material; majority of horizon is E material.
<b>E and Bt</b> <b>B and E</b>	Thin, heavier textured lamellae (Bt) within a dominantly E horizon (or thin E within dominantly B horizon).
<b>BA</b> or <b>BE</b> or <b>BC</b>	Dominantly B characteristics but contains A, E, or C horizon attributes.
<b>B/A</b> or <b>B/E</b> or <b>B/C</b>	Discrete, intermingled bodies of B and A, E, or C material; majority of horizon is B material.
<b>B</b>	Subsurface accumulation of clay, Fe, Al, Si, humus, CaCO <sub>3</sub> , CaSO <sub>4</sub> ; or loss of CaCO <sub>3</sub> ; or accumulation of sesquioxides; or subsurface soil structure.
<b>CB</b> or <b>CA</b>	Dominantly C horizon characteristics but also contains attributes of the B or A horizon.
<b>C/B</b> or <b>C/A</b>	Discrete, intermingled bodies of C and B or A material; majority of horizon is C material.
<b>C</b>	Little or no pedogenic alteration, unconsolidated earthy material, soft bedrock.
<b>L</b>	Limnic soil materials.
<b>W</b>	A layer of liquid water (W) or permanently frozen water (Wf) within or beneath the soil (excludes water/ice above soil).
<b>M</b>	Root-limiting subsoil layers of human-manufactured materials.
<b>R</b>	Bedrock, strongly cemented to indurated.

<sup>1</sup> See "Soil Taxonomy" (p. 4–6) for older horizon nomenclature.

<sup>2</sup> Soil Survey Staff, 2010.

**HORIZON SUFFIXES**—Historically referred to as “Horizon Subscripts,” “Subordinate Distinctions,” <sup>1</sup> “Horizon\_Designation\_Suffix” in NASIS, and as “Suffix Symbols” in soil taxonomy <sup>2</sup>. (Historical designations and conversions are shown in the “Soil Taxonomy” section.)

<b>Horizon Suffix</b>	<b>Criteria <sup>2</sup></b> (expanded details listed in “Soil Taxonomy” section)
<b>a</b>	Highly decomposed organic matter (used only with O)
<b>aa</b> <sup>3</sup>	(proposed) Accumulation of anhydrite (CaSO <sub>4</sub> )
<b>b</b>	Buried genetic horizon (not used with C horizons)
<b>c</b>	Concretions or nodules
<b>co</b>	Coprogenous earth (used only with L)
<b>d</b>	Densic layer (physically root restrictive)
<b>di</b>	Diatomaceous earth (used only with L)
<b>e</b>	Moderately decomposed organic matter (used only with O)
<b>f</b>	Permanently frozen soil or ice (permafrost); continuous subsurface ice; not seasonal ice
<b>ff</b>	Permanently frozen soil (“Dry” permafrost); no continuous ice; not seasonal ice
<b>g</b>	Strong gley
<b>h</b>	Illuvial organic matter accumulation
<b>i</b>	Slightly decomposed organic matter (used only with O)
<b>j</b>	Jarosite accumulation
<b>jj</b>	Evidence of cryoturbation
<b>k</b>	Pedogenic CaCO <sub>3</sub> accumulation (<50% by vol.)
<b>kk</b>	Major pedogenic CaCO <sub>3</sub> accumulation (>50% by vol.)
<b>m</b>	Continuous cementation (pedogenic)
<b>ma</b>	Marl (used only with L)
<b>n</b>	Pedogenic, exchangeable sodium accumulation
<b>o</b>	Residual sesquioxide accumulation (pedogenic)
<b>p</b>	Plow layer or other artificial disturbance
<b>q</b>	Secondary (pedogenic) silica accumulation
<b>r</b>	Weathered or soft bedrock
<b>s</b>	Illuvial sesquioxide and organic matter accumulation
<b>se</b>	Presence of sulfides (in mineral or organic horizons)
<b>ss</b>	Slickensides

<b>Horizon Suffix</b>	<b>Criteria <sup>2</sup></b> (expanded details listed in "Soil Taxonomy" section)
<b>t</b>	Illuvial accumulation of silicate clay
<b>u</b>	Presence of human-manufactured materials (artifacts)
<b>v</b>	Plinthite
<b>w</b>	Weak color or structure within B (used only with B)
<b>x</b>	Fragipan characteristics
<b>y</b>	Accumulation of gypsum
<b>yy</b>	Dominance of gypsum ( $\approx \geq 50\%$ by vol.)
<b>z</b>	Pedogenic accumulation of salt more soluble than gypsum

<sup>1</sup> Soil Survey Division Staff, 1993.

<sup>2</sup> Soil Survey Staff, 2010.

<sup>3</sup> Personal communication with Soil Survey Standards Staff, 2012.

#### **OTHER HORIZON MODIFIERS—**

**Numerical Prefixes** (2, 3, etc.)—Used to denote lithologic discontinuities. By convention, 1 is understood but is *not* shown; e.g., *A, E, Bt1, 2Bt2, 2BC, 3C1, 3C2*. (**NOTE:** Discontinuities have important implications for site history, internal water flow, and soil interpretations [see additional discussion under "Subaqueous Soils," p. 2–103]).

**Numerical Suffixes**—Used to denote subdivisions within a master horizon; e.g., *A1, A2, E, Bt1, Bt2, Bt3, Bs1, Bs2*.

**The Prime (´)** (Called `horz_desgn_master_prime` in NASIS)—Used to indicate the recurrence of identical horizon descriptor(s) in a profile or pedon; e.g., *A, E, Bt, E´ Btx, C*. The prime does not indicate either buried horizons (which are denoted by a lowercase "b"; e.g., *Btb*) or lithologic discontinuities (denoted by numerical prefixes). In NASIS, up to five primes can be used to denote subsequent occurrences of horizon descriptors in a pedon; e.g., *A, E, Bt, E´, Btx, E´´, Cd*.

**The Caret (^) symbol**—Used as a prefix to master horizons to indicate human-transported material; e.g., *^A, ^Bw, C*. (The caret symbol can be applied to all master horizon combinations *except* B and E, B/E, E and B, E/B, EC, L, M, R, or W.)

**HORIZON DEPTH**—Record the depths of both the upper and lower boundary for each horizon; specify units (centimeters preferred); e.g., 15-24 cm. Begin (zero datum) at the ground surface <sup>1</sup>, which is not necessarily the mineral surface. (**NOTE:** Prior to 1993, the zero datum was at the top of the mineral surface, except for thick organic layers, such as peat or muck. Organic horizons were recorded as above and mineral horizons recorded as below, relative to the mineral surface.)

Example: Zero Datum for the same horizons

At Present: Oe 0 - 5 cm, A 5 - 15 cm, E 15 - 24 cm

Before 1993: Oe 5 - 0 cm, A 0 - 10 cm, E 10 - 19 cm

<sup>1</sup> Conventionally, the "soil surface" is considered to be the top boundary of the first layer that can support plant/root growth. This equates to:

- a) (for bare mineral soil) the air/fine earth interface;
- b) (for vegetated mineral soil) the upper boundary of the first layer that can support root growth;
- c) (for organic mantles) the same as b) but *excludes* freshly fallen plant litter and includes litter that has compacted and begun to decompose; e.g., Oi horizon;
- d) (for submerged soil) the same as b) but refers to the water/soil contact and extends out from the shore to the limit of rooted plants;
- e) (for rock mulches; e.g., desert pavement, scree) the same as a) unless the areal percentage of surface rock coverage is greater than 80%, the top of the soil is the mean height of the top of the rocks.

**HORIZON THICKNESS**—Record the average thickness and range in thickness of horizon; e.g., 15 cm (12-21 cm). **NOTE:** Used primarily for irregular soil horizons/layers.

**HORIZON BOUNDARY**—Record **Distinctness** and **Topography** of horizon boundaries. (In NASIS, Distinctness is called Boundary Distinctness). Distinctness is the vertical distance through which the bottom of one horizon grades (transitions) into another. Topography is the lateral undulation and continuity of the boundary between horizons. A complete example is *clear, wavy, or C,W*.

**Distinctness**—The vertical distance (thickness) over which a horizon transitions to the top of the next.


Distinctness Class	Code	Criteria: transitional zone thickness
Very Abrupt	V	< 0.5 cm
Abrupt	A	0.5 to < 2 cm



Distinctness Class	Code	Criteria: transitional zone thickness
Clear	C	2 to < 5 cm
Gradual	G	5 to < 15 cm
Diffuse	D	≥ 15 cm


**Topography**—Cross-sectional shape of the contact between horizons.

Topography	Code	Criteria
Smooth	S	Planar with few or no irregularities
Wavy	W	Width of undulation is > than depth
Irregular	I	Depth of undulation is > than width
Broken	B	Discontinuous horizons; discrete but intermingled, or irregular pockets



## SOIL COLOR

**DECISION FLOWCHART FOR DESCRIBING SOIL COLORS**—Use the following chart to decide how and with which data elements the color patterns of a soil or soil feature should be described.



**NOTE:** Reduced matrix color is described as a *matrix color* and in the associated "(Soil Color) - Location or Condition Described Table."

**(SOIL) MATRIX COLOR**—Record the **Color(s)**, **Moisture State**, and **Location or Condition**.

**(Soil) Matrix Color - (Soil) Color**—Identify the soil matrix color(s) with Munsell® notation (**Hue, Value, Chroma**); e.g., 10YR 3/2. For neutral colors, chroma is zero but not shown; e.g., N 4/. For other gley colors, use appropriate notation (see Munsell® gley pages; e.g., 5GY 6/1). For narrative descriptions (soil survey reports, Official Soil Series Descriptions), both the verbal name and the Munsell® notation are given; e.g., *dark brown*, 10YR 3/3.

**(Soil) Matrix Color - Moisture State**—Record the general moisture condition of the soil described; e.g., *moist*. (Not to be confused with Soil Water State.)

Moisture State	Code
Dry	D
Moist	M

**(Soil) Matrix Color - Location or Condition**—Record pertinent circumstances of the color described (called **color\_physical\_state** in NASIS) in Notes column.

Color Location or Condition	Code
<b>COLOR LOCATION</b>	
interior ( <i>within ped</i> )	IN
exterior ( <i>ped surface</i> )	EX
<b>COLOR, MECHANICAL CONDITION</b>	
broken face	BF
crushed	CR
rubbed ( <i>used only with organic matter</i> )	RU
<b>COLOR, REDOXIMORPHIC CONDITION</b>	
oxidized <sup>1</sup>	OX
reduced <sup>2</sup>	RE
<b>COLOR, INTRICATE MULTICOLORED PATTERN</b>	
variegated <sup>3</sup>	VA

<sup>1</sup> Soil that is reduced *in situ* but has been extracted and exposed to the atmosphere (air) and has oxidized (changed color). A mineral example is vivianite. **NOTE:** Not used for soil that is normally oxidized in place. For indicators of reduction, see **Redoximorphic Features**.

<sup>2</sup> Color determined immediately after extraction from a reduced environment and prior to oxidation; e.g., *FeS*. Also used to record **Reduced Matrix**.

<sup>3</sup> Color pattern is too intricate (banded or patchy) with numerous diverse colors to credibly identify dominant matrix colors (e.g., foliated felsic crystalline saprolite).

## **REDOXIMORPHIC FEATURES—RMFs (DISCUSSION)**

Redoximorphic features (RMFs) are color patterns in a soil caused by loss (depletion) or gain (concentration) of pigment compared to the matrix color, formed by oxidation/reduction of Fe and/or Mn coupled with their removal, translocation, or accrual; or a soil matrix color controlled by the presence of  $Fe^{+2}$ . The composition and process of formation for a soil color or color pattern must be known or inferred before it can be described as an RMF. Because of this inference, RMFs are described separately from mottles, other concentrations (e.g., *salts*), or compositional features (e.g., *clay films*). RMFs generally occur in one or more of these settings:

- a. In the soil matrix, unrelated to surfaces of peds or pores.
- b. On or beneath the surfaces of peds.
- c. As filled pores, as linings of pores, or beneath the surfaces of pores.

### **RMFs include the following:**

1. **Redox Concentrations**—Localized zones of enhanced pigmentation due to an accrual of, or a phase change in, the Fe-Mn minerals; or physical accumulations of Fe-Mn minerals.  
**NOTE:** Iron concentrations may be either  $Fe^{+3}$  or  $Fe^{+2}$ . Types of redox concentrations are:
  - a. **Masses**—Noncemented bodies of enhanced pigmentation that have a redder or blacker color than the adjacent matrix.
  - b. **Nodules or Concretions**—Cemented bodies of Fe-Mn oxides.
2. **Redox Depletions**—Localized zones of “decreased” pigmentation that are grayer, lighter, or less red than the adjacent matrix. Redox depletions include, but are not limited to, what were previously called “low-chroma mottles” (chroma  $\leq 2$ ). Redox depletions of chroma  $\leq 2$  formed through reduction and oxidation processes are strong field indicators of saturation. Types of redox depletions are:
  - a. **Iron Depletions**—Localized zones that have one or more of the following: a yellower, greener, or bluer hue; a higher value; or a lower chroma than the matrix color. Color value is

normally  $\geq 4$ . Loss of pigmentation results from the loss of Fe and/or Mn. Clay content equals that in the matrix.

- b. **Clay Depletions**—Localized zones that have either a yellower, greener, or bluer hue, a higher value, or a lower chroma than the matrix color. Color value is normally  $\geq 4$ . Loss of pigmentation results from a loss of Fe and/or Mn and clay. Silt coats or skeletalans commonly form as depletions but can be nonredox concentrations if deposited as flow material in pores or along faces of peds.
3. **Reduced Matrix**—A soil horizon that has an in situ matrix chroma  $\leq 2$  due to the presence of  $\text{Fe}^{+2}$ . Color becomes redder or brighter (oxidizes) when the sample is exposed to air. The color change usually occurs within 30 minutes. A 0.2% solution of  $\alpha, \alpha'$ -dipyridyl dissolved in 1N ammonium acetate ( $\text{NH}_4\text{OAc}$ ) pH 7 can verify the presence of  $\text{Fe}^{+2}$  in the field (Childs, 1981).

**NOTE:** RMF alters the traditional sequence for describing soil color (see the "Decision Flowchart for Describing Colors for Soil Matrix and Soil Features"). RMFs are described separately from other color variations or concentrations. Mottles (color variations *not* due to loss or accrual of Fe-Mn oxides; e.g., variegated weathered rock) are still described under **Soil Color**. A reduced matrix is recorded as an RMF and as "reduced" in **Soil Color - Location** or **Condition Described**.

## REDOXIMORPHIC FEATURES

Record **Kind, Quantity** (percent of area covered), **Size, Contrast, Color, Moisture State, Shape, Location, Hardness,** and **Boundary**. A complete example is: *common, medium, prominent, black iron-manganese nodules, moist, spherical, in the matrix, weakly cemented, sharp or c, 2, p, 5YR 2.5/1, FMM, M, S, MAT, w, s*. At present, relict RMFs, as supported by geomorphic setting, water table data, etc., are recorded as "relict RMFs" (include horizons and depths) under **Miscellaneous Field Notes**.

### REDOXIMORPHIC FEATURES - KIND—

Kind	Code	Kind	Code
<b>REDUCED MATRIX (chroma <math>\leq 2</math> primarily from Fe<sup>+2</sup>)</b>			
reduced matrix	RMX		
<b>REDOX DEPLETIONS (loss of pigment or material)</b>			
clay depletions	CLD	iron depletions	FED
<b>REDOX CONCENTRATIONS (accumulated pigment, material)</b>			
<b>Masses <sup>1</sup> (noncemented)</b>			
iron (Fe <sup>+2</sup> ) <sup>2</sup>	F2M	jarosite	JAM
iron (Fe <sup>+3</sup> ) <sup>3, 4, 5</sup>	F3M	manganese <sup>4, 5</sup>	MNM
iron-manganese <sup>3, 4, 5</sup>	FMM		
<b>Nodules <sup>1</sup> (cemented; no layers, crystals not visible at 10X)</b>			
ironstone	FSN	jarosite	JAN
iron-manganese <sup>4</sup>	FMN	plinthite	PLN
<b>Concretions <sup>1</sup> (cemented; distinct layers, crystals not visible)</b>			
iron-manganese <sup>4</sup>			FMC
<b>Surface Coats/Films or Hypocoats</b>			
manganese (mangans: flat black, very thin, exterior films)			MNF
ferriargillans (Fe <sup>+3</sup> stained clay film)			FEF

<sup>1</sup> See discussion under **Concentrations** for definitions.

<sup>2</sup> A concentration of reduced iron Fe<sup>+2</sup>; e.g., FeS.

<sup>3</sup> A concentration of oxidized iron Fe<sup>+3</sup>; e.g., hematite (formerly described as *reddish mottles*).

<sup>4</sup> Iron and Mn commonly occur in combination, and field identification of distinct phases is difficult. Use *Mn masses* only for those that are at least *Slightly Effervescent* with H<sub>2</sub>O<sub>2</sub>. Describe nodules and concretions as *iron-manganese* unless colors are unambiguous.

<sup>5</sup> Suggested color guidelines for field description of Fe vs. Mn masses:


Color of RMF or Concentration		Dominant Composition
Value	Chroma	
≤2	≤2	Mn
>2 and ≤4	>2 and ≤4	Fe and Mn
>4	>4	Fe

**REDOXIMORPHIC FEATURES - QUANTITY (Percent of Area Covered)**—See graphics for **% of Area Covered** (2, 20%) beginning on p. 7–1.

Class	Code		Criteria: percent of surface area covered
	Conv.	NASIS	
Few	f	#	< 2
Common	c	#	2 to < 20
Many	m	#	≥ 20

**REDOXIMORPHIC FEATURES - SIZE**—See size class graphic on next page.

Size Class	Code	Criteria
Fine	1	< 2 mm
Medium	2	2 to < 5 mm
Coarse	3	5 to < 20 mm
Very Coarse	4	20 to < 76 mm
Extremely Coarse	5	≥ 76 mm





**REDOXIMORPHIC FEATURES - CONTRAST**—Record the color difference between the RMF and the dominant matrix color; e.g., *Prominent* or *p*. Use this table or the following chart to express the difference. (Also used for **Concentrations** and **Mottles**.)

<b>Contrast Class</b>	<b>Code</b>	<b>Difference in Color Between Matrix and RMF (<math>\Delta</math> means "difference between")</b>		
Faint <sup>1</sup>	F	Hue (h)	Value (v)	<b>Chroma (c)</b>
		$\Delta h = 0;$	$\Delta v \leq 2$	and $\Delta c \leq 1$
		$\Delta h = 1;$	$\Delta v \leq 1$	and $\Delta c \leq 1$
		$\Delta h = 2;$	$\Delta v = 0$	and $\Delta c = 0$
Distinct <sup>1</sup>	D	$\Delta h = 0;$	$\Delta v \leq 2$	and $\Delta c > 1$ to $< 4$
			or $\Delta v > 2$ to $< 4$	and $\Delta c < 4$
		$\Delta h = 1;$	$\Delta v \leq 1$	and $\Delta c > 1$ to $< 3$
		or $\Delta v > 1$ to $< 3$	and $\Delta c < 3$	
		$\Delta h = 2;$	$\Delta v = 0$	and $\Delta c > 0$ to $< 2$
		or $\Delta v > 0$ to $< 2$	and $\Delta c < 2$	
Prominent <sup>1</sup>	P	$\Delta h = 0;$	$\Delta v \geq 4$	or $\Delta c \geq 4$
		$\Delta h = 1;$	$\Delta v \geq 3$	or $\Delta c \geq 3$
		$\Delta h = 2;$	$\Delta v \geq 2$	or $\Delta c \geq 2$
		$\Delta h \geq 3;$		

<sup>1</sup> If compared colors have both a value  $\leq 3$  and a chroma of  $\leq 2$ , the contrast is *Faint*, regardless of hue differences.

## Tabular List for Determination of Color Contrast

**Hues are the same ( $\Delta h = 0$ )<sup>1</sup>**

$\Delta$ Value	$\Delta$ Chroma	Contrast
0	$\leq 1$	Faint
0	2	Distinct
0	3	Distinct
0	$\geq 4$	Prominent
1	$\leq 1$	Faint
1	2	Distinct
1	3	Distinct
1	$\geq 4$	Prominent
$\leq 2$	$\leq 1$	Faint
$\leq 2$	2	Distinct
$\leq 2$	3	Distinct
$\leq 2$	$\geq 4$	Prominent
3	$\leq 1$	Distinct
3	2	Distinct
3	3	Distinct
3	$\geq 4$	Prominent
$\geq 4$	—	Prominent

**Hues differ by 1 ( $\Delta h = 1$ )<sup>1</sup>**

$\Delta$ Value	$\Delta$ Chroma	Contrast
0	$\leq 1$	Faint
0	2	Distinct
0	$\geq 3$	Prominent
1	$\leq 1$	Faint
1	2	Distinct
1	$\geq 3$	Prominent
2	$\leq 1$	Distinct
2	2	Distinct
2	$\geq 3$	Prominent
$\geq 3$	—	Prominent

**Hues differ by 2 ( $\Delta h = 2$ )<sup>1</sup>**

$\Delta$ Value	$\Delta$ Chroma	Contrast
0	0	Faint
0	1	Distinct
0	$\geq 2$	Prominent
1	$\leq 1$	Distinct
1	$\geq 2$	Prominent
$\geq 2$	—	Prominent

**Hues differ by 3 or more ( $\Delta h \geq 3$ )<sup>1</sup>**

$\Delta$ Value	$\Delta$ Chroma	Contrast
Color contrast is Prominent, except for low chroma and value <sup>1</sup>		Prominent

<sup>1</sup> Exception: If both colors have a value  $\leq 3$  and a chroma  $\leq 2$ , the color contrast is *Faint*, regardless of hue differences.

**REDOXIMORPHIC FEATURES - COLOR**—Use standard Munsell® notation from the “Soil Color” section; e.g., *light brownish gray* or *2.5Y 6/2*.

**REDOXIMORPHIC FEATURES - MOISTURE STATE**—Describe the moisture condition of the redoximorphic feature (use “Soil Color - Moisture State” table); e.g., *Moist (M)*.

Moisture State	Code
Dry	D
Moist	M

**REDOXIMORPHIC FEATURES - SHAPE**—Describe the shape of the redoximorphic feature (use “Concentrations - Shape” table); e.g., *Spherical (S)*.

**REDOXIMORPHIC FEATURES - LOCATION**—Describe the location(s) of the redoximorphic feature within the horizon (use “Concentrations - Location” table); e.g., *In the matrix around depletions (MAD)*.

**REDOXIMORPHIC FEATURES - HARDNESS**—Describe the relative force required to crush the redoximorphic feature (use the same classes and criteria as the “Rupture Resistance for Blocks/Peds/Clods-Cementation” column); e.g., *Strongly Cemented (ST)*.

**REDOXIMORPHIC FEATURES - BOUNDARY**—The gradation between the redoximorphic feature and the adjacent matrix (use “Concentrations - Boundary” table; p. 2-27); e.g., *Sharp (S)*.

## MOTTLES

Describe mottles (areas that differ from the matrix color). Mottles commonly have a lithomorphic or lithochromic (e.g., gray shale) geologic origin rather than pedogenic. Mottles do not indicate existing redox conditions. Describe *Redoximorphic Features* and *Ped and Void Surface Features* (e.g., clay films) separately from mottles. Record **Quantity Class** (in NASIS, estimate “Percent of Horizon Area Covered”), **Size**, **Contrast**, **Color**, and **Moisture State** (D or M). **Shape** is an optional descriptor. A complete example is: *common (15%), medium, distinct, reddish yellow, moist, irregular mottles*; or *c, 2, d, 7.5YR 7/8, M, I mottles*.

**MOTTLE QUANTITY (Percent of Area Covered)**—See graphics for **% of Area Covered** (2, 20%), p. 7-1 to p. 7-9.

Quantity Class	Code		Criteria: range in percent
	Conv.	NASIS	
Few	f	%	<2% of surface area
Common	c	%	2 to <20% of surface area
Many	m	%	≥20% of surface area

**MOTTLE SIZE**—Size refers to dimensions as seen on a plane. If mottle length is <3 times the mottle width, record the greater of the two. If length is >3 times width, record the smaller dimension. (See graphic on p. 2-14.)

Size Class	Code	Criteria
Fine	1	0.25 to < 2 mm
Medium	2	2 to < 5 mm
Coarse	3	5 to < 20 mm
Very Coarse	4	20 to < 76 mm
Extremely Coarse	5	≥ 76 mm

**MOTTLE CONTRAST**—Use **Redoximorphic Feature - Contrast** criteria and table (p. 2-15).

**MOTTLE COLOR**—Use standard Munsell® notation of hue, value, and chroma; e.g., *5YR 4/4* (for reddish brown).

**MOTTLE MOISTURE STATE**—Record moisture condition of mottle (don’t confuse with soil water state); e.g., *moist (M)* or *dry (D)*.

**MOTTLE SHAPE (optional)**—Use “Concentrations - Shape” table; e.g., *irregular*.

**MOTTLE LOCATION (optional)**—Use **(Soil) Matrix Color - Location or Condition** table; e.g., *interior*.

## ***CONCENTRATIONS (DISCUSSION)***


---

Concentrations are soil features that form by accumulation of material during pedogenesis. Dominant processes involved are chemical dissolution/precipitation; oxidation and reduction; and physical and/or biological removal, transport, and accrual. Types of concentrations (modified from Soil Survey Division Staff, 1993) include the following:

1. **Finely Disseminated Materials** are physically small precipitates (e.g., salts, carbonates) dispersed throughout the matrix of a horizon. The materials cannot be readily seen (10X lens) but can be detected by a chemical reaction (e.g., effervescence of  $\text{CaCO}_3$  by HCl) or other proxy indicators.
2. **Masses** are noncemented ("Rupture Resistance-Cementation Class" of *Extremely Weakly Cemented* or less) bodies of accumulation of various shapes that cannot be removed as discrete units and do not have a crystal structure that is readily discernible in the field (10X hand lens). This includes finely crystalline salts and *Redox Concentrations* that do not qualify as nodules or concretions.
3. **Nodules** are cemented (*Very Weakly Cemented* or greater) bodies of various shapes (commonly spherical or tubular) that can be removed as discrete units from soil and don't slake. Crystal structure is not discernible with a 10X hand lens.
4. **Concretions** are cemented bodies (*Very Weakly Cemented* or greater) that don't slake and are similar to nodules, except for the presence of visible concentric layers of material around a point, line, or plane. The terms "nodule" and "concretion" are not interchangeable.
5. **Crystals** are macro-crystalline forms of relatively soluble salts (e.g., halite, gypsum, carbonates) that form *in situ* by precipitation from soil solution. The crystalline shape and structure are readily discernible in the field with a 10X hand lens.
6. **Biological Concentrations** are discrete bodies accumulated by a biological process (e.g., fecal pellets) or pseudomorphs of biota or biological processes (e.g., insect casts) formed or deposited in soil.
7. **Inherited Minerals** are field-observable particles (e.g., mica flakes) or aggregates (e.g., glauconite pellets) that impart distinctive soil characteristics and formed by geologic processes in the original parent material and subsequently inherited by the soil rather than formed or concentrated by pedogenic processes. Included here due to historical conventions; not all concentrations descriptors may apply (e.g., shape, color).

General conventions for documenting various types of concentrations:

Type of Distribution	Documentation	Examples
Finely Disseminated (discrete bodies not visible)	Horizon Suffix, Concentrations (finely disseminated)	Carbonates (none) Salts (Bz, Bn)
Masses, Nodules, Concretions, Crystals, Biological Features	Redoximorphic Features, or Concentrations	Mn nodules Fe concretions Insect casts
Continuous Cementation	Terms in Lieu of Texture	Duripan Petrocalcic



## **CONCENTRATIONS**

Record **Kind, Quantity** (percent of area covered), **Size, Contrast, Color, Moisture State, Shape, Location, Hardness, and Boundary**. A complete example is: *many, fine, prominent, white, moist, cylindrical, carbonate nodules in the matrix, moderately cemented, clear, or m, 1, p, 10YR 8/1, M, c, CAN, MAT, M, c.*

**CONCENTRATIONS - KIND**—Identify the composition and the physical state of the concentration in the soil. **NOTE:** Table subheadings (e.g., *Masses*) are a guide to various physical states of materials. Materials with similar or identical chemical composition may occur in multiple physical states (under several subheadings); e.g., *salt masses* and *salt crystals*.

<b>CONCENTRATIONS (NONREDOX)</b> ( <i>accumulations of material</i> )			
Kind	Code	Kind	Code
<b>FINELY DISSEMINATED</b> ( <i>bodies not visible by unaided eye; detectable by chemical tests, e.g., Effervescence</i> )			
finely disseminated carbonates	FDC	finely disseminated salts	FDS
finely disseminated gypsum	FDG		
<b>MASSES</b> ( <i>noncemented; crystals not visible with 10X hand lens</i> )			
barite masses ( $BaSO_4$ )	BAM	gypsum masses ( $CaSO_4 \cdot 2H_2O$ ) <i>crystals not visible</i>	GYM
carbonate masses ( $Ca, Mg, NaCO_3$ )	CAM	salt masses ( $NaCl, Na-Mg$ sulfates)	SAM
clay bodies	CBM	silica masses	SIM
gypsum crystal clusters ( <i>nests</i> ) very fine crystals	GNM		
<b>NODULES</b> ( <i>cemented; noncrystalline at 10X, no layers</i> )			
carbonate nodules <sup>1</sup>	CAN	opal	OPN
durinodes ( $SiO_2$ )	DNN	ortstein nodules	ORT
gibbsite nodules ( $Al_2O_3$ )	GBN		
<b>CONCRETIONS</b> ( <i>cemented; noncrystalline at 10X, distinct layers</i> )			
carbonate concretions <sup>1</sup>	CAC	silica concretions	SIC
gibbsite concretions	GBC	titanium oxide concretions	TIC
<b>CRYSTALS</b> ( <i>crystals visible with 10X hand lens or larger</i> )			
barite crystals ( $BaSO_4$ )	BAX	salt crystals ( $NaCl, Na-Mg$ sulfates)	SAX
calcite crystals ( $CaCO_3$ )	CAX	satin spar crystals ( $CaSO_4 \cdot 2H_2O$ )	SSC
gypsum crystals ( <i>unspecified; <math>CaSO_4 \cdot 2H_2O</math></i> )	GYX	selenite crystals ( $CaSO_4 \cdot 2H_2O$ )	SEC

<b>CONCENTRATIONS (NONREDOX)</b> ( <i>accumulations of material</i> )			
<b>Kind</b>	<b>Code</b>	<b>Kind</b>	<b>Code</b>
<b>BIOLOGICAL CONCENTRATIONS</b> ( <i>entities, byproducts, or pseudomorphs</i> )			
diatoms <sup>2</sup>	DIB	root sheaths	RSB
fecal pellets	FPB	shell fragments ( <i>terrestrial or aquatic</i> )	SFB
insect casts <sup>3</sup> ( <i>e.g., cicada mold</i> )	ICB	sponge spicules <sup>2</sup>	SSB
plant phytoliths <sup>2</sup> ( <i>plant opal</i> )	PPB	worm casts <sup>3</sup>	WCB
<b>INHERITED MINERALS</b> ( <i>geogenic</i> ) <sup>4</sup>			
glauconite pellets	GLI	volcanic glass	VOG
mica flakes	MIC		
<b>MISCELLANEOUS</b> <sup>5</sup>			
carbonate bands	CBA	carbonate ooliths	CAO
carbonate beds	CBE	carbonate pisoliths	CAP
carbonate laminae	CAL	carbonate root casts	CRC

<sup>1</sup> For example: *loess doll* (aka "*loess kindchen*," "*loess puppies*," etc.).

<sup>2</sup> Commonly requires magnification >10X to be observed.

<sup>3</sup> Worm casts are ovoid, fecal pellets excreted by earthworms. Insect casts are cemented (e.g., CaCO<sub>3</sub>) molds of insect bodies or burrows.

<sup>4</sup> Minerals inherited from parent material rather than formed in soil.

<sup>5</sup> See Discussion on carbonate stages (p. 2–28).



**CONCENTRATIONS - QUANTITY (PERCENT OF AREA COVERED)**—See graphics for **% of Area Covered** (2, 20%) beginning on p. 7-1.

Class	Code		Criteria: percent of surface area covered
	Conv.	NASIS	
Few	f	#	< 2
Common	c	#	2 to < 20
Many	m	#	≥ 20


**CONCENTRATIONS - SIZE** (Same as “RMFs” and “Mottle Size Classes”)—See graphic on page p. 2-24.)

Size Class	Code	Criteria
Fine	1	0.25 to < 2 mm
Medium	2	2 to < 5 mm
Coarse	3	5 to < 20 mm
Very Coarse	4	20 to < 76 mm
Extremely Coarse	5	≥ 76 mm

**CONCENTRATIONS - CONTRAST**—Use “RMF - Contrast” table or chart; e.g., *distinct*.

**CONCENTRATIONS - COLOR**—Use standard Munsell® notation; e.g., 7.5YR 8/1.












**CONCENTRATIONS - MOISTURE STATE**—Use “Soil Color - Moisture State” table; i.e., *Moist (M)* or *Dry (D)*.



**CONCENTRATIONS - SHAPE <sup>1</sup>** (also used for **Mottles, Redoximorphic Features**)

Shape <sup>1</sup>	Code	Criteria
cubic	CU	crudely equidimensional blocklike units
cylindrical	C	tubular and elongated bodies; e.g., filled wormholes and insect burrows
dendritic	D	tubular, elongated, and branched bodies; e.g., pipestems (root pseudomorphs)
irregular	I	bodies of nonrepeating spacing or shape
lenticular	L	disk-shaped forms with thicker centers and thinning towards outer edge (e.g., double-convex lens)
pendular	PE	irregular drapes, coatings, or nodules suspended from underside of coarse fragments (e.g., pendular gypsum masses)
platy	P	relatively thin, tabular sheets, lenses; e.g., lamellae
reticulate	R	crudely interlocking bodies with similar spacing; e.g., plinthite
rosettelike	RO	interlocking blades radiating out from a central point forming petal-like clusters; e.g., barite
spherical	S	well-rounded to crudely spherical bodies; e.g., Fe/Mn "shot"
threadlike	T	thin (e.g., <1 mm diam.) elongated filaments; generally not dendritic (e.g., very fine CaCO <sub>3</sub> stringers)

<sup>1</sup> Shape terms are presented as adjectives due to the typical data string output (e.g., *dendritic carbonate concretions*).

<p><b>Examples of Mottles, Concentrations, and RMF Shapes</b></p>	<p>cubic (e.g., halite)</p> 
<p>cylindrical (e.g., filled worm holes)</p> 	<p>dendritic (e.g., branched root pseudo-morphs)</p> 
<p>irregular</p> 	<p>lenticular (e.g., gypsum)</p> 
<p>pendular (e.g., CaCO<sub>3</sub>, CaSO<sub>4</sub>, SiO<sub>2</sub>)</p> 	<p>platy (e.g., lamellae)</p> 
<p>reticulate (e.g., plinthite)</p> 	<p>rosettelike (e.g., barite, gypsum)</p> 
<p>spherical (e.g., Fe/Mn shot)</p> 	<p>threadlike (e.g., very fine CaCO<sub>3</sub> stringers and filaments)</p> 

**CONCENTRATIONS - LOCATION**—Describe the location(s) of the concentration (or depletion for RMFs) within the horizon. Historically called **Concentrations - Distribution**.

Location	Code
<b>MATRIX (in soil matrix; not associated with ped faces or pores)</b>	
In the matrix ( <i>not associated with peds/pores</i> )	MAT
In matrix surrounding redox depletions	MAD
In matrix surrounding redox concentrations	MAC
Throughout ( <i>e.g., finely disseminated carbonates</i> )	TOT
<b>PEDS (on or associated with faces of peds)</b>	
Between peds	BPF
Infused into the matrix along faces of peds ( <i>hypocoats</i> )	MPF
On faces of peds ( <i>all orientations</i> )	APF
On horizontal faces of peds	HPF
On vertical faces of peds	VPF

Location	Code
<b>PORES (in pores or associated with surfaces along pores)</b>	
Infused into the matrix adjacent to pores ( <i>hypocoats</i> ; see <i>Coats/Films graphic p. 2-34</i> )	MPO
Lining pores (see <i>Coats/Films graphic p. 2-34</i> )	LPO
On surfaces along pores	SPO
On surfaces along root channels	RPO
<b>OTHER</b>	
In cracks	CRK
At top of horizon	TOH
Around rock fragments	ARF
On bottom of rock fragments ( <i>e.g., pendants</i> )	BRF
On slickensides	SSS
Along lamina or strata surfaces	ALS

**CONCENTRATIONS - HARDNESS**—Describe the relative force required to crush the concentration body (use the same criteria and classes as in the **"Rupture Resistance for Blocks, Peds, and Clods – Cementation"** column (exclude the *Noncemented* class); e.g., *Moderately Cemented*.)

**CONCENTRATIONS - BOUNDARY**—The gradation between feature and matrix. (Also used to describe **Redoximorphic Features - Boundary**.)

Class	Code	Criteria
Sharp	S	Color changes in <0.1 mm between the feature and the soil matrix; change is abrupt even under a 10X hand lens.
Clear	C	Color changes within 0.1 to <2 mm between the feature and the soil matrix; gradation is visible without 10X lens.
Diffuse	D	Color changes in ≥2 mm between the feature and the soil matrix; gradation is easily visible without 10X hand lens.

## ***PEDOGENIC CARBONATE STAGES (DISCUSSION)***

---

**Pedogenic Carbonate Development:** In arid, semiarid, and subhumid environments, pedogenic carbonate accumulation is of overarching and unifying importance. The present morphological concepts and intellectual basis for soil-carbonate horizons and stages that follow originate from the seminal contributions of Leland Gile (Gile, 1961, 1970, 1975, 1993; Gile and Grossman, 1968; Gile and Grossman, 1979; Gile et al., 1966; Gile et al., 2007).

Calcium carbonate ( $\text{CaCO}_3$ ) mediates or controls key chemical and physical soil properties (e.g., pH, nutrient availability, dispersion-flocculation, organic matter stabilization). Calcium carbonate in soil may be inherited from parent material and/or may accumulate via pedogenic processes. Climate (precipitation, temperature, evapotranspiration), carbonate solubility, and microbial biomineralization control and constrain the rate and quantity of pedogenic carbonate accumulation. Major carbonate accumulation occurs mainly in subhumid to arid regions (precipitation  $\sim < 750\text{mm}$ /annum). Pedogenic carbonate formation requires a Ca source, such as mineral weathering, concentration by soil or ground water evaporation, and/or input via dust and/or precipitation. The carbonate ( $\text{CO}_3^{2-}$ ) source is plant and microbial respiration ( $\text{CO}_2$ ) via the  $\text{CO}_2\text{-HCO}_3^{-1}\text{-CO}_3^{2-}$  equilibria. Beyond the major climatic control, pedogenic carbonate accumulation depends on a balance among geomorphic age or landscape stability, soil water movement (at both profile and landscape scales), soil texture, and vegetation type and quantity.

Pedogenic carbonate accumulation follows a morphogenetic development sequence starting as horizon features, such as carbonate coatings, masses, and fine nodules. If carbonate continues to accumulate, it may entirely engulf, plug, and cement soil horizons. Carbonate-cemented soil horizons are generically termed caliche or calcrete and are recognized in Soil Taxonomy as petrocalcic horizons. Pedogenic carbonate accumulation is closely linked to soil age (Gile et al., 1981; Machette, 1985). Soils on progressively older geomorphic surfaces contain sequentially more pedogenic carbonate. The progression of carbonate development and morphology has been defined as Stages I through VI (see figures A and B) (Gile et al., 1966; Gile et al., 1981; Brock, 2007; Machete, 1985; Bachman and Machette, 1977).


Pedogenic carbonate stage morphology and expression are initially different for a fine earth matrix (fig. A; e.g., lacustrine, distal fan deposits, eolian deposits) than for a coarse fragment matrix (fig. B; e.g., proximal alluvial fan deposits, channel deposits) soils (Gile et al., 1966; Flach et al., 1969; Gile et al., 1981). The time required for carbonate stage development depends on soil texture and its control on porosity. A fine-grained soil has greater surface area and total porosity that requires a correspondingly greater

carbonate quantity to fill voids and form equivalent carbonate stages compared to a coarse-textured soil (Gile, 1993; Gile et al., 1981). Thus, it takes more time for a fine-grained soil to reach the same carbonate stage as a coarse-textured soil under the same conditions.

**Multiple Carbonate Stages:** Pedogenic carbonates (especially Stages IV through VI) are durable and persistent in arid environments. If different sedimentation events and subsequent pedogenesis (separated by time) occur to produce a stacked soil sequence (paleosols), a pedon may contain multiple carbonate stages. Each soil sequence (sediment package) is evaluated independently, and a carbonate stage is attached to horizons in that sequence. For example, Stage II carbonate may occur in a soil overlying a buried soil with Stage IV carbonate; e.g., A, 0-12 cm; Bt, 12-22 cm; Bk (Stage II  $\text{CaCO}_3$  nodules), 22-65cm; 2Bkkm1 (Stage IV  $\text{CaCO}_3$ ), 65-150 cm; 2Bkkm2 (Stage IV  $\text{CaCO}_3$ ), 150-260 cm. Moreover, on stable geomorphic surfaces climatic shifts may superpose younger carbonate forms into preexisting, more advanced stages within the same soil. For example, carbonate nodules may occur above and into a stage IV Bkkm; e.g., A, 0-11 cm; Bt, 11-22 cm; Bk ( $\text{CaCO}_3$  nodules), 22-65 cm; Bkkm (Stage IV  $\text{CaCO}_3$ ), 65-150 cm; 2Bkkm (Stage IV  $\text{CaCO}_3$ ), 150-260 cm. Despite the more recent nodules, this soil is *Stage IV* in the morphogenetic sequence.

**Pedogenic Carbonate Stage Description:** Evaluate and record the *Pedogenic Carbonate Stage* by pedon. More than one  $\text{CaCO}_3$  stage may exist in a pedon as a result of multiple sediment layers or shifting climate conditions (see discussion). Pedogenic carbonates are described under both *Concentrations* and *Ped and Void Surface Features*. In arid and semiarid regions, a pedogenic carbonate stage is commonly based upon the overall carbonate morphology in relation to texture and coarse fragment content (see figures A and B). The “stage” is recorded as an interpretive text note following the conventional carbonate concentration description; e.g., *100-165 cm, 2Bkkm (massive indurated  $\text{CaCO}_3$ ), Stage IV*. Assigning a  $\text{CaCO}_3$  stage is an interpretive complement to, but not a replacement for, conventional soil horizonation.

**Figure A - Pedogenic Carbonate Development Stages - Fine Earth Matrix**




Modified from Gile, 1966; Machete, 1985; Brock 2010

DAW 2012



**Figure B - Pedogenic Carbonate Development Stages - Coarse Fragment Matrix**



Modified from Gile, 1966; Machete, 1985; Brock, 2010

DAW 2012

## **PED and VOID SURFACE FEATURES**

These features are coats/films, hypocoats, or stress features formed by translocation and deposition or by shrink-swell processes on or along surfaces. Describe **Kind**, **Amount Class** (percent in NASIS), **Distinctness**, **Location**, and **Color** (dry or moist). An example is: *many, faint, brown 10YR 4/6 (Moist) clay films on all faces of peds or m, f, 10YR 4/6 (M), CLF, PF.*

### **PED and VOID SURFACE FEATURES - KIND** (nonredoximorphic)

<b>Kind</b>	<b>Code</b>	<b>Field Criteria</b>
<b>COATS, FILMS (exterior, adhered to surface)</b>		
carbonate coats	CAF	off-white, effervescent with HCl
silica ( <i>silans, opal</i> )	SIF	off-white, noneffervescent with HCl
clay films ( <i>argillans</i> )	CLF	waxy, exterior coats
clay bridges	BRF	"wax" between sand grains
ferriargillans <i>described as RMF-Kind</i>	<i>see RMFs</i>	Fe <sup>+3</sup> stained clay film
gibbsite coats ( <i>sesquan</i> )	GBF	AlOH <sub>3</sub> , off-white, noneffervescent with HCl
gypsum coats	GYF	CaSO <sub>4</sub> • 2H <sub>2</sub> O
manganese ( <i>mangans</i> ) <i>described as RMF-Kind</i>	<i>see RMFs</i>	black, thin films effervescent with H <sub>2</sub> O <sub>2</sub>
organic stains	OSF	dark organic films
organoargillans	OAF	dark, organic stained clay films
sand coats	SNF	separate grains visible with 10X
silt coats <sup>1</sup>	SLF	separate grains not visible at 10X
skeletans <sup>2</sup> (sand or silt)	SKF	clean sand or silt grains as coats
<b>HYPOCOATS <sup>3</sup> (a stain infused beneath a surface)</b>		
<b>STRESS FEATURES (a smeared exterior face)</b>		
pressure faces (i.e., stress cutans)	PRF	look like clay films; sand grains uncoated

Kind	Code	Field Criteria
slickensides (pedogenic)	SS	shrink-swell shear features (e.g., grooves, striations, glossy surface) on pedo-structure surfaces (e.g., wedges, bowls); can be horizontal
slickensides (geogenic)	SSG	vertical/oblique, roughly planar shear face from external stress (e.g., faults; mass movement); striations, grooves

<sup>1</sup> Individual silt grains are not discernible with a 10X lens. Silt coats occur as a fine, off-white, noneffervescent, “grainy” coat on ped surfaces.


<sup>2</sup> Skeletans are (pigment) stripped grains >2 μm and <2 mm (Brewer, 1976). Preferably describe either *silt coats* (grains not discernible with 10X lens) or *sand coats* (grains discernible with 10X lens).

<sup>3</sup> Hypocoats, as used here, are field-scale features commonly expressed only as redoximorphic features. Micromorphological hypocoats include nonredoximorphic features (Bullock et al., 1985).



**Coats/Films**

**Peds**

**Pores**





General  
On surface





argillans

(sand grains "waxed" over)




skeletalans



**Hypocoats**




Below surface




**Stress Features**

pressure face

slickensides



(sand grains stand clear)



(striations, grooves, glossy sheen)

**PED and VOID SURFACE FEATURES - AMOUNT**—Estimate the relative percent of the visible surface area that a ped surface feature occupies in a horizon. (See graphics for **% of Area Covered** [5, 25, 50, 90%] beginning on page 7.1.) In NASIS, record the estimate as a numeric percent; e.g., 20%.

Amount Class	Code		Criteria: percent of surface area
	Conv.	NASIS	
Very Few	vf	%	< 5 percent
Few	f	%	5 to < 25 percent
Common	c	%	25 to < 50 percent
Many	m	%	50 to < 90 percent
Very Many	vm	%	≥ 90 percent

**PED and VOID SURFACE FEATURES - CONTINUITY** (Obsolete in NRCS; replaced with **Ped and Void Surface Features - Amount** in NASIS.)

Continuity Class	Conv. Code	Criteria: features occur as
Continuous	C	Entire Surface Cover
Discontinuous	D	Partial Surface Cover
Patchy	P	Isolated Surface Cover

**PED and VOID SURFACE FEATURES - DISTINCTNESS**—The relative extent to which a ped surface feature visually stands out from adjacent material.

Distinctness Class	Code	Criteria:
Faint	F	Visible only with magnification (10X hand lens); little contrast between materials.
Distinct	D	Visible without magnification; significant contrast between materials.
Prominent	P	Markedly visible without magnification; sharp visual contrast between materials.

**PED and VOID SURFACE FEATURES - LOCATION**—Specify where ped surface features occur within a horizon; e.g., *Between sand grains*.

Location	Code
<b>PEDS</b>	
On all faces of peds (vertical and horizontal)	PF
On bottom faces of peds	BF
On top faces of peds	TF
On tops of soil columns	TC
On vertical faces of peds	VF
<b>OTHER (NONPED)</b>	
Between sand grains (bridging)	BG
On bedrock	BK
On bottom surfaces of rock fragments	BR
On concretions	CC
On nodules	NO
On rock fragments	RF
On slickensides	SS
On surfaces along pores	SP
On surfaces along root channels	SC
On top surfaces of rock fragments	TR

**PED and VOID SURFACE FEATURES - COLOR**—Use standard Munsell® notation (hue, value, chroma) to record feature color. Indicate whether the color is Moist (M) or Dry (D); e.g., *7.5R 5/8 M*.

## **SOIL TEXTURE**

---

Soil texture is the numerical proportion (weight percentage) of the sand, silt, and clay separates in the fine-earth fraction ( $\leq 2$  mm). Soil texture is field estimated by hand or lab measured by hydrometer or pipette and placed within the textural triangle to obtain **Texture Class**.


Record the **Texture Class**; e.g., *loam*; or Subclass; e.g., *fine sandy loam*; or choose a **Term in Lieu of Texture**; e.g., *gravel*. If appropriate, use a **Texture Class Modifier**; e.g., *gravelly loam*.

**NOTE: Soil Texture** includes only the fine-earth fraction ( $\leq 2$  mm). "Whole-soil Particle-Size Distribution" includes the fine-earth fraction ( $\leq 2$  mm, wt %) and coarse fragments ( $> 2$  mm). (**NOTE:**

For fragments  $\geq 76$  mm in diameter, visually estimate the volume percent, which is then converted to a weight basis using the estimated particle density [pd] and bulk density [ $B_d$ ].)

**TEXTURE CLASS—**

Texture Class or Subclass	Code	
	Conv.	NASIS
Coarse Sand	cos	COS
Sand	s	S
Fine Sand	fs	FS
Very Fine Sand	vfs	VFS
Loamy Coarse Sand	lcos	LCOS
Loamy Sand	ls	LS
Loamy Fine Sand	lfs	LFS
Loamy Very Fine Sand	lvfs	LVFS
Coarse Sandy Loam	cosl	COSL
Sandy Loam	sl	SL
Fine Sandy Loam	fsl	FSL
Very Fine Sandy Loam	vfsl	VFSL
Loam	l	L
Silt Loam	sil	SIL
Silt	si	SI
Sandy Clay Loam	scl	SCL
Clay Loam	cl	CL
Silty Clay Loam	sicl	SICL
Sandy Clay	sc	SC
Silty Clay	sic	SIC
Clay	c	C



**TEXTURE MODIFIERS**—Conventions for using “Rock Fragment Texture Modifiers” and for using textural adjectives that convey the “% volume” ranges for **Rock Fragments - Quantity and Size**.

Frag. Content Vol. %	Rock Fragment Modifier Usage
<15	No texture class modifier (noun only; e.g., <i>loam</i> ).
15 to <35	Use fragment-size adjective with texture class; e.g., <i>gravelly loam</i> .
35 to <60	Use “ <b>very</b> ” with fragment-size adjective with texture class; e.g., <i>very gravelly loam</i> .
60 to <90	Use “ <b>extremely</b> ” with fragment-size adjective with texture class; e.g., <i>extremely gravelly loam</i> .
≥90	No adjective or modifier. If ≤10% fine earth, use the appropriate fragment-size class name for the dominant size class; e.g., <i>gravel</i> . Use <b>Terms Used in Lieu of Texture</b> (see table on p. 2-43).



**TEXTURE MODIFIERS—Quantity and Size adjectives.**

Rock Fragments: Quantity and Size <sup>1</sup>	Code		Criteria: total (rock) fragment volume % dominated by (name size) <sup>1</sup>
	Conv.	NASIS	
<b>ROCK FRAGMENTS (&gt;2 mm; ≥ Strongly Cemented)</b>			
Gravelly	GR	GR	≥15% but <35% gravel
Fine Gravelly	FGR	GRF	≥15% but <35% fine gravel
Medium Gravelly	MGR	GRM	≥15% but <35% med. gravel
Coarse Gravelly	CGR	GRC	≥15% but <35% coarse gravel
Very Gravelly	VGR	GRV	≥35% but <60% gravel
Extremely Gravelly	XGR	GRX	≥60% but <90% gravel
Cobbly	CB	CB	≥15% but <35% cobbles
Very Cobbly	VCB	CBV	≥35% but <60% cobbles
Extremely Cobbly	XCB	CBX	≥60% but <90% cobbles
Stony	ST	ST	≥15% but <35% stones
Very Stony	VST	STV	≥35% but <60% stones
Extremely Stony	XST	STX	≥60% but <90% stones
Bouldery	BY	BY	≥15% but <35% boulders
Very Bouldery	VBY	BYV	≥35% but <60% boulders
Extremely Bouldery	XBY	BYX	≥60% but <90% boulders
Channery	CN	CN	≥15% but <35% channers
Very Channery	VCN	CNV	≥35% but <60% channers
Extremely Channery	XCN	CNX	≥60% but <90% channers
Flaggy	FL	FL	≥15% but <35% flagstones
Very Flaggy	VFL	FLV	≥35% but <60% flagstones
Extremely Flaggy	XFL	FLX	≥60% but <90% flagstones
<b>PARAROCK FRAGMENTS (&gt;2 mm; &lt; Strongly Cemented) <sup>2, 3</sup></b>			
Paragravelly	PGR	PGR	(same criteria as for gravelly)
Very Paragravelly	VPGR	PGRV	(same criteria as for very gravelly)
Extr. Paragravelly	XPGR	PGRX	(same criteria as for extr. gravelly)
etc.	etc.	etc.	(same criteria as for nonpara)

Rock Fragments: Quantity and Size <sup>1</sup>	Code		Criteria: total (rock) fragment volume % dominated by (name size) <sup>1</sup>
	Conv.	NASIS	
<b>COMPOSITE ROCK AND ARTIFACT FRAGMENTS <sup>4</sup></b>			
Gravelly - Artificial	GRART	GRART	(same criteria as for gravelly)
Very Gravelly - Artificial	VGRART	GRVART	(same criteria as for very gravelly)
Extremely Gravelly - Artificial	XGRART	GRXART	(same criteria as for extr. gravelly)
etc.	etc.	etc.	(same criteria as for noncomposite)

<sup>1</sup> The "Quantity" modifier (e.g., *very*) is the volume % whole soil of the total rock fragment content. The "Size" modifier (e.g., *cobbly*) is independently based on the largest, dominant fragment size. (See "Comparison of Particle-Size Classes" table; p. 2-45, first row.) For a size mixture (e.g., *gravel and stones*), a smaller size class is named if its quantity (%) sufficiently exceeds that of a larger size class. For field texture determination, a smaller rock fragment size class must exceed either 1.5 or 2 times the quantity (volume %) of a larger size class before it is named (e.g., 30% gravel and 14% stones=*very gravelly*, but 20% gravel and 14% stones=*stony*). For detailed naming criteria, see NSSH, Part 618, Subpart B, Exhibits, "Rock Fragment Modifier of Texture" (Soil Survey Staff, 2012c).

<sup>2</sup> Use "Para" prefix if the rock fragments are soft (i.e., meet criteria for "para"). (Rupture Resistance – Cementation Class is < *Strongly Cemented*, and fragments do not slake [slake test: ≈3 cm (1 inch) diam. block, air dried, then submerged in water for ≥1 hour; collapse/disaggregation="slaking"].)

<sup>3</sup> For "Para" codes, add "P" to "Size" and "Quantity" code terms. Precedes noun codes and follows quantity adjectives; e.g., paragravelly=*PGR*; very paragravelly=*PGRV*.

<sup>4</sup> Used if a horizon contains both rock and artifact fragments >2 mm that are both cohesive and persistent and whose combined % by volume is ≥15%; use appropriate *Quantity Class* (the dominant size fraction is named).

**(COMPOSITIONAL) TEXTURE MODIFIERS** <sup>1, 2</sup>—Compositional adjectives (e.g., ashy silt loam).

Types	Code	Criteria
<b><i>VOLCANIC</i></b>		
Ashy	ASHY	Andic soil properties, and is neither hydrous nor medial, or $\geq 30\%$ of the $< 2$ mm fraction is 0.02 to 2.00 mm in size, $\geq 5\%$ is volcanic glass, and the $[Al + 1/2 Fe, \% \text{ by ammonium oxalate}] \times 60 + \% \text{ volc glass}$ is $\geq 30$
Hydrous	HYDR	Andic soil properties, and with field moist 15 bar water content $\geq 100\%$ of the dry weight
Medial	MEDL	Andic soil properties, and with field moist 15 bar water content $\geq 30\%$ to $< 100\%$ of the dry weight, or $\geq 12\%$ water content for air-dried samples
<b><i>ORGANIC SOIL MATERIALS</i></b>		
Grassy <sup>3</sup>	GS	OM $> 15\%$ (vol.) grassy fibers
Herbaceous <sup>3</sup>	HB	OM $> 15\%$ (vol.) herbaceous fibers
Mossy <sup>3</sup>	MS	OM $> 15\%$ (vol.) moss fibers
Woody <sup>3</sup>	WD	OM $\geq 15\%$ (vol.) wood pieces or fibers
<b><i>HIGHLY ORGANIC MINERAL MATERIALS</i></b>		
Highly Organic <sup>4</sup>	HO	Organic carbon (wt %) is: $> 5$ to $< 20\%$ (no mineral clays) $12$ to $< 20\%$ (if mineral clay is $\geq 60\%$ ) or $5+$ (clay % $\times 0.12$ to $< 20\%$ ) (if mineral clay is $< 60\%$ )
Mucky <sup>5</sup>	MK	Mineral soil $> 10\%$ OM and $< 17\%$ fibers
Peaty <sup>5</sup>	PT	Mineral soil $> 10\%$ OM and $> 17\%$ fibers
<b><i>LIMNIC MATERIALS (used only with Histosols)</i></b>		
Coprogenous	COP	Limnic layer with many very small fecal pellets
Diatomaceous	DIA	Limnic layer composed of diatoms
Marly	MR	Light-colored limnic layer composed of $CaCO_3$ mud

Types	Code	Criteria
<b>ANTHROPOGENIC MATERIALS</b>		
Artifactual	ART	≥15% but <35% (vol.) artifacts
Very Artifactual	ARTV	≥35% but <60% (vol.) artifacts
Extremely Artifactual <sup>2</sup>	ARTX	≥60% but <90% (vol.) artifacts
<b>OTHER</b>		
Cemented	CEM	Material is “cemented” by ≥1 cementing agents; does not slake
Gypsiferous	GYP	≥15 to <40% (by weight) gypsum
Permanently Frozen	PF	e.g., Permafrost

<sup>1</sup> **(Compositional) Texture Modifiers** can be used with the **Soil Texture Name** (e.g., *gravelly ashy loam*) or with **Terms Used in Lieu of Texture** (e.g., *mossy peat*). For complete definitions and usage of **(Compositional) Texture Modifiers**, see NSSH, Part 618.67 (Soil Survey Staff, 2012c).

<sup>2</sup> If artifact fragments are >90% (by vol.), no texture is described and a **Term Used in Lieu of Texture** is applied (i.e., *artifacts*).

<sup>3</sup> Used to modify muck, mucky peat, or peat terms in histic epipedons and organic horizons (of any thickness) that are saturated with water for ≥30 consecutive days in normal years (or are artificially drained), including those in Histels and Histosols (except Folists).

<sup>4</sup> Used only with near-surface horizons of mineral soils saturated <30 cumulative days in normal years (and *not* artificially drained).

<sup>5</sup> Designed for near-surface horizons saturated ≥30 cumulative days annually.

**TERMS USED IN LIEU OF TEXTURE**—nouns (used only if fragments or artifacts are >90% by volume). Bedrock, organic terms, gypsum materials, and permanent water have different criteria.

<b>Terms Used in Lieu of Texture</b>	<b>Code</b>
<b>SIZE (ROCK FRAGMENTS) ≥ Strongly Cemented</b>	
Gravel	GR
Cobbles	CB
Stones	ST
Boulders	BY
Channers	CN
Flagstones	FL
<b>SIZE (PARAROCK FRAGMENTS) &lt; Strongly Cemented</b>	
Paragravel	PG
Paracobbles	PCB
Parastones	PST
Paraboulders	PBY
Parachanners	PCN
Paraflagstones	PFL
<b>COMPOSITION</b>	
<b><i>Cemented/Consolidated:</i></b>	
Bedrock	BR
<b><i>Organic Soil Materials:</i></b>	
Highly Decomposed Plant Material (Oa) <sup>1</sup>	HPM
Moderately Decomposed Plant Material (Oe) <sup>1</sup>	MPM
Slightly Decomposed Plant Material (Oi) <sup>1</sup>	SPM
Muck <sup>2</sup> (≈Oa)	MUCK
Mucky Peat <sup>2</sup> (≈Oe; saturated, moderately decomposed organic matter)	MPT
Peat <sup>2</sup> (≈Oi)	PEAT
<b><i>Other:</i></b>	
Artifacts <sup>3</sup> (human-manufactured materials)	ART
Coarse Gypsum Material	CGM
Fine Gypsum Material	FGM
Ice <sup>5</sup> (permanent, subsurface)	ICE
Water <sup>4</sup> (permanent, subsurface)	W

- <sup>1</sup> Use only with organic horizons of mineral and organic soils that are saturated <30 cumulative days in normal years (and are *not* artificially drained).
- <sup>2</sup> Use only with organic horizons (of any thickness) of mineral and organic soils that are saturated  $\geq 30$  cumulative days in normal years or are artificially drained.
- <sup>3</sup> "Artifacts" is used only to denote presence of artificial materials associated with human activities (bitumen, bricks, construction debris, garbage, etc.).
- <sup>4</sup> Use only for layers found below the soil surface (e.g., a floating bog).
- <sup>5</sup> Used for permanent (nonseasonal), massive, subsurface ice; e.g., a glacial layer.

[ Footnotes below apply to the following table: ]

- <sup>1</sup> Soil Survey Staff, 2011; p. 489.
- <sup>2</sup> Soil Survey Staff, 2011; p. 33. Note: Mineralogy studies may subdivide clay into three size ranges: fine (<0.08  $\mu\text{m}$ ), medium (0.08–0.2  $\mu\text{m}$ ), and coarse (0.2–2  $\mu\text{m}$ ) (Jackson, 1969).
- <sup>3</sup> The Kellogg Soil Survey Laboratory (Lincoln, NE) uses a no. 300 sieve (0.047-mm opening) for the USDA sand/silt measurement. A no. 270 sieve (0.053-mm opening) is more readily available and widely used.
- <sup>4</sup> Soil Survey Staff, 1951; p. 207.
- <sup>5</sup> ASTM, 2011; ASTM designation D2487–92.
- <sup>6</sup> AASHTO, 1997a.
- <sup>7</sup> AASHTO, 1997b.
- <sup>8</sup> Ingram, 1982.

## Comparison of Particle Size Classes in Different Systems

<b>FINE EARTH</b>																							
<b>ROCK FRAGMENTS</b>																							
channers																							
flagst																							
stones																							
boulders																							
6" 15" 24"																							
1.50 380 600 mm																							
<b>USDA<sup>1</sup></b>	Clay <sup>2</sup>		Silt			Sand			Gravel			Boulders											
	fine	co.	fine	co.	v. fi.	fi.	med.	co.	v. co.	coarse	Cob- bles	Stones											
millimeters:	0.0002	.002 mm	.02	.05	.1	.25	.5	1	2 mm	5	20	76	250 mm	600 mm									
U.S. Standard Sieve No. (opening):																							
<sup>3</sup> 300 140 60 35 18 10 4 (3/4") (3") (10") (25")																							
<b>Inter- national<sup>4</sup></b>	Clay		Silt			Sand			Gravel			Stones											
	fine	co.	fine	co.	coarse	fine	medium	co.	fine	coarse	Gravel	Stones											
millimeters:	.002 mm	.02	.20	2 mm	20 mm																		
U.S. Standard Sieve No. (opening):																							
10 (3/4")																							
<b>Unified<sup>5</sup></b>	Silt or Clay																						
	Sand			Gravel			Cobbles			Boulders													
millimeters:	.074	.42	2 mm	4.8	19	76																	
U.S. Standard Sieve No. (opening):																							
200 40 10 4 (3/4") (3")																							
<b>AASHTO<sup>6,7</sup></b>	Clay		Silt			Gravel or Stones			Broken Rock (angular), or Boulders (rounded)														
	fine	co.	fine	co.	co.	fine	med.	co.															
millimeters:	.005 mm	.074	.42	2 mm	9.5	25	75 mm																
U.S. Standard Sieve No. (opening):																							
200 (3/8") (1") (3")																							
phi #:	12	10	9	8	7	6	5	4	3	2	1	0	-1	-2	-3	-4	-5	-6	-7	-8	-9	-10	-12
<b>Modified<sup>8</sup> Wentworth</b>	clay		silt			sand			pebbles			Cobb- bles			boulders								
millimeters:	.00025	.002	.004	.008	.016	.031	.062	.125	.25	.5	1	2	4	8	16	32	64	128	256	512	1024	2048	4092 mm
U.S. Standard Sieve No.:																							
230 120 60 35 18 10 5																							

## ROCK and OTHER FRAGMENTS

These are discrete, water-stable particles >2 mm. Hard fragments (e.g., rock) have a Rupture Resistance - Cementation Class  $\geq$  *Strongly Cemented*. Softer fragments (e.g., pararock) are less strongly cemented. (**NOTE:** Artifacts are addressed separately following this section [p. 2–49].) Describe **Kind**, **Volume Percent** (classes given below), **Roundness or Shape**, **Size** (mm), and **Hardness**; e.g., *granite, 17%, subangular, gravel, indurated*; or *GRA, 17%, SA, GR, I*.

**ROCK and OTHER FRAGMENTS - KIND** (called **FRAGMENTS** in NASIS)—Use the choice list given for **Bedrock - Kind** and the additional choices in the table below. **NOTE:** Interbedded rocks from the “Bedrock - Kind” table are not appropriate choices or terminology for rock fragments.

Kind	Code	Kind	Code
Includes all choices in <b>Bedrock - Kind</b> (except <i>Interbedded</i> ), plus:			
calcrete (caliche) fragments <sup>1</sup>	CA	metamorphic rock fragments, unspecified <sup>2</sup>	MMR
carbonate concretions	CAC	mixed rock fragments <sup>3</sup>	MXR
carbonate nodules	CAN	ortstein fragments	ORF
carbonate rocks <sup>2</sup>	CAR	petrocalcic fragments	PEF
charcoal fragments	CH	petroferric fragments	TCF
cinders	CI	petrogypsic fragments	PGF
durinodes	DNN	plinthite nodules	PLN
duripan fragments	DUF	quartz fragments	QUA
foliated metamorphic rocks <sup>2</sup>	FMR	quartzite fragments	QZT
gibbsite concretions	GBC	scoria fragments	SCO
gibbsite nodules	GBN	sedimentary rock fragments, unspecified <sup>2</sup>	SED
igneous rock fragments, unspecified <sup>2</sup>	IGR	shell fragments	SHF
iron-manganese concretions	FMC	silica concretions	SIC
iron-manganese nodules	FMN	volcanic bombs	VB



Kind	Code	Kind	Code
ironstone nodules	FSN	volcanic rock fragments, unspecified <sup>2</sup>	VOL
lapilli	LA	wood fragments	WO

<sup>1</sup> Fragments strongly cemented by carbonate; may include fragments derived from petrocalcic horizons.

<sup>2</sup> Generic rock names may be appropriate for identifying fragments (e.g., a cobble) but are too general and should *not* be used to name Bedrock - Kind.

<sup>3</sup> Numerous unspecified fragment lithologies are present, as in till or alluvium; not for use with residuum.

### ROCK and OTHER FRAGMENTS - VOLUME PERCENT

(Quantity)—Estimate the quantity (volume percent) of rock and other fragments present. **NOTE:** Refer to the "Total (rock) fragment volume percent" column found under **Texture Modifiers - Quantity and Size** table (p. 2-39).

### ROCK and OTHER FRAGMENTS - SIZE CLASSES AND DESCRIPTIVE TERMS—

Size <sup>1</sup>	Noun	Adjective <sup>2</sup>
<b>SHAPE—SPHERICAL or CUBELIKE (discoidal, subdiscoidal, or spherical)</b>		
>2 - 76 mm diam.	gravel	gravelly
>2 - 5 mm diam.	fine gravel	fine gravelly
>5 - 20 mm diam.	medium gravel	medium gravelly
>20 - 76 mm diam.	coarse gravel	coarse gravelly
>76 - 250 mm diam.	cobbles	cobbly
>250 - 600 mm diam.	stones	stony
>600 mm diam.	boulders	bouldery
<b>SHAPE—FLAT (prismoidal or subprismoidal)</b>		
>2 - 150 mm long	channers	channery
>150 - 380 mm long	flagstones	flaggy
>380 - 600 mm long	stones	stony
>600 mm long	boulders	bouldery

<sup>1</sup> Fragment sizes measured by sieves; class limits have a greater lower limit.

<sup>2</sup> For a mixture of sizes (e.g., both gravel and stones present), the largest size class (most mechanically restrictive) is named. A smaller size class is named only if its quantity (%) sufficiently exceeds that of a larger size class. For field texture determination, a smaller size class must exceed 2 times the quantity (volume %) of a larger size class before it is named (e.g., 30% gravel and 14% stones=*very gravelly*; but 20% gravel and 14% stones=*stony*). For more explicit naming criteria, see NSSH, Part 618, Subpart B, Exhibits, "Rock Fragment Modifier of Texture" (Soil Survey Staff, 2012c).

**ROCK and OTHER FRAGMENTS - ROUNDNESS**—Estimate the relative roundness of rock fragments; use the following classes.

Roundness Class	Code	Criteria: visual estimate <sup>1</sup>
Very Angular	VA	[Use <b>Roundness</b> graphic on p. 2-49]
Angular	AN	
Subangular	SA	
Subrounded	SR	
Rounded	RO	
Well Rounded	WR	

<sup>1</sup> The criteria consist of a visual estimation; use the following graphic.

Estimate the relative roundness of rock fragments. (Ideally, use the average roundness of 50 or more fragments.) The conventional geologic and engineering approach is presented in the following graphic. **NOTE:** NRCS does *not* quantify **Sphericity**. It is included here for completeness and to show the **Fragment Roundness** range.

		<b>Roundness</b> <sup>1, 2</sup>					
		<b>Very Angular</b> 0.5	<b>Angular</b> 1.5	<b>Sub-angular</b> 2.5	<b>Sub-rounded</b> 3.5	<b>Rounded</b> 4.5	<b>Well Rounded</b> 5.5
<b>Sphericity</b>	Discooidal 0.5						
	Sub-discooidal 1.5						
	Spherical 2.5						
	Sub-prismatic 3.5						
	Prismatic 4.5						

<sup>1</sup> After Powers, 1953.

<sup>2</sup> Numerical values below *Roundness* and *Sphericity* headings are class midpoints (median rho values; Folk, 1955) used in statistical analysis.

**ROCK and OTHER FRAGMENTS - HARDNESS** (called **fragment\_hardness** in NASIS)—Describe the relative force required to crush the fragment. Use the same criteria and classes as the **Rupture Resistance for Blocks, Peds, and Clods - Cementation** column (p. 2–63); e.g., *Moderately Cemented* (exclude the *Noncemented* class).

### **ARTIFACTS (Human-derived)**

These are discrete, water-stable fragments of human origin (cultural byproducts) (called **Human\_artifacts** in NASIS). They are described separately from **Rock and Other Fragments** due to their unique properties and nongeologic origins and due to unique historical and cultural implications.

Describe **Kind, Quantity** (vol. percentage), **Roundness, Shape, Cohesion, Penetrability, Persistence, Safety**.

**ARTIFACTS - KIND**—Record the dominant types of human artifacts present by horizon/layer. (Used in NASIS primarily for % passing sieve calculation.) All fragments  $\geq 2$  mm.

Kind	
bitumen ( <i>asphalt</i> )	fly ash
boiler slag	glass
bottom ash	metal
brick	paper
cardboard	plasterboard
carpet	plastic
cloth	potsherd
coal combustion byproducts	rubber ( <i>tires, etc.</i> )
concrete ( <i>fragments</i> )	treated wood
debitage ( <i>stone tool flakes</i> )	untreated wood

**ARTIFACTS - QUANTITY**—Estimate the relative amount (volume %) of artifacts by horizon/layer. In NASIS, estimate a Representative Value (RV).

Quantity	Criteria
#	(volume percent)

**ARTIFACTS - ROUNDNESS**—Estimate the dominant extent of roundness of the artifacts by horizon/layer. (Refer to **Rock and Other Fragments - Roundness** graphic on p. 2-49.)

Roundness Class	Code
Angular	AN
Rounded	RO
Subangular	SA
Subrounded	SR
Very Angular	VA
Well Rounded	WR

**ARTIFACTS - SHAPE**—Describe the dominant form (shape) of the artifacts by horizon/layer.

Shape Class	Code	Criteria
Elongated	E	One dimension (length, width, or height) is 3X longer than either of the others.
Equidimensional	Q	Length, width, height are approximately the same.
Flat	F	One dimension is <1/3 that of either of the others, and one dimension is <3X that of the intermediate.
Irregular	I	Branching or convoluted form.

**ARTIFACTS - COHESION**—Describe the dominant relative fragment integrity.

Cohesion Class	Code	Criteria
Cohesive	C	Cannot be readily broken to <2 mm pieces.
Noncohesive	N	Easily broken to <2 mm pieces by hand or simple crushing.

**ARTIFACTS - PENETRABILITY**—Describe the prevalent relative ease of penetration of artifacts by external mechanical force by horizon/layer.

Penetrability Class	Code	Criteria
Nonpenetrable	N	Roots cannot penetrate through or between artifacts.
Penetrable	P	Roots can penetrate through or between artifacts.

**ARTIFACTS - PERSISTENCE**—Describe the dominant relative extent.

Persistence Class	Code	Criteria
Nonpersistent	N	Susceptible to relatively rapid weathering or decay (expected loss in <10 years).
Persistent	P	Expected to remain intact in soil for >10 years.

**ARTIFACTS - SAFETY**—Describe the dominant relative level of chemical safety of artifacts present.

Safety Class	Code	Criteria
Innocuous artifacts	IA	Harmless to living beings (e.g., brick, wood, glass, etc.).
Noxious artifacts	NA	Potentially harmful or destructive to living beings (e.g., batteries, garbage, petroleum products).

### ***(SOIL) STRUCTURE***

---










(Soil) structure is the naturally occurring arrangement of soil particles into aggregates that results from pedogenic processes. Record **Grade**, **Size**, and **Type**. For compound structure, list each size and type; e.g., *medium and coarse SBK parting to fine GR*. Lack of structure (structureless) has two end members: *massive (MA)* or *single grain (SG)*. A complete example is: *weak, fine, subangular blocky or 1, f, sbk*.

**(SOIL STRUCTURE) - TYPE** (formerly **Shape**)—Record the dominant type of ped, by layer; e.g., *granular* or *gr*. If a prevailing large shape readily breaks into smaller units, record as “(larger type) parting to (smaller type)”; e.g., *prismatic parting to subangular blocky*.


Type	Code		Criteria: definition
	Conv.	NASIS	
<b>NATURAL SOIL STRUCTURAL UNITS (pedogenic structure)</b>			
Granular	gr	GR	Small polyhedrals with curved or very irregular faces.
Angular Blocky	abk	ABK	Polyhedrals with faces that intersect at sharp angles (planes).
Subangular Blocky	sbk	SBK	Polyhedrals with subrounded and planar faces lacking sharp angles.
Lenticular	lp	LP	Overlapping, lens-shaped peds generally parallel to the soil surface that are thick at the center and taper toward the edges; formed by active or relict periglacial frost processes. Most common in soils with moderate to high water-holding capacity in moist conditions.
Platy	pl	PL	Flat and platelike units.
Wedge	wg	WEG	Elliptical, interlocking lenses that terminate in acute angles, bounded by slickensides; not limited to vertic materials.
Prismatic	pr	PR	Vertically elongated units; flat tops.
Columnar	cpr	COL	Vertically elongated units with rounded tops that commonly are "bleached."
<b>STRUCTURELESS</b>			
Single Grain	sg	SGR	No structural units; entirely noncoherent; e.g., loose sand.
Massive	m	MA	No structural units; material is a coherent mass (not necessarily cemented).
<b>ARTIFICIAL EARTHY FRAGMENTS OR CLODS<sup>1</sup> (nonpedogenic structure)</b>			
Cloddy <sup>1</sup>	—	CDY	Irregular blocks created by artificial disturbance; e.g., tillage or compaction.

<sup>1</sup> Used only to describe oversized, "artificial" earthy units that are not pedogenically derived soil structural units; e.g., the direct result of mechanical manipulation; use **Blocky Structure Size** criteria.

## Examples of Soil Structure Types

<p><b>Granular</b></p>  <p>(Soil aggregates)</p>	<p><b>Blocky</b></p> <p>(Subangular)      (Angular)</p> 
<p><b>Lenticular</b></p> 	<p><b>Platy</b></p> 
<p><b>Wedge</b></p> 	<div style="display: flex; justify-content: space-around;"> <div style="width: 45%;"> <p><b>Prismatic</b></p>  </div> <div style="width: 45%;"> <p><b>Columnar</b></p>  </div> </div>
<p><b>Structureless Types</b></p>	
<p><b>Single Grain</b></p>  <p>(Loose mineral/rock grains)</p>	<p><b>Massive</b></p>  <p>(Continuous, unconsolidated mass)</p>

### Example of Wedge Structure, Gilgai Microfeature, and Microrelief





**(SOIL STRUCTURE) - GRADE—**

Grade	Code	Criteria
Structureless	0	No discrete units observable in place or in hand sample.
Weak	1	Units are barely observable in place or in a hand sample.
Moderate	2	Units well formed and evident in place or in a hand sample.
Strong	3	Units are distinct in place (undisturbed soil) and separate cleanly when disturbed.

**(SOIL STRUCTURE) - SIZE—**

Size Class	Code		Criteria: structural unit size <sup>1</sup> (mm)		
	Conv.	NASIS	Granular, Platy <sup>2</sup> , (Thickness)	Columnar, Prismatic, Wedge <sup>3</sup> (Diameter)	Angular & Subangular Blocky and Lenticular (Diameter)
Very Fine (Very Thin) <sup>2</sup>	vf (vn)	VF (VN)	< 1	< 10	< 5
Fine (Thin) <sup>2</sup>	f (tn)	F (TN)	1 to < 2	10 to < 20	5 to < 10
Medium (Medium)	m (m)	M (M)	2 to < 5	20 to < 50	10 to < 20
Coarse (Thick) <sup>2</sup>	co (tk)	CO (TK)	5 to < 10	50 to <100	20 to < 50
Very Coarse (Very Thick) <sup>2</sup>	vc (vk)	VC (VK)	≥ 10	100 to <500	≥ 50
Extremely Coarse	ec (—)	EC (—)	—	≥500	—


<sup>1</sup> Size limits always denote the *smallest* dimension of the structural units.

<sup>2</sup> For platy structures only, substitute *Thin* for *Fine* and *Thick* for *Coarse* in the Size Class names.

<sup>3</sup> Wedge structure is generally associated with Vertisols (for which it is a requirement) or related soils (e.g., "Vertic" subgroups) with high amounts of smectitic clays.


# Granular

## Codes




# Platy

## Codes




# Lenticular

## Codes





# Angular and Subangular Blocky



# Prismatic and Columnar

## Codes






## CONSISTENCE

Consistence is the degree and kind of cohesion and adhesion that soil exhibits and/or the resistance of soil to deformation or rupture under an applied stress. Soil-water state strongly influences consistence. Field evaluations of consistence include: **Rupture Resistance** (Blocks, Peds, and Clods; or Surface Crusts and Plates), **Manner of Failure** (Brittleness, Fluidity, Smeariness), **Stickiness**, **Plasticity**, and **Penetration Resistance**. Historically, consistence applied to dry, moist, or wet soil as observed in the field. Wet consistence evaluated stickiness and plasticity. **Rupture Resistance** now applies to dry soils and to soils in a water state from moist through wet. **Stickiness** and **Plasticity** of soil are independent evaluations.

**RUPTURE RESISTANCE**—A measure of the strength of soil to withstand an applied stress. Separate estimates of **Rupture Resistance** are made for **Blocks/Peds/Clods** and for **Surface Crusts and Plates** of soil. Block-shaped specimens should be approximately 2.8 cm across. If 2.8-cm cubes (e.g.,  $\approx 2.5$ -3.1 cm, or 1 inch) are not obtainable, use the following equation and the table below to calculate the stress at failure:  $[(2.8 \text{ cm}/\text{cube length cm})^2 \times \text{estimated stress (N) at failure}]$ ; e.g., for a 5.6-cm cube  $[(2.8/5.6)^2 \times 20 \text{ N}] = 5 \text{ N} \Rightarrow \text{Soft Class}$ . Plate-shaped specimens (surface crusts or platy structure) should be approximately 1.0-1.5 cm long by 0.5 cm thick (or the thickness of occurrence, if <0.5 cm thick).





## RUPTURE RESISTANCE FOR:

**Blocks, Peds, and Clods**—Estimate the class by the force required to rupture (break) a soil unit. Select the column for the appropriate soil water state (*dry vs. moist*) and/or the *Cementation* column, if applicable.

Dry <sup>1</sup>		Moist <sup>1</sup>		Cementation <sup>2</sup>		Specimen Fails Under
Class	Code <sup>3</sup>	Class	Code <sup>3</sup>	Class	Code <sup>3</sup>	
Loose	L d(lo)	Loose	L m(lo)	[Not Applicable]		[Intact specimen not obtainable]
Soft	S d(so)	Very Friable	VFR m(vfr)	Non-cemented	NC	Very slight force between fingers. <8 N
Slightly Hard	SH d(sh)	Friable	FR m(fr)	Extremely Weakly Cemented	EW	Slight force between fingers. 8 to <20 N
Mod. Hard	MH d(h)	Firm	FI m(fi)	Very Weakly Cemented	VW	Moderate force between fingers. 20 to <40 N
Hard	HA d(vh)	Very Firm	VFI m(vfi)	Weakly Cemented	W c(w)	Strong force between fingers. 40 to <80 N
Very Hard	VH d(vh)	Extr. Firm	EF m(efi)	Moderately Cemented	M	Moderate force between hands. 80 to <160 N
Extr. Hard	EH d(eh)	Slightly Rigid	SR m(efi)	Strongly Cemented	ST c(s)	Foot pressure by full body weight. 160 to <800 N
Rigid	R d(eh)	Rigid	R m(efi)	Very Strongly Cemented	VS	Blow of <3 J but not body weight. 800 N to <3 J
Very Rigid	VR d(eh)	Very Rigid	VR m(efi)	Indurated	I c(I)	Blow of ≥3 J (3 J = 2 kg weight dropped 15 cm)

<sup>1</sup> Dry Rupture Resistance column applies to soils that are moderately dry or drier (*Moderately Dry* and *Very Dry Soil Water State* subclasses). Moist column applies to soils that are slightly dry or wetter (*Slightly Dry* through *Satiated Soil Water State* subclasses) (Soil Survey Division Staff, 1993, p. 91).

<sup>2</sup> This is not an immediate field test; specimen must first be air dried and then submerged in water for a minimum of 1 hour prior to test; collapse/disaggregation="slaking" (Soil Survey Division Staff, 1993, p. 173).

<sup>3</sup> Codes in parentheses (e.g., d(lo); Soil Survey Staff, 1951) are obsolete.

### Surface Crust and Plates—

Class (air dried)	Code	Force <sup>1</sup> (Newtons)
Extremely Weak	EW	Not Obtainable
Very Weak	VW	Removable, < 1N
Weak	W	1 to < 3N
Moderate	M	3 to < 8N
Moderately Strong	MS	8 to < 20N
Strong	S	20 to < 40N
Very Strong	VS	40 to < 80N
Extremely Strong	ES	≥ 80N

<sup>1</sup> For operational criteria (field estimates of force [N]), use the *Fails Under* column in the "Rupture Resistance for Blocks, Peds, Clods" table.

**CEMENTING AGENTS** (called **rupture\_resist\_cem\_agent** in NASIS)—Record kind of cementing agent, if present.

Kind	Code <sup>1</sup>
carbonates	K
gypsum <sup>2</sup>	G
humus	H
iron	I
silica (SiO <sub>2</sub> )	S

<sup>1</sup> Conventional codes traditionally consist of the entire material name or its chemical symbols; e.g., *silica* or *SiO<sub>2</sub>*. Consequently, the Conv. code column would be redundant and is not shown in this table.

<sup>2</sup> Gypsum is not a true cement but functionally behaves as such.

**MANNER OF FAILURE**—The rate of change and the physical condition soil attains when subjected to compression. Samples are moist or wetter.

<b>Failure Class</b>	<b>Code</b>	<b>Criteria: related field operation</b>
<b>BRITTLENESS</b>		<b>Use a 3-cm block (press between thumb and forefinger)</b>
Brittle	BR	Ruptures abruptly ("pops" or shatters).
Semi-deformable	SD	Rupture occurs before compression to $<1/2$ original thickness.
Deformable	DF	Rupture occurs after compression to $\geq 1/2$ original thickness.
<b>FLUIDITY <sup>1</sup></b>		<b>Use a palmful of soil (squeeze in hand)</b>
Nonfluid	NF	After full compression, no soil flows through the fingers. n value = 0
Slightly Fluid	SF	After full compression is exerted, some soil flows through fingers; most remains in the palm. n value $>0$ to $<0.7$
Mod. Fluid	MF	After full pressure is exerted, most soil flows through fingers; some remains in the palm. n value $>0.7$ to $<1.0$
Very Fluid	VF	Under very gentle pressure, most soil flows through the fingers as a slightly viscous fluid; very little or no residue remains in the palm of the hand. n value $>1.0$
<b>SMEARINESS</b>		<b>Use a 3-cm block (press between thumb and forefinger)</b>
Nonsmeary <sup>2</sup>	NS	At failure, the sample does not change abruptly to fluid, fingers do not skid, no smearing occurs.
Weakly Smeary <sup>2</sup>	WS	At failure, the sample changes abruptly to fluid, fingers skid, soil smears, little or no water remains on fingers.
Moderately Smeary <sup>2</sup>	MS	At failure, the sample changes abruptly to fluid, fingers skid, soil smears, some water remains on fingers.
Strongly Smeary <sup>2</sup>	SM	At failure, the sample abruptly changes to fluid, fingers skid, soil smears and is slippery, water is easily seen on fingers.

<sup>1</sup> See additional comments on fluidity under Subaqueous Soils (p. 2–105).

<sup>2</sup> *Smeariness* failure classes are used dominantly with materials displaying andic soil properties (and some spodic materials).

**STICKINESS**—The capacity of soil to adhere to other objects. Stickiness is estimated at the moisture content that displays the greatest adherence when pressed between thumb and forefinger.


Stickiness Class	Code		Criteria: work moistened soil between thumb and forefinger
	Conv.	NASIS	
Nonsticky	(w) so	SO	Little or no soil adheres to fingers after release of pressure.
Slightly Sticky	(w) ss	SS	Soil adheres to both fingers after release of pressure. Soil stretches little on separation of fingers.
Moderately Sticky <sup>1</sup>	(w) s	MS	Soil adheres to both fingers after release of pressure. Soil stretches some on separation of fingers.
Very Sticky	(w) vs	VS	Soil adheres firmly to both fingers after release of pressure. Soil stretches greatly upon separation of fingers.

<sup>1</sup> Historically, the *Moderately Sticky* class was simply called *Sticky*.

**PLASTICITY**—The degree to which “puddled” or reworked soil can be permanently deformed without rupturing. The evaluation is made by forming a roll (wire) of soil at a water content where the maximum plasticity is expressed.

Plasticity Class	Code		Criteria: make a roll of soil 4 cm long
	Conv.	NASIS	
Nonplastic	(w) po	PO	Will not form a roll 6 mm in diameter, or if a roll is formed, it can’t support itself if held on end.
Slightly Plastic	(w) ps	SP	6 mm diameter roll supports itself; 4 mm diameter roll does not.
Moderately Plastic <sup>1</sup>	(w) p	MP	4 mm diameter roll supports itself; 2 mm diameter roll does not.
Very Plastic	(w) vp	VP	2 mm diameter roll supports its weight.

<sup>1</sup> Historically, the *Moderately Plastic* class was simply called *Plastic*.



**PENETRATION RESISTANCE**—The ability of soil in a confined (field) state to resist penetration by a rigid object of specified size. A pocket penetrometer (Soil-Test Model CL-700) with a rod diameter of 6.4 mm (area 20.10 mm<sup>2</sup>) and insertion distance of 6.4 mm (note line on rod) is used for the determination. An average of five or more measurements should be used to obtain a value for penetration resistance.

**NOTE:** The pocket penetrometer has a scale of 0.25 to 4.5 tons/ft<sup>2</sup> (tons/ft<sup>2</sup>  $\approx$  kg/cm<sup>2</sup>). The penetrometer does *not* directly measure penetration resistance. The penetrometer scale is correlated to and gives a field estimate of *unconfined compressive strength* of soil as measured with a Tri-Axial Shear device. The table below converts the scale reading on the pocket penetrometer to penetration resistance in MPa. Penetrometer readings are dependent on the spring type used. Springs of varying strength are needed to span the range of penetration resistance found in soil.

Penetrometer Scale Reading	Spring Type <sup>1, 2, 3</sup>			
	Original MPa	Lee MPa	Jones 11 MPa	Jones 323 MPa
0.25	<b>0.32 L</b>	<b>0.06 VL</b>	<b>1.00 M</b>	<b>3.15 H</b>
0.75	0.60	<b>0.13 L</b>	1.76	4.20
1.00	0.74	0.17	<b>2.14 H</b>	4.73
1.50	<b>1.02 M</b>	0.24	2.90	5.78
2.75	1.72	0.42	4.80	<b>8.40 EH</b>
3.50	<b>2.14 H</b>	0.53	—	—

<sup>1</sup> On wet or “soft” soils, a larger “foot” may be needed (Soil Survey Division Staff, 1993).

<sup>2</sup> Each bolded value highlights the force associated with a rounded value on the penetrometer scale that is closest to a *Penetration Resistance Class* boundary. The bolded letter represents the *Penetration Resistance Class* from the following table (e.g., **M** indicates the *Moderate* class).

<sup>3</sup> Each spring type spans only a part of the range of penetration resistance possible in soils; various springs are needed to span all *Penetration Resistance Classes*.

**Penetration Resistance Class** (called **Penetration Resistance** in NASIS)—Record the appropriate class, by horizon or layer, based on the average value of five or more measurements with a pocket penetrometer.

<b>Penetration Resistance Class</b>	<b>Code</b>	<b>Criteria: Penetration Resistance (MPa)</b>
Extremely Low	EL	< 0.01
Very Low	VL	0.01 to < 0.1
Low	L	0.1 to < 1
Moderate	M	1 to < 2
High	H	2 to < 4
Very High	VH	4 to < 8
Extremely High	EH	≥ 8

**PENETRATION ORIENTATION**—Record the orientation of the pocket penetrometer used to determine the **Penetration Resistance Class**.

<b>Orientation</b>	<b>Code</b>	<b>Criteria</b>
Horizontal	H	Oriented perpendicular to a vertical pit face
Vertical <sup>1</sup>	V	Oriented perpendicular to the ground surface

<sup>1</sup> The conventional (preferred) orientation.

**EXCAVATION DIFFICULTY**—The relative force or energy required to dig soil out of place. Describe the **Excavation Difficulty Class** and the moisture condition (*moist* or *dry*, but not *wet*); use the “(Soil) Water State” table; e.g., *moderate, moist* or *M, M*. Estimates can be made for either the most limiting layer or for each horizon.

<b>Class</b>	<b>Code</b>	<b>Criteria</b>
Low	L	Excavation by tile spade requires arm pressure only; impact energy or foot pressure is not needed.
Moderate	M	Excavation by tile spade requires impact energy or foot pressure; arm pressure is insufficient.
High	H	Excavation by tile spade is difficult but easily done by pick using over-the-head swing.
Very High	VH	Excavation by pick with over-the-head swing is moderately to markedly difficult. Backhoe excavation by a 50- to 80-hp tractor can be made in a moderate time.
Extremely High	EH	Excavation via pick is nearly impossible. Backhoe excavation by a 50- to 80-hp tractor cannot be made in a reasonable time.

## ROOTS

Record the **Quantity**, **Size**, and **Location** of roots in each horizon. **NOTE:** Describe **Pores** using the same **Quantity** and **Size** classes and criteria as those for **Roots** (use the combined tables). A complete example for roots is: *Many, fine, roots In mat at top of horizon or 3, f (roots), M.*

**ROOTS (and PORES) - QUANTITY**—Describe the quantity (number) of roots for each size class in a horizontal plane. (**NOTE:** Typically, this is done across a vertical plane, such as a pit face.) Record the average quantity from three to five representative unit areas. **CAUTION:** The unit area that is evaluated varies with the *Size Class* of the roots being considered. Use the appropriate unit area stated in the *Soil Area Assessed* column of the "Size (Roots and Pores)" table (also see following graphic). In NASIS, record the actual number (#) of roots/unit area (NASIS then assigns the appropriate class). Use class names in narrative description.

Quantity Class <sup>1</sup>	Code		Average Count <sup>2</sup> (per assessed area)
	Conv.	NASIS	
Few	1	#	<1 per area
Very Few <sup>1</sup>	—	#	<0.2 per area
Moderately Few <sup>1</sup>	—	#	0.2 to <1 per area
Common	2	#	1 to <5 per area
Many	3	#	≥5 per area

<sup>1</sup> The *Very Few* and *Moderately Few* subclasses can be used for roots (optional) but do not apply to pores.

<sup>2</sup> The applicable area for appraisal varies with the size of roots or pores. Use the appropriate area stated in the *Soil Area Assessed* column of the "Size (Roots and Pores)" table or use the following graphic.


**ROOTS (and PORES) - SIZE**—(See the following graphic for size.)

Size Class	Code		Diameter	Soil Area Assessed <sup>1</sup>
	Conv.	NASIS		
Very Fine	vf	VF	< 1 mm	1 cm <sup>2</sup>
Fine	f	F	1 to < 2 mm	1 cm <sup>2</sup>
Medium	m	M	2 to < 5 mm	1 dm <sup>2</sup>
Coarse	co	C	5 to <10 mm	1 dm <sup>2</sup>
Very Coarse	vc	VC	≥10 mm	1 m <sup>2</sup>


<sup>1</sup> One dm<sup>2</sup>=a square that is 10 cm on a side, or 100 cm<sup>2</sup>.



**ROOTS (and PORES) - QUANTITY**—Soil area to be assessed.



## Root and Pore Size Classes



**ROOTS - LOCATION (Roots only)**—Identify where roots occur.

Location	Code
Between peds	P
In cracks	C
In mat at top of horizon <sup>1</sup>	M
Matted around rock fragments	R
Throughout	T

<sup>1</sup> Describing a root mat at the top of a horizon rather than at the bottom or within the horizon flags the horizon that restricts root growth.

## **PORES (DISCUSSION)**

---

Pores are the air- or water-filled voids in soil. Historically, description of soil pores, called “nonmatrix” pores in the *Soil Survey Manual* (Soil Survey Division Staff, 1993), excluded interstructural voids, cracks, and, in some schemes, interstitial pores. *Interstructural voids* (i.e., the subplanar fractures between pedis; also called interpedal or structural faces/planes), which can be inferred from soil structure descriptions, are not recorded directly. *Cracks* can be assessed independently (Soil Survey Division Staff, 1993). *Interstitial pores* (i.e., visible, primary packing voids) may be visually estimated, especially for fragmental soils, or can be inferred from soil porosity, bulk density, and particle-size distribution. Clearly, one cannot assess the smallest interstitial pores (e.g., <0.05 mm) in the field. Field observations are limited to those that can be seen through a 10X hand lens or larger. Field estimates of interstitial pores are considered to be somewhat tenuous but still useful.

## **PORES**

---

Describe the **Quantity** and **Size** of pores for each size class, by horizon, in a horizontal plane. (**NOTE:** Typically, this is actually assessed on a vertical face.) Description of soil pore **Shape** and **Vertical Continuity** is optional. A complete example for pores is: *common, medium, tubular pores, throughout or c, m, TU (pores), T.*

**PORES - QUANTITY**—See and use **Quantity (Roots and Pores)**.

**PORES - SIZE**—See and use **Size (Roots and Pores)**.


**PORES - SHAPE (or Type)**—Record the dominant form (or “type”) of pores discernible with a 10X hand lens and by the unaided eye. (See following graphic.)

<b>Description</b>	<b>Code</b>	<b>Criteria</b>
<b>SOIL PORES</b> <sup>1</sup>		
Dendritic Tubular	DT	Cylindrical, elongated, branching voids; e.g., empty root channels.
Irregular	IG	Nonconnected cavities, chambers; e.g., vughs; various shapes.
Tubular	TU	Cylindrical and elongated voids; e.g., worm tunnels.

Description	Code	Criteria
Vesicular	VE	Ovoid to spherical voids; e.g., solidified pseudomorphs of entrapped gas bubbles concentrated below a crust; most common in arid and semiarid environments.
<b>PRIMARY PACKING VOIDS <sup>2</sup></b>		
Interstitial	IR	Voids between sand grains or rock frags.

<sup>1</sup> Also called "Nonmatrix Pores" (Soil Survey Division Staff, 1993).

<sup>2</sup> *Primary Packing Voids* include a continuum of sizes. As used here, they have a minimum size that is defined as pores that are visible with a 10X hand lens. *Primary Packing Voids*: also called "Matrix Pores" (Soil Survey Division Staff, 1993).




**PORES - VERTICAL CONTINUITY**—The average vertical distance through which the minimum pore diameter exceeds 0.5 mm. Soil must be moist or wetter.

Class	Code		Criteria: vertical distance
	Conv.	NASIS	
Low	—	L	< 1 cm
Moderate	—	M	1 to < 10 cm
High	—	H	≥ 10 cm

## CRACKS

Cracks (also called “Extra-Structural Cracks”; Soil Survey Division Staff, 1993) are fissures other than those attributed to soil structure. Cracks are commonly vertical, subplanar, and polygonal and are the result of desiccation, dewatering, or consolidation of earthy material. Cracks are much longer and can be much wider than planes that surround soil structural units, such as prisms and columns. Cracks are key to preferential flow, also called “bypass flow” (Bouma et al., 1982), and are a primary cause of temporal (transient) changes in ponded infiltration and hydraulic conductivity in soils (Soil Survey Division Staff, 1993). Cracks are primarily associated with, but not restricted to, clayey soils and are most pronounced in high shrink-swell soils (high COLE value). Record the **Relative Frequency** (estimated average number per m<sup>2</sup>), **Depth** (average), and **Kind**. A complete example is: *3, 25 cm deep, reversible trans-horizon cracks*.



**CRACKS - KIND**—Identify the dominant types of fissures.

Kind	Code <sup>1</sup>	General Description
<b><i>CRUST-RELATED CRACKS</i></b> <sup>2</sup> ( <i>shallow, vertical cracks related to crusts; derived from raindrop-splash and soil puddling followed by dewatering/consolidation and desiccation</i> )		
Reversible Crust-Related Cracks <sup>3</sup>	RCR	Very shallow (e.g., 0.1-0.5 cm); very transient (generally persist less than a few weeks); formed by drying from surface down; minimal seasonal influence on ponded infiltration (e.g., raindrop crust cracks).
Irreversible Crust-Related Cracks <sup>4</sup>	ICR	Shallow (e.g., 0.5-2 cm); seasonally transient (not present year-round nor every year); minor influence on ponded infiltration (e.g., freeze-thaw crust and associated cracks).
<b><i>TRANS-HORIZON CRACKS</i></b> <sup>5</sup> ( <i>deep, vertical cracks that commonly extend across more than one horizon and may extend to the surface; derived from wetting and drying or original dewatering and consolidation of parent material</i> )		
Reversible Trans-Horizon Cracks <sup>6</sup>	RTH	Transient (commonly seasonal; close when rewetted); large influence on ponded infiltration and $K_{sat}$ ; formed by wetting and drying of soil (e.g., Vertisols, vertic subgroups).
Irreversible Trans-Horizon Cracks <sup>7</sup>	ITH	Permanent (persist year-round; see <i>Soil Taxonomy</i> ), large influence on ponded infiltration and $K_{sat}$ (e.g., extremely coarse subsurface fissures within glacial till; drained polder cracks).

<sup>1</sup> No conventional codes; use entire term. NASIS codes are shown.

<sup>2</sup> Called "Surface-Initiated Cracks" (Soil Survey Division Staff, 1993).


<sup>3</sup> Called "Surface-Initiated Reversible Cracks" (Soil Survey Division Staff, 1993).

<sup>4</sup> Called "Surface-Initiated Irreversible Cracks" (Soil Survey Division Staff, 1993).

<sup>5</sup> Also called "Subsurface-Initiated Cracks" (Soil Survey Division Staff, 1993).

<sup>6</sup> Called "Subsurface-Initiated Reversible Cracks" (Soil Survey Division Staff, 1993).

<sup>7</sup> Called “Subsurface-Initiated Irreversible Cracks” (Soil Survey Division Staff, 1993).



**CRACKS - DEPTH**—Record the **Average Apparent Depth** (also called a “depth index value” in the *Soil Survey Manual* [Soil Survey Division Staff, 1993]), measured from the surface, as determined by the wire-insertion method ( $\approx 2$  mm diameter wire). **NOTE:** This method commonly gives a standard but conservative measure of the actual fracture depth. Do not record this data element for cracks that are not open to the surface. Depth (and apparent vertical length) of subsurface cracks can be inferred from the *Horizon Depth* column of layers exhibiting subsurface cracks.

**CRACKS - RELATIVE FREQUENCY**—Record the **Average Number of Cracks**, per meter, across the surface or **Lateral Frequency** across a soil profile as determined with a line-intercept method. This data element cannot be assessed from cores or push tube samples.

## **SOIL CRUSTS (DISCUSSION)**

C. Franks, R. Grossman, and P. Schoeneberger, NRCS, Lincoln, NE

A soil crust is a thin (i.e., <1 cm up to 10 cm thick) surface layer of soil particles bound together by living organisms and/or by minerals into a horizontal “mat” or small polygonal plates. Soil crusts form at the soil surface and have different physical and/or chemical characteristics than the underlying soil material. Typically, soil crusts change the infiltration rate of the mineral soil and stabilize loose soil particles and aggregates. There are two general categories of soil crusts: (I) biological crusts and (II) mineral crusts.

(I) **Biological Crust** (also called *biotic*, *cryptogamic*, *microbiotic*, or *microphytic* crust): a thin, biotically dominated surface layer or

mat formed most commonly by cyanobacteria (blue green algae), green and brown algae, mosses, and/or lichens (NRCS, 1997; NRCS, 2001) that forms in or on the soil surface. Various types of microbiotic crusts have been recognized based on the biological communities of which they are composed (no prevailing consensus on types of biological crusts, at present).

(II) **Mineral Crust** (also called *abiotic*, *nonbiotic*, or *nonmicrobiotic* crust): a thin surface layer composed of reversibly bonded soil particles or secondary mineral crystals, sometimes laminated, that is *not* physically dominated by a microbiotic "mat."

1. **Chemical Crust** (e.g., salt incrustations): a thin surface layer that is dominated by macro- or microcrystalline evaporites of halite ( $\text{NaCl}$ ),  $\text{MgSO}_4$ , mirabilite ( $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ ), thenardite ( $\text{Na}_2\text{SO}_4$ ), epsomite ( $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ), hexahydrite ( $\text{MgSO}_4 \cdot 6\text{H}_2\text{O}$ ), bloedite ( $\text{Na}_2\text{Mg}(\text{SO}_4)_2 \cdot 4\text{H}_2\text{O}$ ), konyaite ( $\text{Na}_2\text{Mg}(\text{SO}_4)_2 \cdot 5\text{H}_2\text{O}$ ), loewite ( $\text{Na}_{12}\text{Mg}_7(\text{SO}_4)_{13} \cdot 15\text{H}_2\text{O}$ ), gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) (Singer and Warrington, 1992; Doner and Lynn, 1989), or other minerals. Other surficial mineral incrustations (e.g., from acid mine drainage or other sources) are included within this group.
2. **Physical Crust**: a physically reconstituted, reaggregated, or reorganized surface layer composed predominantly of primary mineral particles.
  - a). **Raindrop Impact Crust** (also called a *structural* crust): a thin layer that forms as a result of *raindrop impact*, which causes the clay in the soil to disperse, and subsequently hardens into a massive structureless or platy surface layer when it dries (Singer and Warrington, 1992).
  - b). **Depositional Crust** (also called a "*fluventic zone*"; Soil Survey Division Staff, 1993): a surface layer, commonly laminated and of variable thickness, consisting of small aggregates or primary mineral grains deposited by short-range runoff and subsequently dried (Singer and Warrington, 1992).
  - c). **Freeze-Thaw Crust** (Soil Survey Division Staff, 1993): a seasonal surface sediment layer 1 to 5 cm thick occurring on bare ground that has been disaggregated or puddled by radiant heating and cooling to produce freeze/thaw cycles while *Very Moist* or *Wet*. Commonly, the layer is composed of interlocking polygonal plates 5 to 20 cm in diameter, separated by cracks 1 to 2 cm wide that extend to the base of the crust and do not completely close upon wetting; Dry Rupture Resistance is  $\leq$  *Moderately Hard*.
  - d). **Vesicular Crust**: a surface soil layer or zone characterized by spherical or ovoid, *discontinuous* pores 0.5 to 2 mm in



diameter that are visible to the naked eye and make up a substantial portion of the matrix volume (i.e.,  $\geq 20\%$  cross-sectional area). These vesicles are believed to form when the pores between clay particles in platy soil structure are subjected to repeated wetting and drying. If soil aggregates become particularly unstable when they become saturated, air pressure may form small round voids (e.g., “bubbles”) that remain when the soil dries (Blackburn et al., 1975). Vesicular crusts occur primarily in arid and semiarid areas.

## SOIL CRUSTS

**Soil Crusts**—Record the presence of any surface crust. No entry implies that no crust is present. (In NASIS, crusts are included under **Pedoderm**.)

**Description**—Soil crusts can be identified and recorded by **Kind**. Additional suggested descriptors may include: **Rupture Resistance (Surface Crusts and Plates)**, **Porosity (Kind)**, **Size**, **Diameter**, **Thickness**, **Amount** (cross-sectional ground coverage), and **Color**.

### SOIL CRUSTS - KIND—

Kind	Code	Criteria
BIOLOGICAL CRUSTS	MC	biotically dominated surface “mat” of algae, lichens, mosses, etc.; also called biotic, cryptogamic, microbiotic, or microphytic crusts; slightly flexible when moist
MINERAL CRUSTS	MI	reversibly bonded primary, secondary mineral grains; not biotically dominated; stiff or rigid when moist or dry
Chemical Crusts	CC	evaporites (e.g., NaCl) or precipitates (e.g., CaCO <sub>3</sub> )
Physical Crusts	—	reorganized, reconstituted
raindrop impact crust	RC	dispersed, puddled, dried
depositional crust	DC	sediments of variable thickness
freeze-thaw crust	FC	bare ground, small polygons
vesicular crust/ zone	VC	substantial discontinuous, spherical or ovoid pores; e.g., 0.5 to 4 mm diameter

## SPECIAL FEATURES

Record the **Kind** and **Area (%) Occupied**. Describe the special soil feature by kind and estimate the cross-sectional area (%) of the horizon that the feature occupies; e.g., *lamellae*, 15%.

**SPECIAL FEATURES - KIND** [Called **Horizon Feature Kind** in NASIS]—Identify the kind of special soil feature.

Kind	Code <sup>1</sup>	Criteria
desert pavement <sup>2</sup>	DP	A natural concentration of closely packed and polished stones at the soil surface in a desert (may or may not be an erosional lag).
water repellent layer	HL	Either a surface or subsurface layer that repels water (e.g., dry organic materials; scorch layers in chaparral). See p. 7–14.
ice wedge cast	IC	A vertical, often trans-horizon, wedge-shaped or irregular form caused by infilling of a cavity as an ice wedge melts; commonly stratified.
krotovinas	KR	Filled faunal burrows.
lamellae <sup>3</sup>	LA	Thin (e.g., >0.5 cm), pedogenically formed plates or intermittent layers.
lamina	LN	Thin (e.g., <1 cm), geogenically deposited strata or layers of alternating texture (e.g., silt and fine sand or silt and clay).
stone line	SL	A natural concentration of rock fragments caused by water erosion or transport erosional lag (i.e., carpedolith).
tongues of albic material	E	Small areas or lobes of albic material that dip down (interfinger) more than 5 cm into nonalbic material.
tongues of argillic material	B	Small areas or lobes of argillic material that dip down (interfinger) more than 5 cm into nonargillic material.

<sup>1</sup> Conventional codes consist of the entire name; e.g., *Tongues of Albic Material*. Consequently, no *Conv. code* is shown.

<sup>2</sup> In NASIS, proposed to be moved to a new descriptor (data element) called **Pedoderm** (in NASIS 6.2) (Soil Survey Staff, 2012b).

<sup>3</sup> In NASIS, described under **Diagnostic Horizon or Property - Kind**.

**SPECIAL FEATURES - AREA (%) OCCUPIED**—Estimate the cross-sectional area (%) of the horizon that the feature occupies (see graphics, p. 7–1).

## ***SATURATED HYDRAULIC CONDUCTIVITY AND PERMEABILITY (DISCUSSION)***

---

Saturated hydraulic conductivity ( $K_{sat}$ ) is the single most scientifically valuable parameter for phenomena related to soil-water flow and transport.  $K_{sat}$  quantitatively defines a soil's capacity to transmit water. Traditionally, NRCS (formerly SCS) used the term "permeability" for water-flow phenomena and used *Permeability Classes* (PC), which have prescribed percolation rate ranges (originally inches/hr). The PC and associated percolation rates are commonly mistaken to be  $K_{sat}$ . The confusion between  $K_{sat}$ , the term "permeability," and the *Permeability Classes* arises for several reasons (Wysocki et al., 2002). A primary reason is that the term "permeability" has three meanings in soil science.

- 1) "Permeability" in a qualitative sense describes a soil's capacity to transmit fluids, including water, or gases. No quantitative measure is implied. For example, sandy soils are more "permeable" than clayey soils.
- 2) "Permeability" ( $k$ ) (Richards, 1952) (also known as intrinsic permeability) is an exclusive, quantitative porous material parameter controlled by pore geometry. In a stable porous material, ( $k$ ) is independent of the fluid. Permeability ( $k$ ) is the hydraulic conductivity ( $K$ ) times the fluid viscosity ( $\eta$ ) divided by the fluid density ( $\rho$ ) and the gravitational constant ( $g$ ) (**Eq. 1**). Permeability ( $k$ ) has area units (e.g.,  $m^2$ ).
- 3) "Permeability" is short for permeability coefficient, which is hydraulic conductivity ( $K$ ), or in saturated soil ( $K_{sat}$ ). The Darcy equation quantitatively defines (**Eq. 2**) hydraulic conductivity ( $K$ ) as the factor that relates flux ( $q$ ) to the hydraulic gradient ( $\Delta h/l$ ).  $K_{sat}$  depends upon both soil and fluid attributes. Measurement units for  $K_{sat}$  depend on the input units. With flux expressed as volume ( $cm^3$ ), head change ( $\Delta h$ ) as cm ( $cm\ H_2O/cm$ ), and length as cm, the  $K_{sat}$  units are length/time ( $cm/s$ ). Note that both variables  $q$  and  $K$  have units of length/time ( $cm/s$ ), but they are distinctly different entities. Flux ( $q$ ), when expressed as length/time, is an apparent rate that varies with  $\Delta h/l$ .  $K_{sat}$  is a proportionality factor that relates  $q$  to  $\Delta h/l$ .  $K_{sat}$  remains constant when the hydraulic gradient ( $\Delta h/l$ ) varies. It is a physical parameter, not a rate.

**Eq. 1**  $k = Kn/\rho g$ k = permeability (cm<sup>2</sup>)

K = hydraulic conductivity (cm/sec)

n = fluid (water) viscosity (dyne-sec/cm)

ρ = fluid (water) density (gm/cm<sup>3</sup>)g = gravitation acceleration (cm/sec<sup>2</sup>)**Eq. 2**  $q = V/At = -K(\Delta h/l)$ **Darcy's Equation** (one dimensional flow)

q = fluid flux (cm/s)

V = fluid volume (cm<sup>3</sup>)A = cross-sectional area (cm<sup>2</sup>)

t = time (s)

K = hydraulic conductivity (cm/s)

Δh = change in hydraulic head (cm)

l = length (cm)

The different permeability meanings have important distinctions that are not scientifically interchangeable. Most importantly, the intended meaning of "permeability" is *not* specifically discernible from written or verbal context alone. Meaning #1 carries no quantitative implications; meanings #2 and #3 have defined scientific applications. Uhland and O'Neal (1951) developed seven *Permeability Classes* (PC) from measurements on about 10,000 3-inch cores collected from 900 sites. They chose the original PC ranges such that each class represented an equal number of measured values from the sample population (Mason et al., 1957). Uhland and O'Neal (1951) measured discharge volume and calculated flux (q) as follows.

**Eq. 3**  $q = V/At$ V = fluid volume (in<sup>3</sup>)A = cross-sectional area (in<sup>2</sup>)

t = time (hr)

Uhland and O'Neal (1951) specifically noted that the calculated value was a "percolation rate" with units of inches hr<sup>-1</sup>. These percolation rates defined the *Permeability Class* ranges. The Uhland and O'Neal study did not calculate  $K_{sat}$  from Darcy's equation (Eq. 2). The study method employed both a falling and constant head phase during measurement; the hydraulic gradient (Δh/l) varied and was undefined. Darcy's equation requires a defined hydraulic gradient to solve for  $K_{sat}$ . The PC, therefore, are a set of soil "permeability" (meaning #1) classes arrayed by a method-specific percolation rate. To obtain an approximation of  $K_{sat}$  for the Uhland and O'Neal (1951) study, one can use the constant head hydraulic gradient (0.857 in) and the flux (V/At) as a general solution of Darcy's equation, which yields V/At times 0.857 =  $K_{sat}$ . This solution shows that PC percolation rates, at a minimum, exceed  $K_{sat}$  by

about 15%. Two errors result if percolation rates are used as a proxy for  $K_{sat}$ . One is added uncertainty when estimating a soil  $K_{sat}$ . Do you decrease the estimate by 15%? Secondly, measured  $K_{sat}$  values would not be equivalent to the PC percolation rates.  $K_{sat}$  is the scientific standard for soil-water flow calculations (Hillel, 1980), and there is great scientific merit in using  $K_{sat}$  over PC for soil-water flow interpretations.

Uhland and O'Neal (1951) also developed a set of field-observable properties to link the large number of unmeasured soils to a permeability class. The field properties included structure size and type, aggregate overlap, texture, pores, compaction, and clay mineralogy (O'Neal, 1952). NRCS soil scientists assigned soils to the permeability classes based on this characteristic set, or extrapolation from soils measured in the initial study. The original percolation rate ranges for the PC were altered (Soil Survey Staff, 1971) and an eighth class added. The PC have historic merit and are retained for selected uses.

To avoid the confounding difficulties inherent in the PC and the term "permeability," the Soil Survey Division Staff (1993) developed  $K_{sat}$  classes. To summarize:

- 1)  $K_{sat}$  and the percolation rates (that defined the *Permeability Classes*) are different physical parameters. Both  $K_{sat}$  and percolation rates are commonly expressed in units length/time, which presents a false equivalency.
- 2) Darcy's equation relates  $K_{sat}$  to PC percolation rates. Core percolation rates used in *Permeability Class* development exceed  $K_{sat}$  by a minimum 15%.  $K_{sat}$  is not a rate.
- 3) No simple transformation exists to reliably convert PC percolation rates to  $K_{sat}$ . Soils with slower percolation rates have a greater difference between  $K_{sat}$  than those with more rapid percolation rates.
- 4) To prevent confusion and avoid scientific inaccuracies, NRCS now emphasizes  $K_{sat}$  rather than the term "permeability" and  $K_{sat}$  classes rather than *Permeability Classes*.

## **SATURATED HYDRAULIC CONDUCTIVITY ( $K_{SAT}$ )**

**Saturated hydraulic conductivity ( $K_{sat}$ )** is the ease with which a saturated soil can transmit water through the pore space.  $K_{sat}$  is formally defined as the proportionality factor that relates water flow rate to the hydraulic gradient in Darcy's equation (see Discussion).  $K_{sat}$  is a measurable soil property, or it may be estimated from other properties (texture, structure, bulk density, etc.). Direct field  $K_{sat}$  measurement is possible with various devices (Amoozometer, Guelph Permeameter, double-ring infiltrometer). Multiple (e.g.,

≥5) measurement replications are needed on a horizon or layer to capture the natural variation.

Record an estimated **K<sub>sat</sub> class** or a measured K<sub>sat</sub> value for each horizon/layer. If measured, record the **Average K<sub>sat</sub>, Standard Deviation, Replication Number (n), and Method**. See NSSH, Exhibit 618.88 (Soil Survey Staff, 2012c) for guidelines for K<sub>sat</sub> Class estimation using texture and bulk density (see p. 7–10).

K <sub>sat</sub> Class	NASIS Code <sup>1</sup>	Criteria <sup>2</sup>		
		µm/s	cm/hr	in/hr
Very Low	#	< 0.01	< 0.0036	< 0.001417
Low	#	0.01 to < 0.1	0.00360 to < 0.036	0.001417 to < 0.01417
Mod. Low	#	0.1 to < 1.0	0.0360 to < 0.360	0.01417 to < 0.1417
Mod. High	#	1.0 to < 10	0.360 to < 3.60	0.1417 to < 1.417
High	#	10 to < 100	3.60 to < 36.0	1.417 to < 14.17
Very High	#	≥ 100	≥ 36.0	≥ 14.17

<sup>1</sup> For alternative units commonly used for these class boundaries (e.g., Standard International Units [Kg s/m<sup>3</sup>]), see the *Soil Survey Manual* (Soil Survey Division Staff, 1993, p. 107).

<sup>2</sup> To convert µm/sec to in/hr, multiply µm/sec by 0.1417; e.g., (100 µm/sec) x (0.1417)=14.17 in/hr. To convert in/hr to µm/sec, multiply by 7.0572.

## PERMEABILITY CLASSES

---

NRCS deemphasizes the use of *Permeability Classes*. Use  $K_{sat}$ . The *Permeability Classes* are listed here because of historic usage and because they are needed for selected soil interpretations.

Permeability Class	Code	Criteria: estimated in/hr <sup>1</sup>
Impermeable	IM	< 0.0015
Very Slow	VS	0.0015 to < 0.06
Slow	SL	0.06 to < 0.2
Moderately Slow	MS	0.2 to < 0.6
Moderate	MO	0.6 to < 2.0
Moderately Rapid	MR	2.0 to < 6.0
Rapid	RA	6.0 to < 20
Very Rapid	VR	≥ 20

<sup>1</sup> These class breaks were originally defined in English units and are retained here as no convenient metric equivalents are available.

## CHEMICAL RESPONSE

---

Chemical response is the reaction of a soil sample to an applied chemical solution or a measured chemical value. Responses are used to identify the presence or absence of certain materials, to obtain a rough assessment of the amount present, to measure the intensity of a chemical parameter (e.g., pH), or to identify the presence of chemical species (e.g.,  $Fe^{+2}$ ) in the soil.

**REACTION (pH)** - (Called **Field\_pH** in NASIS)—Record **pH** and **Method**; record the pH value to the precision limit of the method (e.g., to the nearest tenth). The preferred method is pH meter for 1:1 (water:soil). In NASIS, record **pH numerical value** and the method used (e.g., *pH 6.5; 1:1 water:soil*).

Descriptive Term	Code <sup>1</sup>	Criteria: pH range
Ultra Acid	#	<3.5
Extremely Acid	#	3.5 to 4.4
Very Strongly Acid	#	4.5 to 5.0
Strongly Acid	#	5.1 to 5.5
Moderately Acid	#	5.6 to 6.0
Slightly Acid	#	6.1 to 6.5
Neutral	#	6.6 to 7.3

Descriptive Term	Code <sup>1</sup>	Criteria: pH range
Slightly Alkaline	#	7.4 to 7.8
Moderately Alkaline	#	7.9 to 8.4
Strongly Alkaline	#	8.5 to 9.0
Very Strongly Alkaline	#	>9.0

<sup>1</sup> No codes; enter the measured value.

**pH METHOD** (called **ph\_determination\_method** in NASIS)—  
Record the method used to measure pH.

pH Method <sup>1</sup>		Code
<b>INDICATOR SOLUTION</b> <sup>2</sup>		(pH range) <sup>1</sup>
Bromocresol green	3.8 - 5.4	BG
Bromocresol purple	5.2 - 6.8	BP
Bromophenol blue	3.0 - 4.6	BL
Bromothymol blue	6.0 - 7.6	BB
Chlorophenol red	5.2 - 6.8	CHR
Cresol red	7.0 - 8.8	CR
Methyl red	4.8 - 6.0	MR
Phenol red	6.8 - 8.4	PR
Phenolphthalein	8.2 - 10.0	PT
Thymol blue	8.0 - 9.6	TB
<b>COMMERCIAL COLORIMETRIC KITS</b>		
Hellige-Truog (kit)		HT
Lamotte-Morgan (kit)		LM
Soil Test (kit)		ST
<b>pH METER</b> <sup>2</sup>		
pH meter 1:1 water <sup>3</sup>		M11
pH meter 1:2 water (0.01 M CaCl <sub>2</sub> ) <sup>3</sup>		C12
pH meter 1N KCl		M12
pH meter, saturated paste		MSD
<b>INDICATOR STRIPS</b> <sup>2</sup>		
indicator paper strip 1N NaF <sup>1</sup>		NF
pH indicator strip (unspecified) <sup>2</sup>		STR
(H)ydrion (unspecified; = hydrogen ion paper strip)		YD
pH unspecified <sup>2, 3</sup>		PHU

<sup>1</sup> Soil Survey Staff, 2009.

<sup>2</sup> The pH method options in NASIS, release 6.2.

<sup>3</sup> Preferred method.



**EFFERVESCENCE**—The gaseous response (seen as bubbles) of soil to applied HCl (carbonate test), H<sub>2</sub>O<sub>2</sub> (MnO<sub>2</sub> test), or other chemicals. Commonly, ≈1 N HCl is used for carbonate test. Apply the chemical to the soil matrix (for HCl, effervescence class refers only to the matrix; do not include carbonate masses, which are described separately as “Concentrations”). Record the observed response (**Effervescence Class**) and the **Chemical Agent** used. A complete example is: *Strongly Effervescent with 1N HCl; or ST, H2.* (**NOTE:** In NASIS, manganese effervescence [by H<sub>2</sub>O<sub>2</sub>] is handled in separate tables; called **MN\_Effervescence\_Agent** and **Mn\_Effervescence** classes; class codes and criteria are the same as those for **Effervescence Class**.)

**Effervescence - Class**—

Effervescence Class	Code	Criteria
Noneffervescent	NE	No bubbles form.
Very Slightly Effervescent	VS	Few bubbles form.
Slightly Effervescent	SL	Numerous bubbles form.
Strongly Effervescent	ST	Bubbles form a low foam.
Violently Effervescent	VE	Bubbles rapidly form a thick foam.

**Effervescence - Location** (obsolete in NASIS)—Use locations and codes from (**Ped and Void**) **Surface Features - Location**. (**NOTE:** The requirement to apply chemical agents [e.g., HCl] to the soil matrix makes many location choices invalid.)

**Effervescence - Chemical Agent** (In NASIS, the manganese chemical test agent [H<sub>2</sub>O<sub>2</sub>] is recorded in a separate table [**mn\_effervescence\_agent**].)

Effervescence Agent	Code	Criteria
HCl (unspecified) <sup>1</sup>	H1	Hydrochloric Acid: Concentration Unknown
HCl (1N) <sup>1, 2</sup>	H2	Hydrochloric Acid: Concentration=1 Normal
HCl (3N) <sup>1, 3</sup>	H3	Hydrochloric Acid: Concentration=3 Normal
HCl (6N) <sup>1, 4</sup>	H4	Hydrochloric Acid: Concentration=6 Normal
H <sub>2</sub> O <sub>2</sub> (unspecified) <sup>5, 6</sup>	P1	Hydrogen Peroxide: Concentration Unknown
H <sub>2</sub> O <sub>2</sub> <sup>5, 6</sup>	P2	Hydrogen Peroxide: Concentration 3-4%

- <sup>1</sup> Positive reaction indicates presence of carbonates (e.g., CaCO<sub>3</sub>).
- <sup>2</sup> The only HCl concentration used for the effervescence field test. **NOTE:** A (1N HCl) solution is made by combining 1 part concentrated (37%) HCl (which is widely available) with 11 parts distilled H<sub>2</sub>O.
- <sup>3</sup> Use 3N HCl to determine the Calcium Carbonate Equivalent test. It is not used for **Effervescence Class**. An approximately 3N HCl solution (10% HCl or 2.87N) is made by combining 6 parts 37% HCl (which is widely available) with 19 parts distilled H<sub>2</sub>O.
- <sup>4</sup> A 6N HCl solution is used to distinguish between calcium and dolomitic carbonates. Dolomite reaction is slower and less robust than CaCO<sub>3</sub> effervescence. A 6N HCl solution is made by combining 1 part concentrated (37%) HCl (which is widely available) with 1 part distilled H<sub>2</sub>O. Soil sample should be saturated in a spot plate and allowed to react for 1 to 2 minutes; froth=positive response.
- <sup>5</sup> Rapid reaction indicates presence of manganese oxides (e.g., MnO<sub>2</sub>). Not used to determine "Effervescence Class."
- <sup>6</sup> Under ambient conditions, Mn-oxides react rapidly whereas most organic matter reacts slowly with (3-4%) H<sub>2</sub>O<sub>2</sub>.

**REDUCED CONDITIONS** (called **Reaction to alpha-dipyridyl** in NASIS)—Record under "Notes" if evaluated.

Chemical Agent	Code	Criteria
$\alpha, \alpha$ -dipyridyl <sup>1</sup> (0.2% conc. <sup>3</sup> )	P	positive reaction <sup>2</sup> : red or pink color develops
	N	negative reaction: no color develops

<sup>1</sup> Commonly stated as "alpha-alpha dipyridyl."

<sup>2</sup> Positive reaction indicates presence of Fe<sup>+2</sup> (i.e., reduced conditions).

<sup>3</sup> Childs, 1981.

**Dipyridyl - Location**—Describe the location(s) where the chemical test was conducted (use "Concentrations - Location" table); e.g., *In the matrix (MAT)*.

**SALINITY CLASS (DISCUSSION)**—Soil salinity classes are based on electrical conductivity from a saturation paste extract. Gypsum (CaSO<sub>4</sub> • 2H<sub>2</sub>O) and salts more soluble than gypsum (e.g., Na, Mg, and Ca chlorides and sulfates) are the sole or major contributors to the saturated paste extract EC.

**NOTE:** Electrical conductivity may be measured at various soil solution extract ratios (e.g., 1:1, 1:2, 1:5). The resultant EC values are not directly comparable because of the dilution effect. The salinity standard is the saturated paste extract EC. To avoid confusion, saturated paste EC is commonly denoted as E<sub>ce</sub> and other extracts denoted by the dilution ratio (e.g., E<sub>C1:1</sub>).

In addition to solution extracts, field measures of EC exist (e.g., electromagnetic induction [EMI], salinity probes). These measurements obtain an EC value that depends on salinity, moisture content, mineralogy, and texture. Such EC measurements are *not* directly comparable to E<sub>ce</sub> or EC of any extract ratio. For example, the electromagnetic induction (EMI) EC is known as apparent EC, which is denoted as E<sub>ca</sub>.

**SALINITY CLASS**—Estimate the **Salinity Class**. If the electrical conductivity is measured, record the **EC Value** (in the “Notes” column). Salinity class is based on saturated paste extract EC.

Salinity Class	Code	Saturated Paste - E <sub>ce</sub> dS/m
Nonsaline	0	< 2
Very Slightly Saline	1	2 to < 4
Slightly Saline	2	4 to < 8
Moderately Saline	3	8 to < 16
Strongly Saline	4	≥ 16

**SODIUM ADSORPTION RATIO (SAR)**—A measure of ion equilibrium between sodium (Na) in solution and exchangeable Na adsorbed on the soil (Soil Survey Staff, 2011). It is applied to soil solution extracts and irrigation waters.  $SAR = [Na^+] / ([Ca^{+2}] + [Mg^{+2}]) / 2$ <sup>0.5</sup>, where the cation concentration is in milliequivalents per liter. As a field method, it is commonly determined with soil paste and an electronic wand.

## ODOR

---

**ODOR**—Record the **Kind** and relative **Intensity of odor (by horizon)** immediately after soil is exposed to air. The presence of an intense hydrogen sulfide odor ( $H_2S$ ; rotten egg) is commonly associated with a strongly anaerobic horizon where sulfate is reduced to sulfide (Fanning and Fanning, 1989).

Odor - Kind	Code	Criteria
None	N	No odor detected.
Petrochemical	P	Presence of gaseous or liquid gasoline, oil, creosote, etc.
Sulfurous	S	Presence of $H_2S$ (hydrogen sulfide); "rotten egg"; commonly associated with strongly reduced soil containing sulfides.

**Odor Intensity** (*proposed*)—Estimate and record the relative intensity of any odor present.

Odor Intensity	Code	Criteria: relative intensity of odor
Slight	SL	Odor is faint (e.g., only detected when sample is brought close to nose).
Moderate	MD	Odor is readily noticeable at arm's length as one handles the material (e.g., intermediate intensity); only detected as one starts to dig into the material.
Strong	ST	Odor is quite intense and readily detected before or immediately after the sample is exposed to air.

## MISCELLANEOUS FIELD NOTES

---

Use additional descriptors and sketches to capture and convey information and features with no existing data element. Record as freehand notes under **Miscellaneous Field Notes**.

## MINIMUM DATA SET (for a soil description)

---

Purpose, field logistics, habits, and soil materials all influence the specific properties necessary to "adequately" describe a given soil. However, some soil properties or features are so universally essential for interpretations or behavior prediction that they should always be recorded. These include: **Location, Horizon, Horizon**

**Depth, Horizon Boundary, Color, Redoximorphic Features, Texture, Structure, and Consistence.**

## ***PEDON DESCRIPTION DATA SHEET***

---

Over the decades, field data for soils have been documented in various ways. For many years soil descriptions were made on small blue cards (SCS-SOI-232 form: USDA-SCS, various versions, dates, and locations of issuance). Since the NRCS reorganization in 1995, some MLRA Soil Survey Regional Offices (MOs) and other groups have generated informal, locally tailored data sheets.

The following (blank) data sheet is provided as an option to record basic soil description information. This revised data sheet contains the most widely used soil descriptors (e.g., depth, color). Other descriptors (called data elements in NASIS) should be added as needed in blank boxes or in the **Miscellaneous Field Notes** box or in the **Notes** column. See p. 2-93.

## ***PEDON DESCRIPTION EXAMPLE***

---

A completed profile description data sheet is included to demonstrate recording soil information in the field (see p. 2-95).

Most field descriptions will likely be entered into an electronic database by the describer or must be deciphered by other scientists. Therefore, descriptions should use reasonably mnemonic abbreviations, standard codes, or a combination of these or be written in "longhand" (using complete words). The following profile description contains examples of all of these conventions.

Soil descriptions in soil survey reports, Official Soil Series Descriptions (OSDs), or other NRCS products should follow prescribed formats and descriptor sequences (i.e., NSSH, Part 614; Soil Survey Staff, 2012c).



# PEDON DESCRIPTION

PEDON ID#:

<b>Series or Component Name:</b>		<b>Map Unit Symbol:</b>	<b>Photo #:</b>	<b>Classification:</b>			<b>Soil Moist. Regime (Tax.):</b>		
<b>Describer(s):</b>	<b>Date:</b>	<b>Weather:</b>	<b>Temp.: Air:</b>		<b>Latitude:</b> °   '    " N		<b>Geodetic Datum:</b>		<b>Location:</b>
				<i>Soil:    Depth:</i>		<b>Longitude:</b> °   '    " W		Sec.       T.       R.	
<b>UTM:</b> <i>Zone:</i> <i>mE:</i> <i>mN:</i>	<b>Topo Quad:</b>		<b>Site ID:</b> <i>Yr:</i> <i>State:</i> <i>County:</i> <i>Pedon #:</i>		<b>Soil Survey Area:</b>	<b>MLRA/LRU:</b>	<b>Transect:</b> <i>ID:</i>		
								<i>Stop#:</i> <i>Interval:</i>	
<b>Landscape:</b>	<b>Landform:</b>	<b>Microfeature:</b>	<b>Anthro:</b>	<b>Elevation:</b>	<b>Aspect:</b>	<b>Slope (%):</b>	<b>Slope Complexity:</b>	<b>Slope Shape:</b> ( <i>Up &amp; Dn / Across</i> )	
<b>Hillslope Profile Position:</b>	<b>Geom. Component:</b>	<b>Microrelief:</b>	<b>Physio. Division:</b>	<b>Physio. Province:</b>	<b>Physio. Section:</b>	<b>State Physio. Area:</b>	<b>Local Physio. Area:</b>		
<b>Drainage:</b>	<b>Flooding:</b>	<b>Ponding:</b>	<b>Soil Moisture Status:</b>		<b>K<sub>sat</sub>:</b>			<b>Land Cover / Use:</b>	
<b>Parent Material:</b>		<b>Bedrock:</b> <i>Kind:</i> <i>Fract.:</i> <i>Hard.:</i> <i>Depth:</i>			<b>Lithostrat. Units:</b> <i>Group:</i> <i>Formation:</i> <i>Member:</i>				
<b>Erosion:</b> <i>Kind:</i> <i>Degree:</i>		<b>Surface Frag %:</b> <i>GR:</i> <i>CB:</i> <i>ST:</i> <i>BD:</i> <i>CN:</i> <i>FL:</i>			<b>P.S. Control Section:</b>		<i>Ave. Clay%:</i>	<i>Ave. Rock Frag %:</i>	
		<i>Kind:</i>			<i>Depth Range:</i>				
<b>Diagnostic Horz. / Prop.:</b>		<i>Kind:</i> <i>Depth:</i>							

VEGETATION:		
SYMBOL	COMMON NAME	% GD COVER

MISCELLANEOUS FIELD NOTES / SKETCH:											


Component Name:										Map Unit Symbol:				Date:				
Obser. Method	Horizon	Depth		Bnd	Matrix Color		Texture	Rock Frags			Structure			Consistence				Notes
		(in)	(cm)		Dry	Moist		Kind %	Rnd	Sz	Grade	Sz	Type	Dry	Mst	Stk	Pls	
1																		
2																		
3																		
4																		
5																		
6																		
7																		
8																		
9																		
10																		

	Redoximorphic Features								Concentrations								Ped / V. Surface Features				Roots			Pores			pH, method	Effer (agent)	Clay %	Sand %	Notes
	% Sz	Cn	Hd	Sp	Kd	Loc	Bd	Col	% Sz	Cn	Hd	Sp	Kd	Loc	Col	% Dst	Cont	Kd	Loc	Col	Qty	Sz	Loc	Qty	Sz	Shp					
1																															
2																															
3																															
4																															
5																															
6																															
7																															
8																															
9																															
10																															



<b>Series or Component Name:</b> Caveat Emptor		<b>Map Unit Symbol:</b> CaC	<b>Photo #:</b> 127A	<b>Classification:</b> fine, smectitic, mesic Typic Argiudoll			<b>Soil Moist. Regime (Tax.):</b> Udic (≈ 27" annual)	
<b>Describer(s):</b> PJS & DAW	<b>Date:</b> 10/12/2000	<b>Weather:</b> sunny	<b>Temp.: Air:</b> 78°F	<b>Latitude:</b> 40° 49' 10.0" N	<b>Geodetic Datum:</b> NAD '83	<b>Location:</b> NE 1/4, SW 1/4 Sec. 20 T. 10N R. 6E		
<b>UTM:</b> Zone: mE: mN:	<b>Topo Quad:</b> Emerald, NE 7.5 topo 1978	<b>Site ID:</b> Yr: S 2002	<b>State:</b> NE	<b>County:</b> 109	<b>Pedon #:</b> 006	<b>Soil Survey Area:</b> Lancaster Co.	<b>MLRA/LRU:</b> 106	<b>Transect:</b> ID: 2 Stop#: 6 Interval: 10m
<b>Landscape:</b> till plain	<b>Landform:</b> low hill	<b>Microfeature:</b> -----	<b>Anthro:</b> furrow	<b>Elevation:</b> 1240'	<b>Aspect:</b> 34°	<b>Slope (%):</b> 6%	<b>Slope Complexity:</b> simple	<b>Slope Shape:</b> (Up & Dn / Across) V V (convex, convex)
<b>Hillslope Profile Position:</b> shoulder	<b>Geom. Component:</b> nose slope	<b>Microrelief:</b> micro-low	<b>Physio. Division:</b> Interior Plains	<b>Physio. Province:</b> Central Lowland	<b>Physio. Section:</b> Dissected Till Plain	<b>State Physio. Area:</b> -----	<b>Local Physio. Area:</b> Middle Creek Basin	
<b>Drainage:</b> WD (Well Drained)	<b>Flooding:</b> none	<b>Ponding:</b> none	<b>Soil Moisture Status:</b> moist	<b>K<sub>sat</sub>:</b> low (ave. = 0.011 cm/hr; n=3; depth: 35-50 cm)			<b>Land Cover / Use:</b> Soybeans (CCG)	
<b>Parent Material:</b> loess, over pedisediment, over till		<b>Bedrock:</b> Kind: Fract.: Hard.: Depth: Sandstone (in adjacent gully) > 15 m		<b>Lithostrat. Units:</b> Group: Formation: Member: Peoria Loess, Gillman Canyon, unnamed pedisediment, unnamed till				
<b>Erosion:</b> Kind: Degree: R (Rill) 1	<b>Surface Frag %:</b> GR: CB: ST: BD: CN: FL: Kind: -- 0 --	<b>P.S. Control Section:</b> Ave. Clay%: Ave. Rock Frag %: Depth Range: 30-80 cm 37% < 1%						
<b>Diagnostic Horz. / Prop.:</b> Kind: Depth: mollic, 0-30 cm; argillic 30-90 cm								

VEGETATION:		
SYMBOL	COMMON NAME	% GD COVER
GLYCI	Soybean stubble	1%
DIGIT2	Crabgrass	5%



Component Name:							Map Unit Symbol:							Date:			
Obsr. Method	Horizon	Depth (in) (cm)	Bnd	Matrix Color		Texture	Rock Frags			Structure			Consistence				Notes
				Dry	Moist		Knd %	Rnd	Sz	Grade	Sz	Type	Dry	Mst	Stk	Pls	
1	LP*	Ap	0-20	Abrupt Smooth	10YR 4/2	10YR 3/1	silt loam (sil)	--	0	--	common, fine & med. granular	Slightly Hard	Friable	non-sticky	non-plastic	Obsrv. Meth.: LP, 4 x 300 m	
2	LP	A	20-30	CW	10YR 4/2	10YR 3/1	sil	--	0	--	3 f, m abk	MH	FI	SO	PO		
3	LP	Bt1	30-60	GW	2.5YR 6/2	10YR 5/3	sicl	--	0	--	2 m, c sbk	H	VFI	SS	MP		
4	LP	2Bt2	60-90	GW	10YR 6/3	70% 10YR 4/3 30% 10YR 5/3	sicl	2% scattered f, m rounded gr.			2 m pr → 2 m sbk	H	VFI	SS	MP		
5	LP	2Bt3	90-130	AW	10YR 4/4	40% 7.5YR 4/3 60% 7.5YR 3/3	sil	2% scattered f, m rounded gr.			1 m, co. pr → 2 m sbk	MH	FI	SS	SP		
6	LP	3Bw	130-145	AW	7.5YR 5/4	7.5YR 4/6	xgrscl	85% f, m, co. rounded gravels mixed lithology			0 sq	L	L	SO	PO		
7	LP	4Btb1	145-163	GW	7.5YR 5/6	7.5YR 4/6	cl	10% f, rounded gr. mixed lithology			2 m sbk → 3 vf, f sbk	VH	EF	MS	MP		
8	LP	4Btb2	163-210	DW	7.5YR 4/4	7.5YR 4/4	cl	1% f, m rounded gr. mixed lithology			3 co., vco pr → 3 f, m sbk	EH	SR	MS	MP	15% coarse, faint, 10YR 4/3 mottles, M, irregular, on faces of peds	
9	LP	4BC	210-230	DI	2.5YR 7/2	2.5YR 5/2	c	1% f, m rounded gr. mixed lithology			3 co., vco pr → 2 m sbk	EH	R	VS	VP	None	
10	SP	4C	230-260+	---	2.5YR 5/2	2.5YR 5/2	c	1% f, m rounded gr. mixed lithology			3 co., vco pr → 3 f, m abk	R	R	VS	VP	None	

	Redoximorphic Features								Concentrations								Ped / V. Surface Features				Roots			Pores			pH, method	Effer (agent)	Clay %	Sand %	Notes
	% Sz	Cn	Cd	Hn	Sp	Kd	Loc	Bd	Col	% Sz	Cn	Hd	Sp	Kd	Loc	Bd	Col	% Dst	Cont	Kd	Loc	Col	Qty	Sz	Loc	Qty					
1	None								None								None				1 m T 2 vf, f T			few, very fine, dendr. tubular			5.0, **NE, H2 M11				(** pH by pocket meter, 1:1 soil to water)
2																	1 m T 2 vf, f T			2 vf TE			6.0	NE, H2							
3																	60%, faint, clay films (CFL) on ped faces (PF)				2 vf T 1 f T			2 vf, f TE			6.7	NE, H2			
4																	60%, faint, clay films (CFL) on ped faces (PF)				1 vf T 1 f T			2 vf, f TE			6.9	NE, H2			
5																	30%, faint, clay films (CFL) on ped faces (PF)				few, very fine, between peds			2 vf TE			7.2	NE, H2			
6																	20%, prominent, clay films on rock fragments (RF)				None			3 vf, f IR			7.1	NE, H2			gravel pavement with scattered, small gully fills
7																	85%, P, clay films (CLF) on all ped faces (PF)							2 vf, f TE			7.1	NE, H2			till joint ghosts remain; truncated paleosol; strong argillans & pedo. structure
8	common, med., distinct 10YR 6/3 iron depletions in matrix																40%, D, CLF on PF							2 vf TE 1 f TE			7.6	NE, H2			till joints ghost and fade upwards to top at 45°; clay & Fe/Mn coated prisms
9	f, 3, P, 10YR 2/1, MNF, APE																27%, D, CLF on ped faces (PF)							2 vf TE			7.7	SL, H2			polygonal till joints ghost & tip 30° to North (down slope)
10	c, 4, P, 2.5 / N, MNF, on prism faces (APF)								c, vco, P, white, l, CaCO <sub>3</sub> nodules along joints (CRK) & in matrix								7%, P, pressure faces (PRF) on PF throughout							1 vf, f IG			8.2, M11	SL, H2 (nodules VE)			till joints tip 30° to North (down slope)

# SUBAQUEOUS SOILS (SAS) DESCRIPTION

*S. McVey, P.J. Schoeneberger, J. Turenne, M. Payne, and D.A. Wysocki, NRCS, and M. Stolt, URI*

**DISCUSSION:** Permanently submerged mineral or organic substrates covered by relatively shallow water display recognizable soil morphology and meet Simonson's soil formation (1959) model in that chemical and physical additions, losses, transformations, and translocations created the morphology. Such soils are informally known as "subaqueous soils." Kubienna (1953) proposed a comprehensive classification that included subaqueous soils. More recently, Demas (1993, 1998) and Demas et al. (1996) reintroduced subaqueous soil concepts in the U.S. Recent reviews (Stolt and Rabenhorst, 2012; Soil Survey Staff, 2012d) provide comprehensive treatment of subaqueous soil settings and processes. Payne (2010) presents operational methods for subaqueous soil inventory. The 11th edition of *Keys to Soil Taxonomy* (Soil Survey Staff, 2010) presently recognizes subaqueous soils as suborders of Entisols and Histosols (Wassents and Wassists) that meet the criterion of "a positive water potential at the soil surface for more than 21 hours of each day in all years."

The description of subaqueous soils is similar to that of terrestrial soils but differs in several important ways. Many subaqueous soil parameters (color, texture, RMF, etc.) fit traditional descriptive conventions outlined in this Field Book. The unique setting and morphology of subaqueous soil coupled with its recent scientific import warrant a separate section that presents all descriptors in one place. This section includes description forms and subaqueous soil description examples. (**NOTE:** The most prevalent subaqueous settings are coastal marine or brackish estuarine. The descriptive conventions presented here reflect this. Freshwater subaqueous settings may require additional descriptors.)

**SUBAQUEOUS SOILS DESCRIPTION**—Record subaqueous soil profile information using the following parameters. (**NOTE:** Field Book soil descriptors presented elsewhere [e.g., horizon] have page number references. Please refer to the cited page for complete choice lists.)

## **BATHYMETRY**

---

Bathymetry is the measurement of sea- or lake-floor or river bottom relief. Because of nautical importance, bathymetric data are commonly expressed as a depth from the water surface at Mean Lower Low Water (MLLW) tidal datum to the bottom. The water surface reference in a coastal setting is commonly Mean Low Water (MLW) or Mean Tide Level (MTL) (see graphic on p. 2–99). Lack of bathymetric data often requires field collection of such data during

subaqueous soil survey. Protocols for bathymetric data collection are addressed elsewhere (Payne, 2010; Bradley and Payne, 2010).

The inverse of water depth is the subaqueous soil survey relief, which is useful for interpretation of subaqueous geomorphs (landforms) and geomorphic description. Geomorphic description for subaqueous soils follows the same convention as those for terrestrial soils (p. 1–4). A compendium of subaqueous geomorphs exists in the Geomorphic Description System (GDS) (Schoeneberger and Wysocki, 2012) (p. 3–38).

## **SITE**

---

**SITE/PEDON ID**—Record the site/pedon identification number, such as the Soil Survey Site Identification Number (see p. 1–2). A complete example is *S2011RI009014A*. (Translation: This is a pedon sampled [S] for soil characterization during 2011 [2011], from Rhode Island [RI], in Washington County [009]; it is the fourteenth pedon [014] sampled in that county during 2011; and it is a satellite sample [A] related to the primary pedon.)

**DATE**—Record the date the sample was collected; e.g., *MM, DD, YYYY*.

**TIME: START/END**—Record the time that the pedon was opened (Start Time) and exposed to aerobic conditions for description and the time that the description was finished (End Time). (**NOTE:** First describe soil color and other soil properties that can change as a result of oxidation.)

**DESCRIBER(S)**—Record the people who describe the core; e.g., *Herman Munster* or *HM*.

**WAYPOINT** (Number)—Record the GPS waypoint number.

**GPS** (Model)—Record the GPS model used. (In NASIS, this is a text field.)


**COORDINATES**—Record the sample site GPS coordinates. (**NOTE:** For subaqueous soils, location is always obtained as a GPS coordinate.)

**GEODETTIC DATUM**—Record the Geodetic Datum (called **GPS Datum** in NASIS) used; e.g., *WGS84*.

**ELEVATION**—Elevation should be normalized relative to an appropriate vertical or tidal datum. In the U.S., use the North American Vertical Datum, 1988 (NAVD88). This elevation datum is the standard shown on recent USGS topographic maps. **NOTE:** For elevations below Mean Tidal Level (MTL), a minus sign precedes the numerical value (e.g., *-1.2 m*). For terrestrial anthropogenic water

bodies (e.g., reservoir), the elevational reference (datum) is the design pool level.

**TIDAL DATUM (DISCUSSION)**—In a tidal system, a Tidal Datum is the elevation of the contact between open water and the land at a specified mean tidal level. A variety of tidal datums are commonly used (see Tidal Datums graphic). Mean Tide Level (MTL), formerly known as Mean Sea Level (MSL), is the average of all tidal fluctuation in a given area and represents the land-water interface on conventional topographic maps. It is also the datum to which terrestrial elevations are normalized. Mean High Water (MHW) and Mean Low Water (MLW) represent typical 24-hour tidal cycles. Mean Higher High Water (MHHW) and Mean Lower Low Water (MLLW) are based on lunar cycles that increase the amplitude of tides. Nautical charts used for boating are based on Mean Lower Low Water in an attempt to represent minimum water depths for navigation. Highest and Lowest Observed water depths are often related to severe storm events (e.g., storm surge) that exaggerate the typical tidal water depth. Specific vegetative communities are associated with the various tidal datums.



<sup>1</sup> Stylized SAS upper limit: ≥21 hours submerged/day (Soil Survey Staff, 2010a, p. 123).

<sup>2</sup> Tidal datum widely used on nautical maps for navigational context.

**MAP UNIT**—Record the map unit name or symbol in which the sample site occurs.

**LOCATION DESCRIPTION**—Record relevant geographic information (e.g., *Greenwich Bay, Warwick, RI—southeastern shoreline, 1300 m SE of Sally Rock Point*).

**WATER DEPTH**—Record water depth at observation time; e.g., *120 cm*.

**BOTTOM TYPE**—Record the dominant bottom type (used in combination with subaqueous vegetation type) at the sample site; e.g., *sand*.

<b>Bottom Type</b> <sup>1</sup>	<b>Criteria</b>
Mud	A silty, clayey, or organic bottom matrix.
Sand	A sandy bottom matrix.
Shelly	A bottom dominated by shells or shell fragments.
Stony or Bouldery	A bottom sparsely covered by stones or boulders (0.01 to <0.1%).
Very Stony or Very Bouldery	A bottom partially covered by stones or boulders (0.1 to <3%).
Extremely Stony or Extremely Bouldery	A bottom dominated by stones or boulders (3 to <15%).
Gravelly/Cobbly	A bottom sparsely covered by gravel or cobbles (0.01 to <0.1%).
Very Gravelly/ Very Cobbly	A bottom partially covered by gravel or cobbles (0.1 to <3%).
Extremely Gravelly/ Extremely Cobbly	A bottom dominated by gravel or cobbles (3 to <15%).
Rubblly	A bottom substantially covered by large rock fragments of various sizes (15 to <50%).
Very Rubblly	A bottom extensively covered by boulders (>50%) <sup>2</sup> .

<sup>1</sup> These bottom types have been used in coastal SAS mapping. Other types exist and should be added as necessary. A formal ecologically based substrate array is found in the Coastal and Marine Ecological Classification Standard (Federal Geographic Data Committee, 2012).

<sup>2</sup> If the surface rock fragments are >80%, the fragments are described as a distinct horizon and should be described as such (NSSH Amendment #4, 1998).

**SUBMERGED AQUATIC VEGETATION (SAV)** (In NASIS, called **Site Existing Vegetation** table); e.g., *ZOSTE* [*Zostera L.*], eelgrass.—Record the *Plant Symbol*, *Plant Common Name*, and *Plant Scientific Name* of aquatic plants observed at the site (see p. 1–17). It is helpful to record the estimated percent cover for each major plant.

**OBSERVATION METHOD**—For each layer, describe the **Kind** of sampling device or procedure used to make observations of the site. Methods and tools for SAS are listed in the following table.

**KIND** (called **Observation Method** in NASIS)—

Kind	Code	Criteria: Tools and Methods
Bucket Auger	BA	Open, closed, sand, or mud buckets (5-12 cm diam.)
Dutch or Mud Auger	DA	An open, strap-sided bucket auger (5-10 cm diam.) with a sharpened outer edge and a screw tip with a partial twist.
Dive	DV	A visual, onsite assessment performed under water.
Macaulay Sampler	MS	A half-cylinder “gouge” sampler with a hinged door that’s pushed in and partially rotated to obtain a sample of soft sediments (e.g., organics).
Vibracore Tube	VT	A hollow tube (e.g., 4-8 cm diam.) vibrated into wet sands, silts, or organics.
Video	VO	Electronically recorded, photo or sequential digital images of a subaqueous setting/site.

**(SOIL) DRAINAGE CLASS**—Subaqueous soils have, by definition, a *subaqueous* (soil) Drainage Class (p. 1–11). There is a positive water potential at the soil surface for more than 21 hours of each day. The soils have a *peraquic* soil moisture regime.

## WATER COLUMN MEASUREMENTS

---

Water quality measurements are not required but are recommended to provide supplemental information on the specific aquatic environment in which the soil is found.

**pH**—Measure the pH (see p. 2–86) at two depths within the water column above the soil.

<b>pH (#)</b>	<b>Criteria</b>
pH top	Within 10 cm of the water surface.
pH bottom	Within 10 cm of the bottom.

**pH METHOD**—Record the pH method used, as there can be considerable difference between methods (e.g., *pH meter* vs. *pH indicator strip*) (see p. 2–86).

**Dissolved Oxygen or DO (mg/l)**—Measure the dissolved oxygen (DO) at two depths within the water column above the soil.

<b>Dissolved Oxygen (DO) (#)</b>	<b>Criteria (mg/l)</b>
DO top	Within 10 cm of the water surface.
DO bottom	Within 10 cm of the bottom.

**SALINITY (ppt)**—Measure the salinity; record in parts per thousand (ppt) at two depths within the water column above the soil. Measurement methods include handheld salinity meters and refractometers (e.g., *YSI salinity meter*, *Vee Gee refractometer*).

<b>Salinity (#)</b>	<b>Criteria</b>
Salinity top	Within 10 cm of the water surface.
Salinity bottom	Within 10 cm of the bottom.

**WATER TEMPERATURE**—Record the water temperature at two depths within the water column above the soil.

<b>Water Temperature (#)</b>	<b>Criteria: Degrees</b> (Celsius [°C] or Fahrenheit [°F])
Water Temperature top	Within 10 cm of the water surface.
Water Temperature bottom	Within 10 cm of the bottom.



## **SOIL PROFILE MEASUREMENTS AND DESCRIPTION**

**HORIZON AND LAYER DESIGNATIONS**—See p. 2–2.

### **HORIZON SUFFIXES FOR SUBAQUEOUS SOILS**

**(DISCUSSION)**—Amongst the conventional list of horizon subscripts (see *Horizon Subscripts*; p. 2–4), several suffixes are used extensively for subaqueous soils and warrant brief clarification:

**g Strong gleying:** The suffix g is used for soil horizons (including subaqueous soils) where Fe has been reduced and pedogenically removed, resulting in a chroma of 2 or less. (**NOTE:** Subaqueous soils are permanently submerged, and most are dominated by reduced conditions and subsequent gray colors, indicated by the use of the g suffix. The g is not applied to soil materials with gray colors attributed to the natural color of the geologic material from which they are derived [geogenic colors; e.g., gray shales].)

**se Presence of sulfides:** This symbol indicates the presence of sulfides in mineral or organic horizons. Horizons with sulfides typically have dark colors (e.g., value  $\leq 4$ , chroma  $\leq 2$ ).

**DISCUSSION:** An se horizon typically forms in coastal soil environments that are permanently saturated or submerged (e.g., low tidal marshes, lagoons, and some freshwater marshes or swamps). Soil materials in which sulfur reduction actively occurs release hydrogen sulfide gas ( $H_2S$ ), which is detectable by its distinctive odor (Fanning and Fanning, 1989; Fanning et al., 2002). (**NOTE:** Not all sulfide-bearing soil materials produce hydrogen sulfide gas. Sulfides may also occur in drier [oxidized] upland environments that have a geologic sulfide source. Examples include soils formed in parent materials derived from coal deposits [e.g., lignite] or soils that formed in coastal plain deposits [e.g., marcasite or pyrite] that have not been oxidized because of thick layers of overburden.)

**LITHOLOGIC DISCONTINUITIES (DISCUSSION)**—Describe the presence of any lithologic discontinuities (see p. 2–5; Numerical Prefixes; The Prime). Contrasting changes in parent material are indicated in Soil Horizon nomenclature by a sequential numeric prefix (e.g., *A-Bw-2Bw-3C*). (**NOTE:** The prefix “1” is implied and not shown by convention.) Discontinuities in subaqueous soils are described when there is a significant change in particle size or mineralogy that indicates the material was deposited by a different process. Important examples are a discontinuity at the change from material deposited in a marine environment to older material deposited on land and later inundated, or sand deposited over marsh organics. In contrast, deposits of similar particle size from multiple washover events on a washover-fan flat behind a barrier

island would not be described as discontinuities (analogous to finely stratified alluvium).

**USE OF PRIMES**—A prime (') is used for horizons with identical characters that are separated by a horizon with different designations. The prime symbol ('), where appropriate, is placed after the master horizon designation and before the lowercase suffix letter symbols that follow it; e.g., *B't*. In cases where three to five horizons have identical letter symbols, three to five prime symbols can be used for the other horizons (e.g., *A, Cg, Aseb, Cseg1, Cseg2, A'seb, Cseb, A''seb, C'seb*).

**HORIZON THICKNESS**—See p. 2–6.

**HORIZON BOUNDARY**—Record **HORIZON BOUNDARY DISTINCTNESS** when possible (see p. 2–6). (**NOTE:** The **HORIZON BOUNDARY TOPOGRAPHY** cannot be adequately determined from small auger, push tube, or vibracore samples.)

**ROCKS AND OTHER FRAGMENTS (Coarse Frags)**—Describe the **Kind**, **Size**, and **Quantity** (% vol.), **Roundness**, etc., of rock and other coarse fragments in each horizon (see p. 2–46). Shell fragments > 2 mm are considered to be coarse fragments.

**FIELD TEXTURE CLASS**—Estimate the **Field Texture Class** by hand for each horizon (see p. 2–36).

**SOIL COLOR**—Record the Munsell color of the soil matrix (see p. 2–9). Include **Color Condition** as needed (e.g., *reduced*).

**REACTION TO H<sub>2</sub>O<sub>2</sub> (or Peroxide Color Change—see SAS Description Form)** (In NASIS, called **Reduced Monosulfide Presence**)—Record the soil color response, as either **yes (Y)** or **no (N)**, to the application of 3% H<sub>2</sub>O<sub>2</sub> solution immediately after exposure to the air (e.g., a freshly broken ped or core interior). A positive reaction (color change) indicates the presence of reduced monosulfides (FeS), which quickly oxidize and change color upon application of hydrogen peroxide. “Peroxide Color Change” is an immediate (within 10 seconds), discernible color change upon addition of H<sub>2</sub>O<sub>2</sub>. (**NOTE:** This method is for monosulfide detection only and is *not* applicable to other sulfides [e.g., pyrite, marcasite, FeS<sub>2</sub>].)

Monosulfides, often in the form of Fe(II) monosulfides (FeS), are visible in reduced soil as a black color (e.g., 10YR 2/1 or N 2.5/0). When a sulfidic soil is oxidized, either in place due to oxidized water conditions or when the soil is drained or excavated and thus exposed to air (oxidized), Fe(II) converts to Fe(III) and the typical black color is lost, leaving a gray or brown color (Lyle, 1983). An example of a common monosulfide oxidation reaction is:

$$4\text{FeS} + 9\text{O}_2 + 10\text{H}_2\text{O} = 4\text{Fe}(\text{OH})_3 + 8\text{H}^+ + 4\text{SO}_4$$

**REACTION BY OXIDIZED pH (DISCUSSION)** (Not a field test)—Oxidation of sulfides creates sulfuric acid as a byproduct that lowers pH. Monosulfidic materials are typically identified in the laboratory using an “oxidized pH” measurement (pH 1:1 water by pH meter), in which a soil undergoes aerobic incubation for at least 16 weeks and the change in pH measurements are compared over time (especially initial vs. final). Sulfidic materials are indicated when 1) the initial pH >3.5 and 2) after oxidation the pH decreases by ≥0.5 unit to a value ≤4.0 within 16 weeks (or longer if the pH is still dropping after 16 weeks) until the pH reaches a nearly constant value. Exposure and oxidation of sulfidic materials (acid sulfate weathering) result in a sulfuric horizon via the formation of sulfuric acid. Field pH is initially measured either immediately after sampling or after thawing a frozen sample. Care should be taken to prevent oxidation of the sample prior to starting the aerobic incubation period and measurements. Hydrogen peroxide has also been used to determine the presence of reduced sulfides in soil samples with pH measurements made after complete oxidation with H<sub>2</sub>O<sub>2</sub> (Finkelman and Giffin, 1986; Jennings et al., 1999). Hydrogen peroxide speeds up the natural oxidation reaction and can be represented in the following reaction as:

$$2\text{FeS} + 9\text{H}_2\text{O}_2 = 2\text{Fe}(\text{OH})_3 + 2\text{SO}_4^{2-} + 4\text{H}^+ + 4\text{H}_2\text{O}.$$

**REACTION BY OXIDIZED pH** (laboratory test)—Measure the pH over time; report the initial and final pH (after 16 weeks) and compare results for evidence of pH reduction over time.

**FLUIDITY CLASS**—See p. 2–65, under **Manner of Failure**. Record the Fluidity class of each horizon/layer. (**NOTE:** Fluidity is estimated by squeezing a moist to wet palmful of soil and observing the extent to which the soil flows out between clenched fingers. Fluidity classes are based on the degree of “flow.” Soil bearing capacity decreases as fluidity increases.)

Fluidity Class	Code	Criteria
Nonfluid	NF	After full compression, no soil flows through the fingers.
Slightly Fluid	SF	After full compression is exerted, some soil flows through fingers; most remains in the palm.
Moderately Fluid	MF	After full pressure is exerted, most soil flows through fingers; some remains in the palm.
Very Fluid	VF	Under very gentle pressure, most soil flows through the fingers as a slightly viscous fluid; very little or no residue remains in the palm of the hand.

**ODOR**—Record the **Kind** and **Intensity** immediately after soil is exposed to air (see p. 2–90).

**ORIGIN**—Record the source of parent material from which the soil is derived (e.g., *estuarine deposit*).

**NOTES**—See **Miscellaneous Field Notes** (see p. 2–93). Describe supplemental information in the “Miscellaneous Field Notes” area or the “Notes” column on the description form. For example, describe the dominant plant fragment type and % (e.g., black needlerush fragments, 25%).

## ***SALINITY (of Subaqueous Soils)***

---

### **ELECTRICAL CONDUCTIVITY OF SUBAQUEOUS SOILS**

**(DISCUSSION)**—Salinity in terrestrial soils is evaluated by electrical conductivity (ds/m) of a saturated paste extract (Saturated Paste method, SSIR 51; Soil Survey Staff, 2009) or a given soil:water (by weight) ratio (e.g., 1:2, 1:5). The saturated paste EC is the standard salinity measure for terrestrial soils and is placed into *Salinity Classes* (see p. 2–89). Salinity is a crucial property of subaqueous soils and is also evaluated via EC measurement. The EC measurement, however, is conducted on a 1:5 by *volume* soil:solution mixture (1:5 vol. method, SSIR 51; Soil Survey Staff, 2009). The resultant EC is **not** placed into the conventional (terrestrially focused) *Salinity Classes*.

Electrical conductivity of subaqueous soils is measured on samples that typically have been stored in a refrigerator or freezer immediately after sampling to prevent sulfide oxidation, which can influence the EC value. Terrestrial soil samples are dried prior to preparing a saturated paste extract. Subaqueous soils cannot be dried because of the sulfide oxidation potential. Hence, a 1:5 volume method is used as follows: Measure 10 ml of moist sample; add 50 ml distilled water (5 times the soil volume). Stir the mixture briefly (10 seconds) and let settle (15 to 60 minutes). Electrical conductivity of the unfiltered supernatant is measured using a hand-held conductivity meter.

**Electrical Conductivity of SAS (1:5 vol method)**—Measure and record the **Electrical Conductivity (EC)** in dS/m and record the **Measurement Method** used (e.g., *11.2 dS/m by hand-held electrical conductivity meter*).

**SULFIDES (DISCUSSION)**—Identifying the presence of sulfate is important to both pH and salinity (see **Reaction by Oxidized pH**). Oxidation of sulfides may generate salts that can alter salinity.

**BULK DENSITY SATIATED (DISCUSSION)** (These are variations of the Soil Core Method. See Section 3.3.1.4 in SSIR 51, Soil Survey Staff, 2009)—It is generally not possible to collect subaqueous soil

samples using the clod method for bulk density determination. Recommended alternative methods are:

- 1) Collect a known volume at the field moisture state (satiated). Bulk density is then calculated based on the dried weight of a known volume of soil at the field moisture status.

**Calculations** (Soil Survey Staff, 2009)

$Db = (ODW - RF - CW) / [CV - (RF/PD)]$ , where:

Db = Bulk density of <2-mm fabric at sampled, field water state ( $\text{g cm}^{-3}$ )

ODW = Oven-dry weight

RF = Weight of rock fragments

CW = Empty core weight

CV = Core volume

PD = Density of rock fragments

- 2) For vibracore samples (opened by cutting the sampling tube rather than by compressive extrusion), a 50-ml syringe with the end removed and shaped to fit the curved core is used as a mini-corer to extract a 10- to 30-ml volume sample. The cylinder is removed, extracting a sample of known volume. The sample is then analyzed following method 1 (above).
- 3) Samples collected in a peat sampler (e.g., Macaulay sampler) can be analyzed for bulk density following method 2 if a known volume (e.g., a core segment) is collected and dried.



## SUBAQUEOUS SOILS PROFILE DESCRIPTION

<b>Site/Pedon ID:</b>		<b>Map Unit:</b>			
<b>Date:</b>		<b>Location Description:</b>	<b>Water Column Measurements</b>		
<b>Start Time:</b>					
<b>End Time:</b>		<b>Water Depth (cm):</b>		<b>pH:</b>	
<b>Describer(s):</b>		<b>Bottom Type:</b>		<b>DO (mg/l):</b>	
<b>Waypoint (#):</b>		<b>Submerged Aq. Veg:</b>		<b>salinity (ppt):</b>	
<b>GPS (unit #):</b>		<b>Observation Method:</b>		<b>temp (°C):</b>	
<b>Coordinates 1:</b>		<b>Site Notes:</b>			
<b>Coordinates 2:</b>					
<b>Geodetic Datum:</b>					

Depth (cm)	Horizon	Horizon Boundary Distinctness	Soil Color (matrix)	Field Texture Class	Coarse Frags (%)	Fluidity Class	RMFs	Peroxide Color Change (Y/N)	Oxidized pH		Odor (Intensity, Kind)	Origin	Notes
									init.	16 wks.			

## SUBAQUEOUS SOILS PROFILE DESCRIPTION

<b>Site/Pedon ID:</b>	S2011R1009014A	<b>Map Unit:</b>	Frankensoil mucky silt loam				
<b>Date:</b>	8/16/2011	<b>Location Description:</b>	Ninigret Pond; 1000 m. E. of intersection of Route 1 and Route 1A at Ninigret Park, RI		<b>Water Column Measurements</b>		
<b>Start Time:</b>	8:30 AM					<b>Top</b>	<b>Bottom</b>
<b>End Time:</b>	11:45 AM	<b>Water Depth (cm):</b>	120 cm	<b>pH:</b>	7.7	7.7	
<b>Describer(s):</b>	Herman Munster	<b>Bottom Type:</b>	mud	<b>DO (mg/l):</b>	6	5	
<b>Waypoint (#):</b>	4	<b>Submerged Aq. Veg:</b>	thick macroalgae	<b>salinity (ppt):</b>	27	29	
<b>GPS (unit #):</b>	Trimble Geo XH	<b>Observation Method:</b>	Vibracore tube	<b>temp (°C):</b>	20 °C	18 °C	
<b>Coordinates 1:</b>	N 41° 22' 13.0" Lat	<b>Site Notes:</b>					
<b>Coordinates 2:</b>	W 71° 39' 4.0" Lon						
<b>Geodetic Datum:</b>	WGS 84						

Depth (cm)	Horizon	Horizon Boundary Distinctness	Soil Color (matrix)	Field Texture Class	Coarse Frags (%)	Fluidity Class	RMFs	Peroxide Color Change (Y/N)	Oxidized pH		Odor (Intensity, Kind)	Origin	Notes
									init.	16 wks.			
0-12	A	Abrupt	5Y 6/1	mucky silt loam	0	Very Fluid		Y	7.8	4.7	strong sulfurous	marine silt	pH by pH meter
12-53	C1	Clear	5Y 2.5/1	mucky silt loam	0	Moderately Fluid		Y	7.7	4.9	strong sulfurous	marine silt	
53-88	C2	Abrupt	5Y 3/1	mucky silt loam	0	Moderately Fluid		Y	8.0	2.6	strong sulfurous	marine silt	
88-98	20a1	Abrupt	N 2.5/	muck	14 % gr	Slightly Fluid		N	7.8	6.6	slight sulfurous	organics, fresh	
98-130	20a2	Abrupt	10YR 2/1	muck	1 % wood frags	Slightly Fluid		N	7.7	6.5	none	organics, fresh	
130-191	20a3	—	10YR 2/2	muck	1 % wood frags	Slightly Fluid		N	7.7	6.5	none	organics, fresh	



# **VIBRACORE SAMPLING FOR SUBAQUEOUS SOILS**

---

## **Discussion**

Subaqueous soils are challenging to observe and sample because of the positive pore pressure of free water in the soil and the water above the soil; therefore, a slightly different protocol for sampling them is helpful. Vibracore sampling is particularly well suited to obtain minimally disturbed samples from sandy, silty, or organic subaqueous materials lacking large or substantial coarse fragments. The principal concern is accounting for sample compression (compaction, repacking, or “core rot”), especially in material with *Moderately Fluid* or *Very Fluid* fluidity classes. (**NOTE:** A Vibracore Log Sheet [p. 2–113] must be paired with a Subaqueous Soil Profile Description [p. 2–109] or a conventional Soil Profile Description [p. 2–93].)

## **Site Description**

Record subaqueous soil site information much the same as you record subaerial soil site information (see p. 1–1). Additional items to evaluate and describe include the following:

**WATER DEPTH (UNIT)**—Record the depth of water (and the units used) above the soil surface at the time the core is collected. (This information is used to develop/verify the map unit name; e.g., *Billington silt loam, 0-1 m water depth.*)

**TIDAL PERIOD**—Record the tidal period (*incoming, high, outgoing, low, none*) at the time of sample extraction. (**NOTE:** Most freshwater lakes do not exhibit appreciable tidal fluctuations.)

## **Core Descriptions**

**TOTAL PIPE LENGTH**—Describe the total length of the collection pipe prior to coring. (This information is used as a check for depth of sample collection. The information may also be used to track how much pipe is consumed during a field season.)

**RISER LENGTH**—After insertion of the pipe tube, record the external length (cm) from the top of the collection pipe to the soil surface outside the pipe.

**INSIDE LENGTH**—After insertion of the pipe tube, record the length (cm) from the top of the collection pipe to the soil surface *inside* the pipe. (**NOTE:** A sinker tied to a string and lowered to the soil surface inside the pipe facilitates measurement.)

**CORE SETTLEMENT**—Calculate the sample settlement/compaction (also called “rot”) by subtracting the Riser Length from the Inside Length. (**NOTE:** Settlement of the soil sample inside the pipe is

common and difficult to precisely account for; therefore, vibracore samples only provide reasonable estimates of horizon depths.)

**FINAL CORE LENGTH**—Record the calculated length of the soil profile collected. (See graphic on Vibracore Log Sheet, p. 2–113.)

**WHERE CORE IS STORED**—Describe where the core is stored for future retrieval, description, and analysis (e.g., *shed 2, core # 2011-25*). (**NOTE:** Subaqueous soils should be kept in refrigerated storage to slow chemical reactions, such as conversion of sulfides to sulfates, which influence pH or other soil properties.)

# VIBRACORE LOG SHEET


## SITE

<b>Site/Pedon ID</b> (YYYYSTFIPS####)	
<b>Date/Time Sampled</b>	
<b>Soil Type</b>	
<b>Map Unit</b>	
<b>Location</b> (geographic)	
<b>Waypoint</b> (#)	
<b>GPS</b> (model/unit #)	
<b>Lat.</b>	
<b>Lon.</b>	
<b>UTM Easting</b>	
<b>UTM Northing</b>	
<b>UTM Zone</b>	
<b>Elevation</b> (NAVD 88)	
<b>Water Depth</b> (cm)	
<b>Tidal Period</b>	

## CORE LOG

<b>a) TOTAL Pipe Length</b> (before coring)	
<b>b) RISER Length</b> (after coring)	
<b>c) INSIDE Length</b> (sinker length: surface to bottom)	
<b>d) Core Settlement</b> (= c - b)	
<b>Final Core Length</b> (after core completed: = a - c)	
<b>Where Is Core Stored?</b>	
<b>Date Described</b>	

**Core Sketch**



# VIBRACORE LOG SHEET EXAMPLE


## SITE

<b>Site/Pedon ID</b> (YYYYSTFIPS###)	S2011R1009014A
<b>Date/Time Sampled</b>	8/16/2011 8:30 AM
<b>Soil Type</b>	Frankensoil
<b>Map Unit</b>	Frankensoil mucky silt loam
<b>Location</b> (geographic)	Ninigret Pond: 1000 m E. of intersection of Route 1 and Route 1A at Ninigret Park, RI
<b>Waypoint</b> (#)	4
<b>GPS</b> (model/unit #)	Trimble Geo XH
<b>Lat.</b>	41° 22' 13.0"
<b>Lon.</b>	W 71° 39' 4.0"
<b>UTM Easting</b>	721720 m
<b>UTM Northing</b>	4583254 m
<b>UTM Zone</b>	19
<b>Elevation</b> (NAVD 88)	- 1.2 m
<b>Water Depth</b> (cm)	120 cm
<b>Tidal Period</b>	Outgoing

## CORE LOG

<b>a) TOTAL Pipe Length</b> (before coring)	390 cm
<b>b) RISER Length</b> (after coring)	260 cm
<b>c) INSIDE Length</b> (sinker length: surface to bottom)	264 cm
<b>d) Core Settlement</b> (= c - b)	4 cm
<b>Final Core Length</b> (after core completed: = a - c)	126 cm
<b>Where Is Core Stored.</b>	URI Bay Campus cold storage
<b>Date Described</b>	8/17/2011

**Core Sketch**



## REFERENCES

---

- AASHTO. 1997a. The classification of soils and soil-aggregate mixtures for highway construction purposes. AASHTO designation M145–91. *In* Standard specifications for transportation materials and methods of sampling and testing, Part 1: Specifications (18th ed.). American Association of State Highway and Transportation Officials, Washington, DC.
- AASHTO. 1997b. Terms relating to subgrade, soil aggregate, and fill materials. AASHTO designation M146–91 (1995). *In* Standard specifications for transportation materials and methods of sampling and testing, Part 1: Specifications (18th ed.). American Association of State Highway and Transportation Officials, Washington, DC.
- Amoozegar, A., and A.W. Warrick. 1986. Hydraulic conductivity of saturated soils: Field methods. *In* A. Klute (ed.) Methods of soil analysis, Part 1: Physical and mineralogical methods, 2nd ed. ASA, Agron. Monogr. 9, Madison, WI.
- ASTM. 2011. Standard practice for classification of soils for engineering purposes (Unified Soil Classification System). ASTM designation D2487–92. *In* Soil and rock; dimension stone; geosynthetics. Annual book of ASTM standards.
- Bachman, G.O., and M.W. Machette. 1977. Calcic soils and calcretes in the southwestern U.S. U.S. Geol. Soc. Open-File Rep. 77–794.
- Blackburn, W.H., R.E. Eckert, Jr., M.K. Wood, and F.F. Peterson. 1975. Influence of vesicular horizons on watershed management. p. 494–515. *In* Proc. ASCE Watershed Management Symposium. Logan, UT, 11–13 Aug. 1975. ASAE, New York.
- Bouma, J., R.F. Paetzold, and R.B. Grossman. 1982. Measuring hydraulic conductivity for use in soil survey. Soil Surv. Invest. Rep. 38. USDA, SCS. U.S. Gov. Print. Office, Washington, DC.
- Bradley, M.P., and M. Payne. 2010. Bathymetric mapping and landscape units in shallow subtidal coastal environments: Integrating fathometry, GPS, and GIS. *In* 2nd National Workshop on Subaqueous Soils. Rhode Island, 9–12 Aug. 2010. Available online at [http://nesoil.com/sas/15A\\_Bradley\\_Bathymetry\\_GIS.pdf](http://nesoil.com/sas/15A_Bradley_Bathymetry_GIS.pdf).
- Brewer, R. 1976. Fabric and mineral analysis of soils. Krieger Publ. Co., Huntington, NY.
- Brock, A.L. 2007. Characterization of stage VI petrocalcic horizons—S. Nevada and N.W. Arizona. Ph.D. diss. Univ. of Nevada Las Vegas, Department of Geoscience.
- Brock, A.L. 2010. Personal communication (carbonate stages source graphic).

Bullock, P., N. Fedoroff, A. Jongerius, G. Stoops, and T. Tursina. 1985. Handbook for soil thin section description. Waine Res. Publ., Wolverhampton, England.

Bureau of Chemistry and Soils. 1937. Soil survey manual. USDA Misc. Publ. 274. Washington, DC.

Childs, C.W. 1981. Field tests for ferrous iron and ferric-organic complexes (on exchange sites or in water-soluble forms) in soils. *Aust. J. Soil Res.* 19:175–180.

Cruden, D.M., and D.J. Varnes. 1996. Landslide types and processes. *In* A.K. Turner and R.L. Schuster (ed.) *Landslides: Investigation and mitigation*. Spec. Rep. 247, Transportation Research Board, National Research Council. National Academy Press, Washington, DC.

Demas, G.P. 1993. Submerged soils: A new frontier in soil survey. *Soil Surv. Horiz.* 34:44–46.

Demas, G.P. 1998. Subaqueous soils of Sinepuxent Bay, MD. Ph.D. diss. Univ. of Maryland, College Park.

Demas, G.P., R.B. Rabenhorst, and J.C. Stevenson. 1996. Subaqueous soils: A pedological approach to the study of shallow-water habitats. *Estuaries* 19:229–237.

Doner, H.E., and W.C. Lynn. 1989. Carbonate, halide, sulfate, and sulfide minerals. *In* J.B. Dixon and S.B. Weed (ed.) *Minerals in the soil environment*, 2nd ed. SSSA Book Ser. 1. SSSA, Madison, WI.

Fanning, D.S., and M.C.B. Fanning. 1989. *Soil: Morphology, genesis, and classification*. John Wiley & Sons, New York.

Fanning, D.S., M.C. Rabenhorst, S.N. Burch, K.R. Islam, and S.A. Tangren. 2002. Sulfides and sulfates. *In* J.B. Dixon et al. (ed.) *Soil mineralogy with environmental applications*. SSSA, Madison, WI.

Federal Geographic Data Committee. 2012. Coastal and marine ecological classification standard, ver. 4.0. Federal Geographic Data Committee Secretariat, U.S. Geol. Surv., Reston, VA.

Finkelman, R.B., and D.E. Giffin. 1986. Hydrogen peroxide oxidation: An improved method for rapidly assessing acid-generation potential of sediments and sedimentary rocks. *Recl. Reveg. Res.* 5:521–534.

Flach, K.W., W.D. Nettleton, L.H. Gile, and J.G. Cady. 1969. Pedocementation: Induration by silica, carbonates, and sesquioxides in the Quaternary. *Soil Sci.* 107:442–453.

Folk, R.L. 1955. Student operator error in determination of roundness, sphericity and grain size. *J. Sediment. Petrol.* 25:297–301.

- Gile, L.H. 1961. A classification of Ca horizons in soils of a desert region, Doña Ana County, New Mexico. *Soil Sci. Soc. Am. Proc.* 25(1):52–61.
- Gile, L.H. 1970. Soils of the Rio Grande valley border in southern New Mexico. *Soil Sci. Soc. Am. Proc.* 34(3):465–472.
- Gile, L.H. 1975. Causes of soil boundaries in an arid region, I. Age and parent materials. *Soil Sci. Soc. Am. Proc.* 39:316–323.
- Gile, L.H. 1993. Carbonate stages in sandy soils of the Leasburg surface, southern New Mexico. *Soil Sci.* 156:101–110.
- Gile, L.H., and R.B. Grossman. 1968. Morphology of the argillic horizon in desert soils of southern New Mexico. *Soil Sci.* 106(1):6–15.
- Gile, L.H., and R.B. Grossman. 1979. The desert project soil monograph. USDA, Natl. Tech. Inf. Serv. Doc. PB80–13534. Springfield, VA.
- Gile, L.H., J.W. Hawley, and R.B. Grossman. 1981. Soils and geomorphology in the Basin and Range area of southern New Mexico—Guidebook to the desert project. New Mexico Bureau of Mines and Mineral Resources, Memoir 39.
- Gile, L.H., H.C. Monger, R.B. Grossman, R.J. Ahrens, J.W. Hawley, F.F. Peterson, R.P. Gibbens, J.M. Lenz, B.T. Bestelmeyer, and B.A. Nolen. 2007. A 50th anniversary guidebook for the desert project. USDA, NRCS, National Soil Survey Center, Lincoln, NE.
- Gile, L.H., F.F. Peterson, and R.B. Grossman. 1966. Morphological and genetic sequences of carbonate accumulation in desert soils. *Soil Sci.* 101:347–360.
- Guthrie, R.L., and J.E. Witty. 1982. New designations for soil horizons and layers and the new soil survey manual. *Soil Sci. Soc. Am. J.* 46:443–444.
- Hillel, D. 1980. *Fundamentals of soil physics*. Academic Press, New York, NY.
- Ingram, R.L. 1982 (updated by M. Johnson, 2007). Modified Wentworth scale. *In* Grain-size scales. AGI data sheet 8.2. J.D. Walker and H.A. Cohen (ed.) 2005 (reprinted 2007). *The geoscience handbook*. AGI data sheets, 4th ed. Am. Geol. Inst., Alexandria, VA.
- Jackson, J.A. (ed.) 1997. *Glossary of geology*, 4th ed. Am. Geol. Inst., Alexandria, VA.
- Jackson, M.L. 1969. *Soil chemical analysis—Advanced course*. Madison, WI.
- Jennings, S.R., D.J. Dollhopf, and W.P. Inskeep. 1999. Acid production from sulfide minerals using hydrogen peroxide weathering. *Appl. Geochem.* 15:235–243.

- Kubienna, W.M. 1953. *The soils of Europe*. T. Murby, London.
- Lyle, M. 1983. The brown-green color transition in marine sediments: A marker of the Fe(III)-Fe(II) redox boundary. *Limnol. Oceanogr.* 28:1026–1033.
- Lynn, W., and D. Williams. 1992. The making of a Vertisol. *Soil Surv. Horiz.* 33:23–52.
- Machette, M.N. 1985. Calcic soils of the southwestern United States. *In* D.L. Weide (ed.) *Soils and Quaternary geomorphology of the southwestern United States*. Geol. Soc. Am. Spec. Pap. 203.
- Mason, D.D., J.F. Lutz, and R.G. Petersen. 1957. Hydraulic conductivity as related to certain soil properties in a number of great soil groups—Sampling errors involved. *Soil Sci. Soc. Am. Proc.* 21:554–561.
- National Institute of Standards and Technology. 1990. Counties and equivalent entities of the United States, its possessions and associated areas. U.S. Dep. Commerce, Federal Information Processing Standards Publ. (FIPS PUB 6–4).
- Natural Resources Conservation Service. 1997. Introduction to microbotic crusts. USDA, NRCS, Soil Quality Institute and Grazing Lands Technology Institute, Fort Worth, TX.
- Natural Resources Conservation Service. 2001. Physical and biological soil crusts. Soil Quality Information Sheet—Rangeland Information Sheet 7. USDA, NRCS, Soil Quality Institute and National Soil Survey Center, Lincoln, NE.
- Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. USDA, Agric. Handb. 296. U.S. Gov. Print. Office, Washington, DC.
- Natural Resources Conservation Service. 2012 (or most current date) [Online]. Complete PLANTS checklist. (The national PLANTS database.) USDA, National Plant Data Center, Baton Rouge, LA (<http://plants.usda.gov>).
- Neuendorf, K., J.P. Mehl, and J.A. Jackson. 2005. *Glossary of geology*, 5th ed. Am. Geol. Inst., Alexandria, VA.
- O’Neal, A.M. 1952. A key for evaluating soil permeability by means of certain field clues. *Soil Sci. Soc. Am. Proc.* 16:312–315.
- Payne, M. 2010. Bathymetry data collection for subaqueous soil mapping. *In* 2nd National Workshop on Subaqueous Soils. Rhode Island, 9–12 Aug. 2010 ([http://nesoil.com/sas/15\\_Payne\\_Bathymetry.pdf](http://nesoil.com/sas/15_Payne_Bathymetry.pdf)).
- Powers, M.C. 1953. A new roundness scale for sedimentary particles. *J. Sediment. Petrol.* 23:117–119.



Public Building Service. Sept. 1996. Worldwide geographic location codes. U.S. Gen. Serv. Admin., Washington, DC.

Richards, L.A. (chairman) 1952. Report of the subcommittee on permeability and infiltration, committee on terminology, Soil Science Society of America. *Soil Sci. Soc. Am. Proc.* 16:85–88.

Ruhe, R.V. 1975. *Geomorphology: Geomorphic processes and surficial geology*. Houghton-Mifflin Co., Boston, MA.

Schoeneberger, P.J., and D.A. Wysocki. 2012. Geomorphic description system, ver. 4.2. USDA, NRCS, National Soil Survey Center, Lincoln, NE ([ftp://ftp-fc.sc.egov.usda.gov/NSSC/GDS/GDS\\_v4\\_14.pdf](ftp://ftp-fc.sc.egov.usda.gov/NSSC/GDS/GDS_v4_14.pdf)).

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and W.D. Broderson. 1998. Field book for describing and sampling soils (ver. 1.1). USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Simonson, R.W. 1959. Outline of a generalized theory of soil genesis. *Soil Sci. Soc. Am. Proc.* 23:152–156.

Singer, M.J., and D.N. Warrington. 1992. Crusting in the western United States. *In* M.E. Sumner and B.A. Stewart (ed.) *Advances in soil science: Soil crusting—Chemical and physical processes*. Lewis Publishers, Boca Raton, FL.

Soil Conservation Service. 1981. Land resource regions and major land resource areas of the United States. USDA, Agric. Handb. 296. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Division Staff. 1993. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1951. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1962. Identification and nomenclature of soil horizons. Supplement to Agric. Handb. 18, Soil Survey Manual (replacing pages 173–188). USDA, SCS. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1971. Guide for interpreting engineering uses of soils. USDA Handb. NB–1. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1975. Soil taxonomy. USDA, SCS, Agric. Handb. 436. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1999. Soil taxonomy, 2nd ed. USDA, NRCS, Agric. Handb. 436. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 2009. Soil survey field and laboratory methods manual. Soil Surv. Invest. Rep. 51, ver. 1.0. R. Burt (ed.) USDA, NRCS.

Soil Survey Staff. 2010. Keys to soil taxonomy, 11th ed. USDA, NRCS. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 2011. Soil survey laboratory information manual. Soil Surv. Invest. Rep. 45, ver. 2.0. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 2012a. Data dictionary. *In* National soil information system (NASIS), release 6.2 [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 2012b. National soil information system (NASIS), release 6.2 [Online]. USDA, NRCS, Lincoln, NE.

Soil Survey Staff. 2012c. National soil survey handbook (NSSH) [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE (<http://soils.usda.gov/technical/handbook/>).

Soil Survey Staff. 2012d. Subaqueous soils primer (draft). USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Stolt, M.H., and M.C. Rabenhorst. 2012. Introduction and historical development of subaqueous soil concepts. *In* P.M. Huang et al. (ed.) Handbook of soil science. CRC Press, LLC, Boca Raton, FL. ISBN: 978-1-4398-0305-9.

Sumner, M.E., and B.A. Stewart (ed.) 1992. Soil crusting: Chemical and physical processes. Advances in soil science. Lewis Publishers, Boca Raton, FL.

Uhland, R.E., and A.M. O'Neal. 1951. Soil permeability determinations for use in soil and water conservation. SCS-TP-101. Washington, DC.

Vepraskas, M.J. 1992. Redoximorphic features for identifying aquic conditions. North Carolina Agric. Res. Serv. Tech. Bull. 301. North Carolina State University, Raleigh.

Wysocki, D.A., P.J. Schoeneberger, and H.E. LaGarry. 2000. Geomorphology of soil landscapes. *In* M.E. Sumner (ed.) Handbook of soil science. CRC Press, LLC, Boca Raton, FL. ISBN: 0-8493-3136-6.

Wysocki, D.A., P.J. Schoeneberger, B. Lowery, and R. Paetzold. 2002. Hydraulic conductivity vs. soil permeability classes: Facts, fiction, and the future. ASA, Indianapolis, IN.

# GEOMORPHIC DESCRIPTION

## **GEOMORPHIC DESCRIPTION SYSTEM**

*(Version 5.0—08/14/2017)*

*P.J. Schoeneberger and D.A. Wysocki, NRCS, Lincoln, NE*

### **PART I: PHYSIOGRAPHIC LOCATION**

- A) Physiographic Division
- B) Physiographic Province
- C) Physiographic Section
- D) State Physiographic Area
- E) Local Physiographic/Geographic Name

### **PART II: GEOMORPHIC DESCRIPTION**

- A) Landscape
- B) Landform
- C) Microfeature
- D) Anthropogenic Features

### **PART III: SURFACE MORPHOMETRY**

- A) Elevation
- B) Slope Aspect
- C) Slope Gradient
- D) Slope Complexity
- E) Slope Shape
- F) Hillslope—Profile Position
- G) Geomorphic Component
  - 1. Hills
  - 2. Terraces, Stepped Landforms
  - 3. Mountains
  - 4. Flat Plains
- H) Microrelief
- I) Drainage Pattern

**NOTE:** *Italicized NASIS shorthand codes, if available, follow each choice. Conventionally, the entire term is recorded.*

## **PART I: PHYSIOGRAPHIC LOCATION**

---

References for items **A**, **B**, and **C**: *Physical Divisions of the United States* (Fenneman, 1946); *Physiographic Divisions of Alaska* (Wahrhaftig, 1965).

<b>Physiographic Divisions (A)</b>		<b>Physiographic Provinces (B)</b>	<b>Physiographic Sections (C)</b>
<b>Laurentian Upland</b>	<i>LU</i>	1. Superior Upland	<i>SU</i>
<b>Atlantic Plain</b>	<i>AP</i>	2. Continental Shelf	<i>CS</i>
		3. Coastal Plain	<i>CP</i>
		a. Embayed section	<i>EMS</i>
		b. Sea Island section	<i>SIS</i>
		c. Floridian section	<i>FLS</i>
		d. East Gulf Coastal plain	<i>EGC</i>
		e. Mississippi alluvial valley	<i>MAV</i>
		f. West Gulf Coastal plain	<i>WGC</i>
<b>Appalachian Highlands</b>	<i>AH</i>	4. Piedmont Province	<i>PP</i>
		a. Piedmont upland	<i>PIU</i>
		b. Piedmont lowlands	<i>PIL</i>
		5. Blue Ridge Province	<i>BR</i>
		a. Northern section	<i>NOS</i>
		b. Southern section	<i>SOS</i>
		6. Valley and Ridge Province	<i>VR</i>
		a. Tennessee section	<i>TNS</i>
		b. Middle section	<i>MIS</i>
		c. Hudson Valley	<i>HUV</i>
		7. St. Lawrence Valley	<i>SL</i>
		a. Champlain section	<i>CHS</i>
		b. St. Lawrence Valley, Northern section	<i>NRS</i>
		8. Appalachian Plateau	<i>AP</i>
		a. Mohawk section	<i>MOS</i>
		b. Catskill section	<i>CAS</i>
		c. Southern New York sect.	<i>SNY</i>
		d. Allegheny Mountain sect.	<i>AMS</i>
		e. Kanawaha section	<i>KAS</i>
		f. Cumberland Plateau sect.	<i>CPS</i>
		g. Cumberland Mtn. sect.	<i>CMS</i>



**Physiographic Divisions (A)      Physiographic Provinces (B)  
 Physiographic Sections (C)**

<b>Rocky Mountain System</b>	<i>RM</i>	16. Southern Rocky Mountains	<i>SR</i>
		17. Wyoming Basin	<i>WB</i>
		18. Middle Rocky Mountains	<i>MR</i>
		19. Northern Rocky Mountains	<i>NR</i>

This division includes portions of Alaska  
 (see "Alaskan Physiographic Areas" section).

<b>Intermontane Plateaus</b>	<i>IP</i>	20. Columbia Plateau	<i>CR</i>	
		a. Walla Walla Plateau	<i>WWP</i>	
		b. Blue Mountain section	<i>BMS</i>	
		c. Payette section	<i>PAS</i>	
		d. Snake River Plain	<i>SRP</i>	
	e. Harney section	<i>HAS</i>		
	21. Colorado Plateau			<i>CO</i>
		a. High Plateaus of Utah		<i>HPU</i>
		b. Uinta Basin		<i>UIB</i>
		c. Canyon Lands		<i>CAL</i>
		d. Navajo section		<i>NAS</i>
		e. Grand Canyon section		<i>GCS</i>
	f. Datil section		<i>DAS</i>	
	22. Basin and Range Province			<i>BP</i>
		a. Great Basin		<i>GRB</i>
		b. Sonoran Desert		<i>SOD</i>
		c. Salton Trough		<i>SAT</i>
		d. Mexican Highland		<i>MEH</i>
	e. Sacramento section		<i>SAS</i>	

This division includes portions of Alaska  
 (see "Alaskan Physiographic Areas" section).

<b>Pacific Mountain</b>	<i>PM</i>	23. Cascade-Sierra Mountains	<i>CM</i>	
		a. Northern Cascade Mtns.	<i>NCM</i>	
		b. Middle Cascade Mtns.	<i>MCM</i>	
		c. Southern Cascade Mtns.	<i>SCM</i>	
	d. Sierra Nevada	<i>SIN</i>		
	24. Pacific Border Province			<i>B</i>
		a. Puget Trough		<i>PUT</i>
		b. Olympic Mountains		<i>OLM</i>
		c. Oregon Coast Range		<i>OCR</i>



<b>Physiographic Divisions (A)</b>	<b>Physiographic Provinces (B) Physiographic Sections (C)</b>
------------------------------------	---

<b>Intermontane Plateaus</b>	<i>IP</i>	Northern Plateaus Province	—
		8. Porcupine Plateau section	—
		a. Thazzik Mountain	
		9. Old Crow Plain section (noted but not described)	—
		10. Olgivie Mountains section	—
		11. Tintina Valley (Eagle Trough) sect.	—
		12. Yukon-Tanana Upland section	—
		a. Western Part	
		b. Eastern Part	
		13. Northway-Tanacross Lowland sect.	—
		14. Yukon Flats section	—
		15. Rampart Trough section	—
		16. Kokrine-Hodzana Highlands sect.	—
		a. Ray Mountains	
		b. Kokrine Mountains	
		Western Alaska Province	—
		17. Kanuti Flats section	—
		18. Tozitna-Melozitna Lowland sect.	—
		19. Indian River Upland section	—
		20. Pah River Section	—
		a. Lockwood Hills	
		b. Pah River Flats	
		c. Zane Hills	
		d. Purcell Mountains	
		21. Koyukuk Flats section	—
		22. Kobuk-Selawik Lowland sect.	—
		a. Waring Mountains	
		23. Selawik Hills section	—
		24. Buckland River Lowland sect.	—
		25. Nulato Hills section	—
		26. Tanana-Kuskowin Lowland sect.	—
		27. Nowitna Lowland section	—
		28. Kuskokwim Mountains section	—
		29. Innoko Lowlands section	—
		30. Nushagak-Big River Hills sect.	—
		31. Holitna Lowland section	—
		32. Nushagak-Bristol Bay Lowland sect.	—



<b>Physiographic Divisions (A)</b>	<b>Physiographic Provinces (B)</b>	<b>Physiographic Sections (C)</b>
------------------------------------	------------------------------------	-----------------------------------

- |     |                                       |     |
|-----|---------------------------------------|-----|
|     | 33. Seward Peninsula Province         | SEP |
|     | a. Bendeleben Mountains               |     |
|     | b. Kigluaik Mountains                 |     |
|     | c. York Mountains                     |     |
|     | Bering Shelf Province                 | BES |
| 34. | Yukon-Kuskokwim Coastal Lowland sect. | —   |
|     | a. Norton Bay Lowland                 |     |
| 35. | Bering Platform section               | —   |
|     | a. St. Lawrence Island                |     |
|     | b. Pribilof Island                    |     |
|     | c. St. Matthew Island                 |     |
|     | d. Nunivak Island                     |     |
| 36. | Ahklun Mountains Province             | —   |

**NOTE:** The map unit numbering sequence shown here is from Wahrhaftig (1965) and is independent of, and not consistent with, that of Fenneman.

- |                                |           |  |            |
|--------------------------------|-----------|--|------------|
| <b>Pacific Mountain System</b> | <i>PM</i> | Alaska-Aleutian Province                           | <i>AAC</i> |
|                                |           | 37. Aleutian Islands section                       | —          |
|                                |           | 38. Aleutian Range section                         | —          |
|                                |           | 39. Alaska Range (Southern Part) sect.             | —          |
|                                |           | 40. Alaska Range (Central & Eastern Parts) section | —          |
|                                |           | a. Mentasta-Nutzotin Mtn. segment                  |            |
|                                |           | 41. Northern Foothills of the Alaska Range section | —          |
|                                |           | Coastal Trough Province                            | —          |
|                                |           | 42. Cook Inlet-Susitna Lowland sect.               | —          |
|                                |           | 43. Broad Pass Depression section                  | —          |
|                                |           | 44. Talkeetna Mountains section                    | —          |
|                                |           | a. Chulitna Mountains                              |            |
|                                |           | b. Fog Lakes Upland                                |            |
|                                |           | c. Central Talkeetna Mountains                     |            |
|                                |           | d. Clarence Lake Upland                            |            |
|                                |           | e. Southeastern Talkeetna Mountains                |            |
|                                |           | 45. Upper Matanuska Valley sect.                   | —          |
|                                |           | 46. Clearwater Mountains section                   | —          |



**Physiographic Divisions (A) Physiographic Provinces (B)  
 Physiographic Sections (C)**

<b>Caribbean Basin</b>	<i>CB</i>	Caribbean Islands Province	<i>CI</i>
		a. Greater Antilles (Puerto Rico, Cuba, Hispaniola, Jamaica)	<i>GRA</i>
		b. Lesser Antilles (U.S. Virgin Islands, Barbados, Grenada, Martinique, etc.)	<i>LEA</i>
		c. Other	
<b>Undesignated</b>	<i>UN</i>	Other	<i>OT</i>
		(reserved for temporary or international designations)	

**STATE PHYSIOGRAPHIC AREA (D)**

e.g., *Des Moines Lobe* (IA)

(OPTIONAL) (Entries presently undefined; to be developed in conjunction with each State Geological Survey; target scale is approximately 1:100,000.)

**LOCAL PHYSIOGRAPHIC/GEOGRAPHIC NAME (E)**

e.g., *Pilot's Knob* (IA)

(OPTIONAL) (Entries presently undefined; to be developed in conjunction with each State Geological Survey; may include area names found on USGS 7.5- and 15-minute topographic maps; target scale is approximately 1:24,000.)

## **PART II: GEOMORPHIC DESCRIPTION (OUTLINE)**

**I) COMPREHENSIVE LISTS:** Alphabetical rosters of all terms currently recognized in a given category.

**A) LANDSCAPES**

**B) LANDFORMS**

**C) MICROFEATURES**

**D) ANTHROPOGENIC FEATURES**

**II) GEOMORPHIC ENVIRONMENTS and OTHER GROUPINGS:**

Landscape, landform, and microfeature terms grouped by geomorphic process (e.g., Fluvial) or by common settings (e.g., Water Bodies). These lists are *not* mutually exclusive; some features occur in more than one environment or setting (e.g., *hill*).

- |                                 |   |                                |
|---------------------------------|---|--------------------------------|
| 1. Coastal Marine and Estuarine | } | <b>GEOMORPHIC ENVIRONMENTS</b> |
| 2. Lacustrine                   |   |                                |
| 3. Fluvial                      |   |                                |
| 4. Solution                     |   |                                |
| 5. Eolian                       |   |                                |
| 6. Glacial                      |   |                                |
| 7. Periglacial                  |   |                                |
| 8. Mass Movement                |   |                                |
| 9. Volcanic and Hydrothermal    |   |                                |
| 10. Tectonic and Structural     |   |                                |
| <hr/>                           |   |                                |
| 11. Slope                       | } | <b>OTHER GROUPINGS</b>         |
| 12. Erosional                   |   |                                |
| 13. Depressional                |   |                                |
| 14. Wetlands                    |   |                                |
| 15. Water Bodies                |   |                                |
| 16. Subaqueous Features         |   |                                |

## **PART II: GEOMORPHIC DESCRIPTION**

---

Codes: Conventionally, the entire land-feature term is used (e.g., dune field). Some data storage programs (e.g., NASIS) may have shorthand codes developed for some terms. If available, an italicized code follows each term (e.g., meander belt MB); these are shown for historical purposes.

### **I) COMPREHENSIVE LISTS:**

#### **A) LANDSCAPES** (broad assemblages or unique groups of natural, spatially associated features). (LF=Landform; w=water body)

alluvial plain	<i>AP</i>	fold-thrust hills	<i>FTH</i>
alluvial plain remnant	<i>AR</i>	foothills	<i>FH</i>
badlands	<i>BA</i>	glaciokarst	<i>GK</i>
bajada (also LF)	<i>BJ</i>	gulf (w; also LF)	<i>GU</i>
barrier island (also LF)	<i>BI</i>	hills (singular=LF)	<i>HI</i>
basin	<i>BS</i>	ice-margin complex	<i>IC</i>
basin floor (also LF)	<i>BC</i>	intermontane basin (also LF)	<i>IB</i>
batholith	<i>BL</i>	island (also LF)	<i>IS</i>
bay [coast] (w; also LF)	<i>BY</i>	karst	<i>KR</i>
bolson	<i>BO</i>	kegel karst	<i>KK</i>
breached anticline (also LF)	<i>BD</i>	lagoon (w; also LF)	<i>LG</i>
breaklands	<i>BR</i>	lake plain (also LF)	<i>LP</i>
breaks (also LF)	<i>BK</i>	lava field (also LF)	<i>LF</i>
caldera (also LF)	<i>CD</i>	lava plain (also LF)	<i>LV</i>
canyonlands	<i>CL</i>	lava plateau (also LF)	<i>LL</i>
coastal plain (also LF)	<i>CP</i>	lowland	<i>LW</i>
cockpit karst	<i>CPK</i>	marine terrace (also LF)	<i>MT</i>
cone karst	<i>CK</i>	meander belt	<i>MB</i>
continental glacier	<i>CG</i>	mountain range	<i>MR</i>
delta plain (also LF)	<i>DP</i>	mountains (singular=LF)	<i>MO</i>
dissected breaklands	<i>DB</i>	mountain system	<i>MS</i>
dissected plateau	<i>DI</i>	ocean (w)	<i>OC</i>
drumlin field	<i>DF</i>	outwash plain (also LF)	<i>OP</i>
dune field (also LF)	<i>DU</i>	peninsula	<i>PE</i>
estuary (w; also LF)	<i>ES</i>	piedmont	<i>PI</i>
everglades	<i>EG</i>	piedmont slope	<i>PS</i>
fan piedmont (also LF)	<i>FP</i>	plain (singular=LF)	<i>PL</i>
fault-block mountains	<i>FM</i>	plateau (also LF)	<i>PT</i>
fluviokarst	<i>FK</i>	rift valley	<i>RF</i>
fluviomarine terrace (also LF)	<i>FT</i>	river valley (also LF)	<i>RV</i>

sand plain	<i>SP</i>	strait (w; also LF)	<i>ST</i>
sandhills	<i>SH</i>	tableland	<i>TB</i>
scabland	<i>SC</i>	thermokarst	<i>TK</i>
sea (w; also LF)	<i>SEA</i>	till plain (also LF)	<i>TP</i>
semi-bolson	<i>SB</i>	tower karst	<i>TW</i>
shield volcano (also LF)	<i>SV</i>	upland	<i>UP</i>
shore complex (also LF)	<i>SX</i>	valley (also LF)	<i>VA</i>
sinkhole karst	<i>SK</i>	volcanic field (also LF)	<i>VF</i>
sound (w; also LF)	<i>SO</i>		

**B) LANDFORMS** (discrete, natural, individual earth-surface features mappable at common survey scales).  
(LS=Landscape; Micro=Microfeature; w=water body.  
Italicized NASIS code follows each term.)

aa lava flow	<i>ALF</i>	basin floor (also LS)	<i>BC</i>
alas	<i>AA</i>	basin-floor remnant	<i>BD</i>
alluvial cone	<i>AC</i>	bay [coast] (w; also LS)	<i>BAY</i>
alluvial fan	<i>AF</i>	bay [geom.]	<i>BYG</i>
alluvial flat	<i>AP</i>	bay bottom	<i>BOT</i>
alpine glacier	<i>AG</i>	bayou (w)	<i>WC</i>
anticline	<i>AN</i>	beach	<i>BE</i>
arete	<i>AR</i>	beach plain	<i>BP</i>
arroyo	<i>AY</i>	beach ridge	<i>BG</i>
ash field	<i>AQ</i>	beach terrace	<i>BT</i>
ash flow	<i>AS</i>	berm	<i>BM</i>
atoll	<i>AT</i>	beveled base	<i>BVB</i>
avalanche chute	<i>AL</i>	blind valley	<i>VB</i>
axial stream	<i>AX</i>	block field	<i>BW</i>
back-barrier beach	<i>BBB</i>	block glide	<i>BLG</i>
back-barrier flat	<i>BBF</i>	block lava flow	<i>BLF</i>
backshore	<i>AZ</i>	block stream	<i>BX</i>
backswamp	<i>BS</i>	blowout	<i>BY</i>
bajada (also LS)	<i>BJ</i>	bluff	<i>BN</i>
ballena	<i>BL</i>	bog	<i>BO</i>
ballon	<i>BV</i>	box canyon	<i>BOX</i>
bar	<i>BR</i>	braided stream	<i>BZ</i>
barchan dune	<i>BQ</i>	breached anticline (also LS)	<i>BRL</i>
barrier beach	<i>BB</i>	breaks (also LS)	<i>BK</i>
barrier beach [relict]	<i>BBR</i>	broad interstream divide	<i>BID</i>
barrier cove	<i>BAC</i>	butte	<i>BU</i>
barrier flat	<i>BF</i>	caldera (also LS)	<i>CD</i>
barrier island (also LS)	<i>BI</i>	canyon	<i>CA</i>

canyon bench	<i>CYB</i>	deflation flat	<i>DFL</i>
canyon wall	<i>CW</i>	delta	<i>DE</i>
Carolina Bay	<i>CB</i>	delta plain (also LS)	<i>DC</i>
channel (also Micro)	<i>CC</i>	depression	<i>DP</i>
chenier	<i>CG</i>	diapir	<i>DD</i>
chenier plain	<i>CH</i>	diatrema	<i>DT</i>
cinder cone	<i>CI</i>	dike	<i>DK</i>
cirque	<i>CQ</i>	dip slope	<i>DL</i>
cirque floor	<i>CFL</i>	disintegration moraine	<i>DM</i>
cirque headwall	<i>CHW</i>	distributary	<i>DIS</i>
cirque platform	<i>CPF</i>	divide	<i>DN</i>
cliff	<i>CJ</i>	dome	<i>DO</i>
climbing dune	<i>CDU</i>	drainageway	<i>DQ</i>
closed depression (also Micro)	<i>CLD</i>	drainhead complex	<i>DRC</i>
coastal plain (also LS)	<i>CP</i>	draw	<i>DW</i>
cockpit	<i>COC</i>	drumlin	<i>DR</i>
col	<i>CL</i>	drumlinoid ridge	<i>DRR</i>
collapse sinkhole	<i>CSH</i>	dune	<i>DU</i>
collapsed ice-floored lakebed	<i>CK</i>	dune field (also LS)	<i>DUF</i>
collapsed ice-walled lakebed	<i>CN</i>	dune lake (w)	<i>DUL</i>
collapsed lake plain	<i>CS</i>	dune slack (also Micro)	<i>DUS</i>
collapsed outwash plain	<i>CT</i>	earthflow	<i>EF</i>
colluvial apron	<i>COA</i>	earth spread	<i>ESP</i>
complex landslide	<i>CLS</i>	earth topple	<i>ETO</i>
coral island	<i>COR</i>	end moraine	<i>EM</i>
coulee	<i>CE</i>	ephemeral stream (also Micro)	<i>EPS</i>
cove	<i>CO</i>	eroded fan remnant	<i>EFR</i>
cove [water] (w)	<i>COW</i>	eroded fan-remnant sideslope	<i>EFS</i>
crag and tail	<i>CAT</i>	erosion remnant	<i>ER</i>
creep	<i>CRE</i>	escarpment	<i>ES</i>
crevasse filling	<i>CF</i>	esker	<i>EK</i>
cuesta	<i>CU</i>	estuary (w; also LS)	<i>WD</i>
cuesta valley	<i>CUV</i>	faceted spur	<i>FS</i>
cutoff	<i>CV</i>	fall	<i>FB</i>
debris avalanche	<i>DA</i>	falling dune	<i>FDU</i>
debris fall	<i>DEF</i>	fan	<i>FC</i>
debris flow	<i>DF</i>	fan apron	<i>FA</i>
debris slide	<i>DS</i>	fan collar	<i>FCO</i>
debris spread	<i>DES</i>	fanhead trench	<i>FF</i>
debris topple	<i>DET</i>	fan piedmont (also LS)	<i>FG</i>
deflation basin	<i>DB</i>	fan remnant	<i>FH</i>

fan skirt	<i>FI</i>	gulf (w; also LS)	<i>GU</i>
fault block	<i>FAB</i>	gut [channel]; (w; also Micro)	<i>WH</i>
fault-line scarp	<i>FK</i>		
fault zone	<i>FAZ</i>	gut [valley]	<i>GV</i>
fen	<i>FN</i>	half graben	<i>HG</i>
fissure vent	<i>FIV</i>	hanging valley	<i>HV</i>
fjord (w)	<i>FJ</i>	headland	<i>HE</i>
flat	<i>FL</i>	head-of-outwash	<i>HD</i>
flatwoods	<i>FLW</i>	headwall	<i>HW</i>
flood plain	<i>FP</i>	high hill	<i>HH</i>
flood-plain playa	<i>FY</i>	highmoor bog	<i>HB</i>
flood-plain splay	<i>FM</i>	hill (plural=LS)	<i>HI</i>
flood-plain step	<i>FO</i>	hillslope	<i>HS</i>
flood-tidal delta	<i>FTD</i>	hogback	<i>HO</i>
flood-tidal delta flat	<i>FTF</i>	homoclinal ridge	<i>HCR</i>
flood-tidal delta slope	<i>FTS</i>	homocline	<i>HC</i>
flow	<i>FLO</i>	horn	<i>HR</i>
flute (also Micro)	<i>FU</i>	horst	<i>HT</i>
fluviomarine bottom	<i>FMB</i>	hot spring	<i>HP</i>
fluviomarine terrace (also LS)	<i>FMT</i>	ice-contact slope	<i>ICS</i>
fold	<i>FQ</i>	ice-marginal stream	<i>IMS</i>
foredune	<i>FD</i>	ice pressure ridge	<i>IPR</i>
fosse	<i>FV</i>	ice-pushed ridge	<i>IPU</i>
free face ( <i>also Geom. Component—Hills, Mountains</i> )	<i>FW</i>	inlet	<i>IL</i>
fringe-tidal marsh	<i>FTM</i>	inselberg	<i>IN</i>
gap	<i>GA</i>	inset fan	<i>IF</i>
geyser	<i>GE</i>	interdrumlin	<i>IDR</i>
geyser basin	<i>GEB</i>	interdune (also Micro)	<i>ID</i>
geyser cone	<i>GEC</i>	interfluve ( <i>also Geom. Component—Hills</i> )	<i>IV</i>
giant ripple	<i>GC</i>	interior valley	<i>INV</i>
glacial drainage channel	<i>GD</i>	intermittent stream (also Micro)	<i>INT</i>
glacial lake (w)	<i>WE</i>	intermontane basin (also LS)	<i>IB</i>
glacial lake [relict]	<i>GL</i>	island (also LS)	<i>IS</i>
glacial-valley floor	<i>GVF</i>	kame	<i>KA</i>
glacial-valley wall	<i>GVW</i>	kame moraine	<i>KM</i>
glacier	<i>GLA</i>	kame terrace	<i>KT</i>
gorge	<i>GO</i>	karst cone	<i>KC</i>
graben	<i>GR</i>	karstic marine terrace	<i>KMT</i>
ground moraine	<i>GM</i>	karst lake	<i>KAL</i>
gulch	<i>GT</i>	karst tower	<i>KTO</i>



karst valley	<i>KVA</i>	marine terrace (also LS)	<i>MT</i>
kettle	<i>KE</i>	marsh	<i>MA</i>
kipuka	<i>KIP</i>	mawae	<i>MAW</i>
knob	<i>KN</i>	meander	<i>MB</i>
knoll	<i>KL</i>	meandering channel	<i>MC</i>
lagoon (w; also LS)	<i>WI</i>	meander scar	<i>MS</i>
lagoon bottom	<i>LBO</i>	meander scroll	<i>MG</i>
lagoon channel	<i>LCH</i>	medial moraine	<i>MH</i>
lagoon [relict]	<i>LAR</i>	mesa	<i>ME</i>
lahar	<i>LA</i>	meteorite crater	<i>MEC</i>
lake (w)	<i>WJ</i>	mogote	<i>MOG</i>
lakebed (w)	<i>LB</i>	monadnock	<i>MD</i>
lakebed [relict]	<i>LBR</i>	monocline	<i>MJ</i>
lake plain (also LS)	<i>LP</i>	moraine	<i>MU</i>
lakeshore	<i>LF</i>	mountain (plural=LS)	<i>MM</i>
lake terrace	<i>LT</i>	mountain slope	<i>MN</i>
landslide	<i>LK</i>	mountain valley	<i>MV</i>
lateral moraine	<i>LM</i>	mudflow	<i>MW</i>
lateral spread	<i>LS</i>	mud pot	<i>MP</i>
lava dome	<i>LD</i>	muskeg	<i>MX</i>
lava field (also LS)	<i>LFI</i>	natural levee	<i>NL</i>
lava flow	<i>LC</i>	nearshore zone	<i>NZ</i>
lava flow unit (also Micro)	<i>LFU</i>	nearshore zone [relict]	<i>NZR</i>
lava plain (also LS)	<i>LN</i>	notch	<i>NO</i>
lava plateau (also LS)	<i>LL</i>	nunatak	<i>NU</i>
lava trench (also Micro)	<i>LTR</i>	open depression (also Micro)	<i>ODE</i>
lava tube	<i>LTU</i>	outwash delta	<i>OD</i>
ledge (also Micro)	<i>LE</i>	outwash fan	<i>OF</i>
levee	<i>LV</i>	outwash plain (also LS)	<i>OP</i>
loess bluff	<i>LO</i>	outwash terrace	<i>OT</i>
loess hill	<i>LQ</i>	overflow stream channel	<i>OSC</i>
longitudinal dune	<i>LDU</i>	oxbow	<i>OX</i>
longshore bar	<i>LON</i>	oxbow lake (w)	<i>WK</i>
longshore bar [relict]	<i>LR</i>	paha	<i>PA</i>
louderback	<i>LU</i>	pahoehoe lava flow	<i>PAF</i>
low hill	<i>LH</i>	paleoterrace	<i>PTR</i>
lowmoor bog	<i>LX</i>	parabolic dune	<i>PB</i>
maar	<i>MAA</i>	parna dune	<i>PD</i>
mainland cove	<i>MAC</i>	partial ballena	<i>PF</i>
main scarp (also Micro)	<i>MAS</i>	patterned ground	<i>PG</i>
mangrove swamp	<i>MAN</i>	pavement karst	<i>PAV</i>
marine lake (w)	<i>ML</i>		

peak	<i>PK</i>	river valley (also LS)	<i>RVV</i>
peat plateau	<i>PJ</i>	roche moutonnée (also Micro)	<i>RN</i>
pediment	<i>PE</i>	rockfall (also Micro)	<i>ROF</i>
perennial stream (w; also Micro)	<i>PS</i>	rockfall avalanche	<i>RFA</i>
pillow lava flow	<i>PIF</i>	rock glacier	<i>RO</i>
pingo	<i>PI</i>	rock pediment	<i>ROP</i>
pinnacle (also Micro)	<i>PIN</i>	rock spread	<i>ROS</i>
pitted outwash plain	<i>PM</i>	rock topple	<i>ROT</i>
pitted outwash terrace	<i>POT</i>	rotational debris slide	<i>RDS</i>
plain (plural=LS)	<i>PN</i>	rotational earth slide	<i>RES</i>
plateau (also LS)	<i>PT</i>	rotational rock slide	<i>RRS</i>
playa	<i>PL</i>	rotational slide	<i>RTS</i>
playa dune (also Micro)	<i>PDU</i>	sabkha	<i>SAB</i>
playa floor (also Micro)	<i>PFL</i>	saddle	<i>SA</i>
playa lake (w)	<i>WL</i>	sag (also Micro)	<i>SAG</i>
playa rim (also Micro)	<i>PRI</i>	sag pond (w; also Micro)	<i>SGP</i>
playa slope (also Micro)	<i>PSL</i>	salt marsh	<i>SM</i>
playa step (also Micro)	<i>PST</i>	salt pond (w; also Micro)	<i>WQ</i>
plug dome	<i>PP</i>	sand flow (also Micro)	<i>RW</i>
pluvial lake (w)	<i>PLL</i>	sand ramp	<i>SAR</i>
pluvial lake [relict]	<i>PQ</i>	sand sheet	<i>RX</i>
pocosin	<i>PO</i>	scarp	<i>RY</i>
point bar	<i>PR</i>	scarp slope	<i>RS</i>
point bar [coastal]	<i>PRC</i>	scree slope	<i>SCS</i>
pothole (also Micro)	<i>PH</i>	sea (w; also LS)	<i>SEA</i>
pothole lake (w)	<i>WN</i>	sea cliff	<i>RZ</i>
proglacial lake (w)	<i>WO</i>	seep (also Micro)	<i>SEE</i>
proglacial lake [relict]	<i>PGL</i>	seif dune	<i>SD</i>
pyroclastic flow	<i>PCF</i>	semi-open depression	<i>SOD</i>
pyroclastic surge	<i>PCS</i>	shield volcano (also LS)	<i>SHV</i>
raised beach	<i>RA</i>	shoal (w)	<i>WR</i>
raised bog	<i>RB</i>	shoal [relict]	<i>SE</i>
ravine	<i>RV</i>	shore	<i>SHO</i>
recessional moraine	<i>RM</i>	shore complex (also LS)	<i>SHC</i>
reef	<i>RF</i>	sill	<i>RT</i>
ribbed fen	<i>RG</i>	sinkhole	<i>SH</i>
ridge	<i>RI</i>	slackwater (w)	<i>WS</i>
rim	<i>RJ</i>	slickrock (also Micro)	<i>SLK</i>
rise (also Micro) ( <i>also Geom. Component—Flat Plains</i> )	<i>RIS</i>	slide	<i>SJ</i>
		slot canyon	<i>SLC</i>
river (w)	<i>RIV</i>	slough (w)	<i>SL</i>

slump block	<i>SN</i>	terrace remnant	<i>TER</i>
snowfield	<i>SNF</i>	thermokarst depression (also Micro)	<i>TK</i>
soil fall	<i>SOF</i>	thermokarst lake (w)	<i>WV</i>
solution platform	<i>SOP</i>	tidal flat	<i>TF</i>
solution sinkhole	<i>SOS</i>	tidal inlet	<i>TI</i>
sound (w; also LS)	<i>SO</i>	tidal inlet [relict] (w)	<i>TIR</i>
spit	<i>SP</i>	tidal marsh	<i>TM</i>
spur	<i>SQ</i>	till-floored lake plain	<i>TLP</i>
stack [coast]	<i>SRC</i>	till plain (also LS)	<i>TP</i>
stack [geom.]	<i>SR</i>	toe (also Micro)	<i>TOE</i>
star dune	<i>SDU</i>	tombolo	<i>TO</i>
steptoe	<i>ST</i>	topple	<i>TOP</i>
stock	<i>STK</i>	tor	<i>TQ</i>
stoss and lee	<i>SAL</i>	Toreva block	<i>TOR</i>
strait (w; also LS)	<i>STT</i>	translational debris slide	<i>TDS</i>
strand plain	<i>SS</i>	translational earth slide	<i>TES</i>
strath terrace	<i>SU</i>	translational rock slide	<i>TRS</i>
stratovolcano	<i>SV</i>	translational slide	<i>TS</i>
stream (w)	<i>STR</i>	transverse dune	<i>TD</i>
stream terrace	<i>SX</i>	trough	<i>TR</i>
strike valley	<i>STV</i>	tunnel valley	<i>TV</i>
string bog	<i>SY</i>	tunnel-valley lake (w)	<i>TVL</i>
structural bench	<i>SB</i>	underfit stream	<i>US</i>
submerged back-barrier beach	<i>SBB</i>	U-shaped valley	<i>UV</i>
submerged mainland beach	<i>SMB</i>	valley (also LS)	<i>VA</i>
submerged point bar [coast]	<i>SPB</i>	valley-border surfaces	<i>VBS</i>
submerged–upland tidal marsh	<i>STM</i>	valley flat	<i>VF</i>
submerged wave-built terrace	<i>SWT</i>	valley floor	<i>VL</i>
submerged wave-cut platform	<i>SWP</i>	valley-floor remnant	<i>VFR</i>
swale (also Micro)	<i>SC</i>	valley side	<i>VS</i>
swallow hole	<i>TB</i>	valley train	<i>VT</i>
swamp	<i>SW</i>	volcanic cone	<i>VC</i>
syncline	<i>SZ</i>	volcanic crater	<i>CR</i>
talus cone	<i>TC</i>	volcanic dome	<i>VD</i>
talus slope	<i>TAS</i>	volcanic field (also LS)	<i>VOF</i>
tarn (w; also Micro)	<i>TAR</i>	volcanic neck	<i>VON</i>
terminal moraine	<i>TA</i>	volcanic pressure ridge (also Micro)	<i>PU</i>
terrace	<i>TE</i>	volcano	<i>VO</i>
		V-shaped valley	<i>VV</i>
		wash	<i>WA</i>
		washover fan	<i>WF</i>

washover-fan flat	WFF	wind gap	WG
washover-fan slope	WFS	window	WIN
water-lain moraine	WM	wind-tidal flat	WTF
wave-built terrace	WT	yardang (also Micro)	YD
wave-cut platform	WP	yardang trough (also Micro)	YDT
wave-worked till plain	WW		

**C) MICROFEATURES** (discrete, natural earth-surface features typically too small to delineate at common survey scales).

1) **Common Microfeatures** (not used in association with the landform “patterned ground”).

bar	BA	open depression (also LF)	OP
channel (also LF)	CH	perennial stream (w; also LF)	PS
closed depression (also LF)	CD	pinnacle (also LF)	PI
corda	CO	playa dune (also LF)	PD
cutter	CU	playa floor (also LF)	PF
dune slack (also LF)	DS	playa rim (also LF)	PR
dune traces	DT	playa slope (also LF)	PSL
earth pillar	EP	playa step (also LF)	PST
ephemeral stream (also LF)	ES	playette	PL
finger ridge	FR	pond (w)	PON
flute (also LF)	FL	pool (w)	POO
frost boil	FB	pothole (also LF)	PH
glacial groove	GG	rib	RB
groove	GR	rill	RL
gully	GU	ripple mark	RM
gut [channel] (w; also LF)	WH	rise (also LF) ( <i>also Geom. Component—Flat Plains</i> )	RIS
hillock	HI	rockfall (also LF)	ROF
hoodoo	HO	roche moutonnée (also LF)	POC
ice wedge	IWD	sag (also LF)	SAG
ice wedge cast	IWC	sag pond (w; also LF)	SP
interdune (also LF)	ID	salt pond (w; also LF)	WQ
intermittent stream (w; also LF)	INT	sand boil	SB
karren	KA	sand flow (also LF)	RW
lava flow unit (also LF)	LFU	seep (also LF)	SE
lava trench (also LF)	LT	shoreline	SH
main scarp (also LF)	MAS	shrub-coppice dune	SCD
minor scarp	MIS	slickrock (also LF)	SLK
mound	MO	slip face	SF
nivation hollow	NH		

solifluction lobe	<i>SOL</i>	tarn (w; also LF)	<i>TN</i>
solifluction sheet	<i>SS</i>	terraces	<i>TER</i>
solifluction terrace	<i>ST</i>	thermokarst depression	<i>TK</i>
solution chimney	<i>SCH</i>	(also LF)	
solution corridor	<i>SCO</i>	toe [mass mvnt.] (also LF)	<i>TOE</i>
solution fissure	<i>SOF</i>	tree-tip mound	<i>TTM</i>
solution pipe	<i>SOP</i>	tree-tip pit	<i>TTP</i>
spatter cone	<i>SPC</i>	tumulus (pl.: tumuli)	<i>TU</i>
spiracle	<i>SPI</i>	vernal pool (seasonal water)	<i>VP</i>
strandline	<i>SL</i>	volcanic pressure ridge	<i>VPR</i>
swale (also LF)	<i>SW</i>	(also LF)	
swash zone	<i>SZ</i>	yardang (also LF)	<i>YD</i>
tank (w)	<i>TA</i>	yardang trough (also LF)	<i>YDT</i>
		zibar	<i>ZB</i>

## 2) **Periglacial "patterned ground" Microfeatures**

(Singular forms [e.g., *circle*] are used for a single feature [pedon scale], whereas plural forms [e.g., *circles*] are used for map unit components.)

circle	<i>CI</i>	palsa (=peat hummock)	<i>PA</i>
earth hummock	<i>EH</i>	polygon	<i>PYG</i>
high-center polygon	<i>HCP</i>	sorted circle	<i>SCI</i>
ice wedge polygon	<i>IWP</i>	stripe	<i>STR</i>
low-center polygon	<i>LCP</i>	turf hummock	<i>TH</i>
nonsorted circle	<i>NSC</i>		

## 3) **Other "patterned ground" Microfeatures**

(Singular forms [e.g., *hummock*] are used for a single feature [pedon scale], whereas plural forms [e.g., *hummocks*] are used for map unit components.)

bar and channel	<i>BC</i>	linear gilgai	<i>LG</i>
circular gilgai	<i>CG</i>	mima mound	<i>MM</i>
elliptical gilgai	<i>EG</i>	pimple mound	<i>PM</i>
gilgai	<i>GI</i>	puff	<i>PU</i>
hummock	<i>HU</i>		

**D) ANTHROPOGENIC FEATURES** (discrete, artificial [human-made] earth-surface features).

anthroscape	<i>ANT</i>	hillslope terrace ( <i>ancient</i> )	<i>HT</i>
artificial collapsed depression	<i>ACD</i>	impact crater	<i>IC</i>
artificial levee	<i>AL</i>	interfurrow	<i>IF</i>
beveled cut	<i>BC</i>	landfill ( <i>see sanitary landfill</i> )	—
bioswale	<i>BS</i>	leveled land	<i>LVL</i>
borrow pit	<i>BP</i>	log landing	<i>LL</i>
burial mound	<i>BM</i>	midden	<i>MI</i>
conservation terrace (modern)	<i>CT</i>	openpit mine	<i>OM</i>
cut ( <i>e.g., railroad</i> )	<i>CUT</i>	polder	<i>POL</i>
cutbank	<i>CB</i>	pond ( <i>human-made</i> )	<i>PO</i>
ditch	<i>DI</i>	quarry	<i>QU</i>
double-bedding mound ( <i>i.e., bedding mound used for timber; lower Coastal Plain</i> )	<i>DBM</i>	railroad bed	<i>RRB</i>
drainage ditch	<i>DD</i>	reclaimed land	<i>RL</i>
dredge-deposit shoal	<i>DDS</i>	rice paddy	<i>RP</i>
dredge spoil bank	<i>DSB</i>	road bed	<i>RB</i>
dredged channel	<i>DC</i>	road cut	<i>RC</i>
dump	<i>DU</i>	sand pit	<i>SP</i>
fill	<i>FI</i>	sanitary landfill	<i>SL</i>
filled marshland	<i>FM</i>	scalped area	<i>SA</i>
floodway	<i>FW</i>	sewage lagoon	<i>SWL</i>
furrow	<i>FR</i>	skid trail	<i>ST</i>
gravel pit	<i>GP</i>	spoil bank	<i>SB</i>
headwall ( <i>anthro</i> )	<i>HW</i>	spoil pile	<i>SPP</i>
		surface mine	<i>SM</i>
		tillage mound	<i>TM</i>
		truncated soil	<i>TS</i>

## II) GEOMORPHIC ENVIRONMENTS and OTHER GROUPINGS:

Landscape, landform, and microfeature terms grouped by geomorphic process (e.g., Fluvial) or by common setting (e.g., Water Body). LS=Landscape; LF=Landform; Micro=Microfeature. Lists are not mutually exclusive.

### 1. COASTAL MARINE and ESTUARINE (wave or tidal control or near-shore/shallow marine).

#### **Landscapes:**

barrier island (also LF)	<i>BI</i>	lagoon (w; also LF)	<i>LG</i>
bay [coast] (w; also LF)	<i>BY</i>	lowland	<i>LW</i>
coastal plain (also LF)	<i>CP</i>	marine terrace (also LF)	<i>MT</i>
delta plain (also LF)	<i>DP</i>	ocean (w)	<i>OC</i>
estuary (w; also LF)	<i>ES</i>	peninsula	<i>PE</i>
fluviomarine terrace (also LF)	<i>FT</i>	sea (w; also LF)	<i>SEA</i>
gulf (w; also LF)	<i>GU</i>	shore complex	<i>SX</i>
island (also LF)	<i>IS</i>	sound (w; also LF)	<i>SO</i>
		strait (w; also LF)	<i>ST</i>

#### **Landforms:**

atoll	<i>AT</i>	delta	<i>DE</i>
back-barrier beach	<i>BBB</i>	delta plain (also LS)	<i>DC</i>
back-barrier flat	<i>BBF</i>	drainhead complex	<i>DRC</i>
backshore	<i>AZ</i>	estuary (also LS)	<i>WD</i>
bar	<i>BR</i>	flat	<i>FL</i>
barrier beach	<i>BB</i>	flatwoods	<i>FLW</i>
barrier cove	<i>BAC</i>	fluviomarine terrace (also LS)	<i>FMT</i>
barrier flat	<i>BF</i>	foredune	<i>FD</i>
barrier island (also LS)	<i>BI</i>	fringe-tidal marsh	<i>FTM</i>
bay [coast] (w; also LS)	<i>BAY</i>	gulf (w; also LS)	<i>GU</i>
bay bottom	<i>BOT</i>	gut [channel] (w, also Micro)	<i>WH</i>
beach	<i>BE</i>	headland	<i>HE</i>
beach plain	<i>BP</i>	island (also LS)	<i>IS</i>
beach ridge	<i>BG</i>	lagoon (w; also LS)	<i>WI</i>
beach terrace	<i>BT</i>	lagoon [relict]	<i>LAR</i>
berm	<i>BM</i>	longshore bar	<i>LON</i>
bluff	<i>BN</i>	longshore bar [relict]	<i>LR</i>
chenier	<i>CG</i>	mangrove swamp	<i>MAN</i>
chenier plain	<i>CH</i>	marine lake (w)	<i>ML</i>
coastal plain (also LS)	<i>CP</i>	marine terrace (also LS)	<i>MT</i>
coral island	<i>COR</i>	nearshore	<i>NZ</i>
cove [water] (w)	<i>COW</i>		

nearshore zone [relict]	<i>NZR</i>	stack [coast]	<i>SRC</i>
point bar [coastal]	<i>PRC</i>	strait (w; also LS)	<i>STT</i>
raised beach	<i>RA</i>	strand plain	<i>SS</i>
reef	<i>RF</i>	submerged-upland tidal marsh	<i>STM</i>
sabkha	<i>SAB</i>	tidal flat	<i>TF</i>
salt marsh	<i>SM</i>	tidal inlet	<i>TI</i>
sea (w; also LS)	<i>SEA</i>	tidal inlet [relict]	<i>TIR</i>
sea cliff	<i>RZ</i>	tidal marsh	<i>TM</i>
semi-open depression	<i>SOD</i>	tombolo	<i>TO</i>
shoal [relict]	<i>SE</i>	washover fan	<i>WF</i>
shore	<i>SHO</i>	wave-built terrace	<i>WT</i>
shore complex (also LS)	<i>SHC</i>	wave-cut platform	<i>WP</i>
sound (w; also LS)	<i>SO</i>	wind-tidal flat	<i>WTF</i>
spit	<i>SP</i>		

**Microfeatures:**

gut [channel] (w; also LF)	<i>WH</i>	shoreline	<i>SH</i>
ripple mark	<i>RM</i>	swash zone	<i>SZ</i>



## 2. LACUSTRINE (related to inland water bodies).

### **Landscapes:**

bay [coast] (w; also LF)	<i>BY</i>	lake plain (also LF)	<i>LP</i>
delta plain (also LF)	<i>DP</i>	peninsula	<i>PE</i>
island (also LF)	<i>IS</i>	shore complex (also LF)	<i>SX</i>

### **Landforms:**

backshore	<i>AZ</i>	longshore bar [relict]	<i>LR</i>
bar (also Micro)	<i>BR</i>	nearshore zone	<i>NZ</i>
barrier beach	<i>BB</i>	nearshore zone [relict]	<i>NZR</i>
barrier flat	<i>BF</i>	oxbow lake	<i>WK</i>
barrier island	<i>BI</i>	playa	<i>PL</i>
bay [coast] (w; also LS)	<i>BAY</i>	playa floor (also Micro)	<i>PFL</i>
beach	<i>BE</i>	playa lake (w)	<i>WL</i>
beach plain	<i>BP</i>	playa rim (also Micro)	<i>PRI</i>
beach ridge	<i>BG</i>	playa slope (also Micro)	<i>PSL</i>
beach terrace	<i>BT</i>	playa step (also Micro)	<i>PST</i>
berm	<i>BM</i>	pluvial lake (w)	<i>PLL</i>
bluff	<i>BN</i>	pluvial lake [relict]	<i>PQ</i>
delta	<i>DE</i>	raised beach	<i>RA</i>
delta plain (also LS)	<i>DC</i>	sabkha	<i>SAB</i>
flat	<i>FL</i>	salt marsh	<i>SM</i>
flood-plain playa	<i>FY</i>	shoal [relict]	<i>SE</i>
foredune	<i>FD</i>	shore	<i>SHO</i>
headland	<i>HE</i>	shore complex (also LS)	<i>SHC</i>
island (also LS)	<i>IS</i>	spit	<i>SP</i>
karst lake	<i>KAL</i>	stack [coast]	<i>SRC</i>
lagoon	<i>WI</i>	strand plain	<i>SS</i>
lagoon [relict]	<i>LAR</i>	till-floored lake plain	<i>TLP</i>
lake (w)	<i>WJ</i>	tombolo	<i>TO</i>
lakebed [relict]	<i>LBR</i>	water-lain moraine	<i>WM</i>
lake plain (also LS)	<i>LP</i>	wave-built terrace	<i>WT</i>
lakeshore	<i>LF</i>	wave-cut platform	<i>WP</i>
lake terrace	<i>LT</i>	wave-worked till plain	<i>WW</i>
longshore bar	<i>LON</i>		

### **Microfeatures:**

bar (also LF)	<i>BA</i>	ripple mark	<i>RM</i>
playa floor (also LF)	<i>PF</i>	shoreline	<i>SH</i>
playa rim (also LF)	<i>PR</i>	strandline	<i>SL</i>
playa slope (also LF)	<i>PSL</i>	swash zone	<i>SZ</i>
playa step (also LF)	<i>PST</i>	vernal pool	<i>VP</i>
playette	<i>PL</i>		

**3. FLUVIAL** (dominantly related to concentrated water flow [channel flow]; includes both erosional and depositional features with the exceptions of glaciofluvial landforms [see *Glacial*] and permanent water features [see *Water Bodies*]).

**Landscapes:**

alluvial plain	AP	delta plain (also LF)	DP
alluvial plain remnant	AR	dissected breaklands	DB
badlands	BA	fan piedmont	FP
bajada (also LF)	BJ	meander belt	MB
breaklands	BR	river valley (also LF)	RV
breaks	BK	scabland	SC
canyonlands	CL		

**Landforms:**

alluvial cone	AC	fan skirt	FI
alluvial fan	AF	flood plain	FP
alluvial flat	AP	flood-plain playa	FY
arroyo	AY	flood-plain splay	FM
axial stream (w)	AX	flood-plain step	FO
backswamp	BS	giant ripple	GC
bajada (also LS)	BJ	gorge	GO
bar (also Micro)	BR	gulch	GT
basin-floor remnant	BD	gut [valley]	GV
block stream	BX	inset fan	IF
box canyon	BOX	intermittent stream (also Micro)	INT
braided stream	BZ		
canyon	CA	levee	LV
channel	CC	meandering channel	MC
coulee	CE	meander scar	MS
cutoff	CV	meander scroll	MG
delta	DE	natural levee	NL
delta plain (also LS)	DC	overflow stream channel	OSC
drainageway	DQ	oxbow	OX
drainhead complex	DRC	paleoterrace	PTR
draw	DW	point bar	PR
ephemeral stream (also Micro)	EPS	ravine	RV
fan apron	FA	river valley (also LS)	RVV
fan collar	FCO	semi-open depression	SOD
fanhead trench	FF	slot canyon	SLC
fan remnant	FH	strath terrace	SU
		stream terrace	SX

terrace remnant	TER	valley-floor remnant	VFR
valley flat	VF	wash	WA
valley-border surfaces	VBS	wind gap	WG

### **Microfeatures:**

bar (also LF)	BA	gully	GU
bar and channel ( <i>patterned ground</i> )	BC	intermittent stream (also LF)	INT
channel	CH	ripple mark	RM
ephemeral stream (also LF)	ES	swash zone	SZ
groove	GR		

## **4. SOLUTION** (dominated by dissolution and, commonly, subsurface drainage).

### **Landscapes:**

cockpit karst	CPK	kegel karst	KK
cone karst	CK	sinkhole karst	SK
fluviokarst	FK	thermokarst	TK
glaciokarst	GK	tower karst	TW
karst	KR		

### **Landforms:**

blind valley	VB	pavement karst	PAV
cockpit	COC	pinnacle	PIN
collapse sinkhole	CSH	sinkhole	SH
interior valley	INV	solution platform	SOP
karst cone	KC	solution sinkhole	SOS
karstic marine terrace	KMT	swallow hole	TB
karst lake (w)	KAL	thermokarst depression (also Micro)	TK
karst tower	KTO	yardang (also Micro)	YD
karst valley	KVA	yardang trough (also Micro)	YDT
mogote	MOG		

### **Microfeatures:**

cutter	CU	solution pipe	SOP
karren	KA	thermokarst depression (also LF)	TK
solution chimney	SCH	yardang (also LF)	YD
solution corridor	SCO	yardang trough (also LF)	YDT
solution fissure	SOF		

## 5. EOLIAN (dominantly wind related; erosion or deposition).

### **Landscapes:**

dune field (also LF)	<i>DU</i>	sand plain	<i>SP</i>
sandhills	<i>SH</i>		

### **Landforms:**

barchan dune	<i>BQ</i>	longitudinal dune	<i>LDU</i>
blowout	<i>BY</i>	paha	<i>PA</i>
climbing dune	<i>CDU</i>	parabolic dune	<i>PB</i>
deflation basin	<i>DB</i>	parna dune	<i>PD</i>
deflation flat	<i>DFL</i>	playa dune (also Micro)	<i>PDU</i>
dune	<i>DU</i>	sabkha	<i>SAB</i>
dune field (also LS)	<i>DUF</i>	sand ramp	<i>SAR</i>
dune lake (w)	<i>DUL</i>	sand sheet	<i>RX</i>
dune slack (also Micro)	<i>DUS</i>	seif dune	<i>SD</i>
falling dune	<i>FDU</i>	slickrock (also Micro)	<i>SLK</i>
foredune	<i>FD</i>	star dune	<i>SDU</i>
interdune (also Micro)	<i>ID</i>	transverse dune	<i>TD</i>
loess bluff	<i>LO</i>	yardang (also Micro)	<i>YD</i>
loess hill	<i>LQ</i>	yardang trough (also Micro)	<i>YDT</i>

### **Microfeatures:**

dune slack (also LF)	<i>DS</i>	slickrock (also LF)	<i>SLK</i>
dune traces	<i>DT</i>	slip face	<i>SF</i>
interdune (also LF)	<i>ID</i>	yardang (also LF)	<i>YD</i>
playa dune (also LF)	<i>PD</i>	yardang trough (also LF)	<i>YDT</i>
playette	<i>PL</i>	zibar	<i>ZB</i>
shrub-coppice dune	<i>SCD</i>		

## 6. GLACIAL (directly related to glaciers; includes glaciofluvial, glaciolacustrine, glaciomarine, and outwash features).

### **Landscapes:**

continental glacier	<i>CG</i>	ice-margin complex	<i>IC</i>
drumlin field	<i>DF</i>	outwash plain (also LF)	<i>OP</i>
glaciokarst	<i>GK</i>	till plain (also LF)	<i>TP</i>
hills	<i>HI</i>		

**Landforms:**

alpine glacier	AG	kame terrace	KT
arete	AR	kettle	KE
cirque	CQ	lateral moraine	LM
cirque floor	CFL	medial moraine	MH
cirque headwall	CHW	moraine	MU
cirque platform	CPF	nearshore zone	NZ
col	CL	nearshore zone [relict]	NZR
collapsed ice-floored lakebed	CK	nunatak	NU
collapsed ice-walled lakebed	CN	outwash delta	OD
collapsed lake plain	CS	outwash fan	OF
collapsed outwash plain	CT	outwash plain (also LS)	OP
crag and tail	CAT	outwash terrace	OT
crevasse filling	CF	paha	PA
disintegration moraine	DM	pitted outwash plain	PM
drumlin	DR	pitted outwash terrace	POT
drumlinoid ridge	DRR	pothole (also Micro)	PH
end moraine	EM	pothole lake (intermittent water)	WN
esker	EK	proglacial lake (w)	WO
fjord (w)	FJ	proglacial lake [relict]	PGL
flute (also Micro)	FU	recessional moraine	RM
fosse	FV	roche moutonnée (also Micro)	RN
giant ripple	GC	rock glacier	RO
glacial drainage channel	GD	snowfield	SNF
glacial lake (w)	WE	stoss and lee	SAL
glacial lake [relict]	GL	swale (also Micro)	SC
glacial-valley floor	GVF	tarn (w; also Micro)	TAR
glacial-valley wall	GVW	terminal moraine	TA
glacier	GLA	till-floored lake plain	TLP
ground moraine	GM	till plain (also LS)	TP
hanging valley	HV	tunnel valley	TV
head-of-outwash	HD	tunnel-valley lake (w)	TVL
ice-contact slope	ICS	underfit stream	US
ice-marginal stream	IMS	U-shaped valley	UV
ice pressure ridge	IPR	valley train	VT
ice-pushed ridge	IPU	water-lain moraine	WM
interdrumlin	IDR	wave-worked till plain	WW
kame	KA		
kame moraine	KM		

**Microfeatures:**

flute (also LF)	<i>FL</i>	pothole (also LF)	<i>PH</i>
glacial groove	<i>GG</i>	roche moutonnée (also LF)	<i>POC</i>
ice wedge	<i>IWD</i>	swale (also LF)	<i>SW</i>
ice wedge cast	<i>IWC</i>	tarn (w; also LF)	<i>TN</i>
nivation hollow	<i>NH</i>		

**7. PERIGLACIAL** (related to nonglacial, cold climate [modern or relict], including periglacial forms of patterned ground.

**NOTE:** Consider “patterned ground” as a landform, but treat specific types of patterned ground [singular or plural] as microfeatures).

**Landscapes:**

coastal plain	<i>CP</i>	plains	<i>PL</i>
hills	<i>HI</i>	thermokarst	<i>TK</i>

**Landforms:**

alas	<i>AA</i>	pingo	<i>PI</i>
block field	<i>BW</i>	rock glacier	<i>RO</i>
muskeg	<i>MX</i>	string bog	<i>SY</i>
patterned ground (see <i>Microfeatures below for types</i> )	<i>PG</i>	thermokarst depression (also Micro)	<i>TK</i>
peat plateau	<i>PJ</i>	thermokarst lake (w)	<i>WV</i>

**Microfeatures:**

circle	<i>CI</i>	palsa (=peat hummock)	<i>PA</i>
earth hummock	<i>EH</i>	polygon	<i>PYG</i>
frost boil	<i>FB</i>	solifluction lobe	<i>SOL</i>
high-center polygon	<i>HCP</i>	solifluction sheet	<i>SS</i>
ice wedge	<i>IWD</i>	solifluction terrace	<i>ST</i>
ice wedge cast	<i>IWC</i>	sorted circle	<i>SCI</i>
ice-wedge polygon	<i>IWP</i>	stripe	<i>STR</i>
low-center polygon	<i>LCP</i>	thermokarst depression (also LF)	<i>TK</i>
nivation hollow	<i>NH</i>	turf hummock	<i>TH</i>
nonsorted circle	<i>NSC</i>		

**8. MASS MOVEMENT (=MASS WASTING)** (dominated by gravity, including creep forms; also see "Mass Movement Types" table, p. 5-8).

**Landscapes:** These generic landscapes are not mass movement features per se but are commonly modified by and include localized areas of mass movement.

breaklands	<i>BR</i>	hills	<i>HI</i>
dissected breaklands	<i>DB</i>	mountain range	<i>MR</i>
foothills	<i>FH</i>	mountains	<i>MO</i>

**Landforms:**

ash flow	<i>AS</i>	rockfall avalanche	<i>RFA</i>
avalanche chute	<i>AL</i>	rock glacier	<i>RO</i>
block glide	<i>BLG</i>	rock spread	<i>ROS</i>
block stream	<i>BX</i>	rock topple	<i>ROT</i>
colluvial apron	<i>COA</i>	rotational debris slide	<i>RDS</i>
complex landslide	<i>CLS</i>	rotational earth slide	<i>RES</i>
creep	<i>CRE</i>	rotational rock slide	<i>RRS</i>
debris avalanche	<i>DA</i>	rotational slide	<i>RTS</i>
debris fall	<i>DEF</i>	sag (also Micro)	<i>SAG</i>
debris flow	<i>DF</i>	sag pond (w; also Micro)	<i>SGP</i>
debris slide	<i>DS</i>	sand flow	<i>RW</i>
debris spread	<i>DES</i>	scree slope	<i>SCS</i>
debris topple	<i>DET</i>	slide	<i>SJ</i>
earthflow	<i>EF</i>	slump block	<i>SN</i>
earth spread	<i>ESP</i>	soil fall	<i>SOF</i>
earth topple	<i>ETO</i>	talus cone	<i>TC</i>
fall	<i>FB</i>	talus slope	<i>TAS</i>
flow	<i>FLO</i>	toe (also Micro)	<i>TOE</i>
lahar	<i>LA</i>	topple	<i>TOP</i>
landslide	<i>LK</i>	Toreva block	<i>TOR</i>
lateral spread	<i>LS</i>	translational debris slide	<i>TDS</i>
main scarp (also Micro)	<i>MAS</i>	translational earth slide	<i>TES</i>
mudflow	<i>MW</i>	translational rock slide	<i>TRS</i>
rockfall (also Micro)	<i>ROF</i>	translational slide	<i>TS</i>

**Microfeatures:**

main scarp (also LF)	<i>MAS</i>	solifluction lobe	<i>SOL</i>
minor scarp	<i>MIS</i>	solifluction sheet	<i>SS</i>
rockfall (also LF)	<i>ROF</i>	solifluction terrace	<i>ST</i>
sag (also LF)	<i>SAG</i>	terraces	<i>TER</i>
sag pond (w; also LF)	<i>SP</i>	toe (also LF)	<i>TOE</i>
sand boil	<i>SB</i>		

## 9. VOLCANIC and HYDROTHERMAL

### **Landscapes:**

caldera (also LF)	<i>CD</i>	lava plateau (also LF)	<i>LL</i>
foothills	<i>FH</i>	mountains	<i>MO</i>
hills	<i>HI</i>	shield volcano (also LF)	<i>SV</i>
lava field (also LF)	<i>LF</i>	volcanic field (also LF)	<i>VF</i>
lava plain (also LF)	<i>LV</i>		

### **Landforms:**

aa lava flow	<i>ALF</i>	lava trench (also Micro)	<i>LTR</i>
ash field	<i>AQ</i>	lava tube	<i>LTU</i>
ash flow	<i>AS</i>	louderback	<i>LU</i>
block lava flow	<i>BLF</i>	maar	<i>MAA</i>
caldera (also LS)	<i>CD</i>	mawae	<i>MAW</i>
cinder cone	<i>CI</i>	mud pot	<i>MP</i>
diatreme	<i>DT</i>	pahoehoe lava flow	<i>PAF</i>
dike	<i>DK</i>	pillow lava flow	<i>PIF</i>
fissure vent	<i>FIV</i>	plug dome	<i>PP</i>
geyser	<i>GE</i>	pyroclastic flow	<i>PCF</i>
geyser basin	<i>GEB</i>	pyroclastic surge	<i>PCS</i>
geyser cone	<i>GEC</i>	shield volcano (also LS)	<i>SHV</i>
hot spring	<i>HP</i>	steptoe	<i>ST</i>
kipuka	<i>KIP</i>	stratovolcano	<i>SV</i>
lahar	<i>LA</i>	volcanic cone	<i>VC</i>
lava dome	<i>LD</i>	volcanic crater	<i>CR</i>
lava field (also LS)	<i>LFI</i>	volcanic dome	<i>VD</i>
lava flow	<i>LC</i>	volcanic field (also LS)	<i>VOF</i>
lava flow unit (also Micro)	<i>LFU</i>	volcanic neck	<i>VON</i>
lava plain (also LS)	<i>LN</i>	volcanic pressure ridge (also Micro)	<i>PU</i>
lava plateau (also LS)	<i>LL</i>	volcano	<i>VO</i>

### **Microfeatures:**

corda	<i>CO</i>	spiracle	<i>SPI</i>
lava flow unit (also LF)	<i>LFU</i>	tumulus (pl.: tumuli)	<i>TU</i>
lava trench (also LF)	<i>LT</i>	volcanic pressure ridge (also LF)	<i>VPR</i>
spatter cone	<i>SPC</i>		



**10. TECTONIC and STRUCTURAL** (related to regional or local bedrock structures or crustal movement; recognized only if expressed at or near the land surface).

**Landscapes:**

basin floor	<i>BC</i>	mountain range	<i>MR</i>
batholith	<i>BL</i>	mountains	<i>MO</i>
bolson	<i>BO</i>	mountain system	<i>MS</i>
breached anticline (also LF)	<i>BD</i>	pediment slope	<i>PS</i>
dissected plateau	<i>DP</i>	plateau (also LF)	<i>PT</i>
fault-block mountains	<i>FM</i>	rift valley	<i>RF</i>
fold-thrust hills	<i>FTH</i>	semi-bolson	<i>SB</i>
foothills	<i>FH</i>	tableland	<i>TB</i>
hills	<i>HI</i>	valley	<i>VA</i>
intermontane basin	<i>IB</i>		

**Landforms:**

anticline	<i>AN</i>	hogback	<i>HO</i>
breached anticline (also LS)	<i>BRL</i>	homoclinal ridge	<i>HCR</i>
canyon bench	<i>CYB</i>	homocline	<i>HC</i>
cueta	<i>CU</i>	horst	<i>HT</i>
cueta valley	<i>CUV</i>	louderback	<i>LU</i>
diapir	<i>DD</i>	meteorite crater	<i>MEC</i>
dike	<i>DK</i>	monocline	<i>MJ</i>
dip slope	<i>DL</i>	rock pediment	<i>ROP</i>
dome	<i>DO</i>	scarp slope	<i>RS</i>
fault block	<i>FAB</i>	sill	<i>RT</i>
fault-line scarp	<i>FK</i>	stock	<i>STK</i>
fault zone	<i>FAZ</i>	strike valley	<i>STV</i>
fold	<i>FQ</i>	structural bench	<i>SB</i>
graben	<i>GR</i>	syncline	<i>SZ</i>
half graben	<i>HG</i>	window	<i>WIN</i>

**Microfeatures:**

sand boil	<i>SB</i>
-----------	-----------

**11. SLOPE** (generic terms [e.g., hill] or those that describe slope form, geometry, or arrangement of land features rather than any particular genesis or process).

**Landscapes:**

badlands	BA	mountain range	MR
breached anticline (also LF)	BD	mountains	MO
breaklands	BR	mountain system	MS
breaks	BK	piedmont	PI
canyonlands	CL	piedmont slope	PS
dissected breaklands	DB	plains (singular=LF)	PL
dissected plateau	DI	plateau (also LF)	PT
fault-block mountains	FM	tableland	TB
foothills	FH	upland	UP
hills (singular=LF)	HI		

**Landforms:**

beveled base	BVB	interfluve ( <i>also Geom. Component—Hills</i> )	IV
block stream	BX		
bluff	BN	knob	KN
breached anticline (also LS)	BRL	knoll	KL
broad interstream divide	BID	ledge (also Micro)	LE
butte	BU	low hill	LH
canyon bench	CYB	mesa	ME
canyon wall	CW	mountain (plural=LS)	MM
cliff	CJ	mountain slope	MN
colluvial apron	COA	mountain valley	MV
cuesta	CU	notch	NO
dip slope	DL	paha	PA
dome	DO	peak	PK
escarpment	ES	pediment	PE
faceted spur	FS	plain (plural=LS)	PN
fault block	FAB	plateau (also LS)	PT
fault-line scarp	FK	ridge	RI
free face ( <i>also Geom. Component—Hills, Mountains</i> )	FW	rim	RJ
gap	GA	rise (also Micro) ( <i>also Geom. Component—Flat Plains</i> )	RIS
headwall	HW	rock pediment	ROP
high hill	HH	scarp	RY
hill (plural=LS)	HI	scarp slope	RS
hillslope	HS	scree slope	SCS
hogback	HO	slickrock (also Micro)	SLK

spur	<i>SQ</i>	tor	<i>TQ</i>
stack [geom.]	<i>SR</i>	valley	<i>VA</i>
structural bench	<i>SB</i>	valley-floor remnant	<i>VFR</i>
talus cone	<i>TC</i>	wind gap	<i>WG</i>
talus slope	<i>TAS</i>		

***Microfeatures:***

finger ridge	<i>FR</i>	rise (also LF) ( <i>also Geom. Component—Flat Plains</i> )	<i>RIS</i>
mound	<i>MO</i>		
rib	<i>RB</i>	slickrock (also LF)	<i>SLK</i>
rill	<i>RL</i>		

**12. EROSIONAL** (related dominantly to water erosion but excludes perennial, concentrated channel flow [i.e. fluvial, glaciofluvial] or eolian erosion).

**Landscapes:**

badlands	BA	hills	HI
breached anticline (also LF)	BD	mountain range	MR
breaklands	BR	mountains	MO
breaks	BK	piedmont	PI
canyonlands	CL	piedmont slope	PS
dissected breaklands	DB	plateau (also LF)	PT
dissected plateau	DI	tableland	TB
foothills	FH		

**Landforms:**

ballena	BL	monadnock	MD
ballon	BV	notch	NO
basin floor remnant	BD	paha	PA
beveled base	BVB	partial ballena	PF
breached anticline (also LS)	BRL	peak	PK
canyon bench	CYB	pediment	PE
canyon wall	CW	plateau (also LS)	PT
col	CL	rock pediment	ROP
colluvial apron	COA	sabkha	SAB
cuesta	CU	saddle	SA
cuesta valley	CUV	scarp slope	RS
eroded fan remnant	EFR	slickrock (also Micro)	SLK
eroded fan-remnant	EFS	stack [geom.]	SR
sideslope		strike valley	STV
erosion remnant	ER	structural bench	SB
free face (also <i>Geom. Component—Hills, Mountains</i> )	FW	terrace remnant	TER
gap	GA	tor	TQ
hogback	HO	valley-border surfaces	VBS
inselberg	IN	valley-floor remnant	VFR
		wind gap	WG
		window	WIN

**Microfeatures:**

earth pillar	EP	pinnacle	PI
finger ridge	FR	rib	RB
groove	GR	rill	RL
gully	GU	slickrock (also LF)	SLK
hoodoo	HO	swale	SW

**13. DEPRESSIONAL** (low areas or declivity features, excluding permanent water bodies).

**Landscapes:**

basin	<i>BS</i>	breaklands	<i>BR</i>
basin floor (also LF)	<i>BC</i>	dissected breaklands	<i>DB</i>
bolson	<i>BO</i>	semi-bolson	<i>SB</i>
breached anticline (also LF)	<i>BD</i>	valley	<i>VA</i>

**Landforms:**

alluvial flat	<i>AP</i>	mountain valley	<i>MV</i>
basin floor (also LS)	<i>BC</i>	open depression (also Micro)	<i>ODE</i>
basin floor remnant	<i>BD</i>		
box canyon	<i>BOX</i>	playa	<i>PL</i>
breached anticline (also LS)	<i>BRL</i>	playa floor (also Micro)	<i>PFL</i>
canyon	<i>CA</i>	playa rim (also Micro)	<i>PRI</i>
canyon wall	<i>CW</i>	playa slope (also Micro)	<i>PSL</i>
Carolina Bay	<i>CB</i>	playa step (also Micro)	<i>PST</i>
closed depression (also Micro)	<i>CLD</i>	pothole (also Micro)	<i>PH</i>
col	<i>CL</i>	pothole lake (intermittent water)	<i>WN</i>
coulee	<i>CE</i>	ravine	<i>RV</i>
cove	<i>CO</i>	sabkha	<i>SAB</i>
cuesta valley	<i>CUV</i>	saddle	<i>SA</i>
depression	<i>DP</i>	sag (also Micro)	<i>SAG</i>
drainageway	<i>DQ</i>	semi-open depression	<i>SOD</i>
drainhead complex	<i>DRC</i>	slot canyon	<i>SLC</i>
gap	<i>GA</i>	strike valley	<i>STV</i>
gorge	<i>GO</i>	swale (also Micro)	<i>SC</i>
gulch	<i>GT</i>	trough	<i>TR</i>
gut [valley]	<i>GV</i>	U-shaped valley	<i>UV</i>
intermontane basin	<i>IB</i>	valley	<i>VA</i>
kettle	<i>KE</i>	valley floor	<i>VL</i>
		V-shaped valley	<i>VV</i>

**Microfeatures:**

closed depression (also LF)	<i>CD</i>	playette	<i>PL</i>
open depression (also LF)	<i>OP</i>	pothole (also LF)	<i>PH</i>
playa floor (also LF)	<i>PF</i>	sag (also LF)	<i>SAG</i>
playa rim (also LF)	<i>PR</i>	semi-open depression	<i>SOD</i>
playa slope (also LF)	<i>PSL</i>	swale (also LF)	<i>SW</i>
playa step (also LF)	<i>PST</i>	tree-tip pit	<i>TTP</i>

**14. WETLANDS** (related to vegetated and/or shallow water areas and wet soils). (Provisional list: conventional, geologic definitions; not legalistic or regulatory usage.)

**Landscapes:**

estuary (also LF)	<i>ES</i>	everglades	<i>EG</i>
-------------------	-----------	------------	-----------

**Landforms:**

alas	<i>AA</i>	oxbow lake (ephemeral water)	<i>WK</i>
backswamp	<i>BS</i>	peat plateau	<i>PJ</i>
bog	<i>BO</i>	playa (intermittent water)	<i>PL</i>
Carolina Bay	<i>CB</i>	pocosin	<i>PO</i>
dune slack (also Micro)	<i>DUS</i>	pothole (also Micro)	<i>PH</i>
ephemeral stream (also Micro)	<i>EPS</i>	pothole lake (w)	<i>WN</i>
estuary (also LS)	<i>WD</i>	raised bog	<i>RB</i>
fen	<i>FN</i>	ribbed fen	<i>RG</i>
flood-plain playa	<i>FY</i>	sabkha	<i>SAB</i>
fringe-tidal marsh	<i>FTM</i>	salt marsh	<i>SM</i>
highmoor bog	<i>HB</i>	seep (also Micro)	<i>SEE</i>
intermittent stream (also Micro)	<i>INT</i>	semi-open depression	<i>SOD</i>
lowmoor bog	<i>LX</i>	slough (intermittent water)	<i>SL</i>
mangrove swamp	<i>MAN</i>	string bog	<i>SY</i>
marsh	<i>MA</i>	swamp	<i>SW</i>
muskeg	<i>MX</i>	tidal flat	<i>TF</i>
		tidal marsh	<i>TM</i>

**Microfeatures:**

dune slack (also LF)	<i>DS</i>	playette	<i>PL</i>
ephemeral stream (also LF)	<i>ES</i>	pothole (also LF)	<i>PH</i>
intermittent stream (also LF)	<i>INT</i>	vernal pool (seasonal water)	<i>VP</i>

**15. WATER BODIES** (discrete “surface water” features; primarily permanent open water, which in soil survey reports is commonly treated as the generic map unit “water” [e.g., lake] or as a spot/line symbol [e.g., perennial stream]).

**Landscapes:**

bay [coast] (also LF)	<i>BY</i>	ocean	<i>OC</i>
estuary (also LF)	<i>ES</i>	sea (also LF)	<i>SEA</i>
gulf (also LF)	<i>GU</i>	sound (also LF)	<i>SO</i>
lagoon (also LF)	<i>LG</i>	strait (also LF)	<i>ST</i>

**Landforms:**

axial stream	<i>AX</i>	playa lake	<i>WL</i>
bay [coast] (also LS)	<i>BAY</i>	pluvial lake	<i>PLL</i>
bayou	<i>WC</i>	pothole lake	<i>WN</i>
cove [water]	<i>COW</i>	proglacial lake	<i>WO</i>
dune lake	<i>DUL</i>	river	<i>RIV</i>
estuary (also LS)	<i>WD</i>	sag pond (also Micro)	<i>SGP</i>
fjord	<i>FJ</i>	salt pond (also Micro)	<i>WQ</i>
glacial lake	<i>WE</i>	sea (also LS)	<i>SEA</i>
gulf (also LS)	<i>GU</i>	shoal	<i>WR</i>
gut [channel] (also Micro)	<i>WH</i>	slackwater	<i>WS</i>
ice-marginal stream	<i>IMS</i>	slough	<i>SL</i>
inlet	<i>IL</i>	sound (also LS)	<i>SO</i>
lagoon (also LS)	<i>WI</i>	strait (also LS)	<i>STT</i>
lagoon channel	<i>LCH</i>	stream (permanent water)	<i>STR</i>
lake	<i>WJ</i>	tarn (also Micro)	<i>TAR</i>
marine lake	<i>ML</i>	thermocarst lake	<i>WV</i>
nearshore zone	<i>NZ</i>	tidal inlet	<i>TI</i>
oxbow lake	<i>WK</i>	tidal inlet [relict]	<i>TIR</i>
perennial stream (also Micro)	<i>PS</i>	tunnel-valley lake	<i>TVL</i>

**Microfeatures:**

channel (permanent water)	<i>CH</i>	sag pond (also LF)	<i>SP</i>
gut [channel] (also LF)	<i>WH</i>	salt pond (also LF)	<i>WQ</i>
perennial stream (also LF)	<i>PS</i>	tank	<i>TA</i>
pond	<i>PON</i>	tarn (also LF)	<i>TN</i>
pool	<i>POO</i>		

**16. SUBAQUEOUS FEATURES** (discrete underwater features [that commonly can support rooted plants] and adjacent features ordinarily found below permanent open water; *historically, in soil survey reports these underwater features have been included in the generic map unit "water."* Subaqueous "Landscape" terms are obviously not terrestrial but are earth-surface features).

**Landscapes:**

bay [coast] (w; also LF)	BY	ocean (w)	OC
estuary (w; also LF)	ES	sea (w; also LF)	SEA
gulf (w; also LF)	GU	sound (w; also LF)	SO
lagoon (w; also LF)	LA	strait (w; also LF)	ST

**Landforms:**

barrier cove	BAC	marine lake	ML
bay [coast] (w; also LS)	BAY	nearshore zone	NZ
bay bottom	BOT	reef	RF
cove [water] (w)	CO	sea (w; also LS)	SEA
estuary (also LS)	WD	shoal	WR
flood-tidal delta	FTD	sound (w; also LS)	SO
flood-tidal delta flat	FTF	strait (w; also LS)	STT
flood-tidal delta slope	FTS	submerged back-barrier beach	SBB
fluviomarine bottom	FMB	submerged mainland beach	SMB
gulf (w; also LS)	GU	submerged point bar [coast]	SPB
inlet	IL	submerged wave-built terrace	SWT
lagoon (also LS)	WI	submerged wave-cut platform	SWP
lagoon bottom	LBO	tidal inlet	TI
lagoon channel	LCH	tidal inlet [relict]	TIR
lake	WJ	washover-fan flat	WFF
lakebed (w)	LB	washover-fan slope	WFS
longshore bar	LON		
mainland cove	MAC		

**Microfeatures:**

channel (permanent water)	CH	gut [channel] (w)	WH
---------------------------	----	-------------------	----


**Anthropogenic Features:**

dredge-deposit shoal	DDS	dredged channel	DC
----------------------	-----	-----------------	----



## PART III: SURFACE MORPHOMETRY

- A) **Elevation:** The height of a point on the earth's surface relative to mean sea level (msl); indicate units; e.g., *106 m* or *348 ft*.
- B) **Slope Aspect:** The compass bearing (in degrees, corrected for declination) that a slope faces, viewed downslope; e.g., *287°*.
- C) **Slope Gradient:** The angle of the ground surface (in percent) through the site and in the direction that overland water would flow (commonly referred to as slope); e.g., *18%*.
- D) **Slope Complexity:** Describe the relative uniformity (smooth linear or curvilinear=*simple* or *S*) or irregularity (*complex* or *C*) of the ground surface leading downslope through the point of interest; e.g., *simple* or *S*.




- E) **Relative Slope Segment Position** (called **geomorph\_slope\_segment** in NASIS): If useful to subdivide long slopes, describe the relative slope location of the area of interest.

Relative Slope Segment Position	Code	Criteria
lower third	LT	on lower third
middle third	MT	on middle third
upper third	UT	on upper third


- F) **Slope Shape:** Slope shape is described in two directions: 1) up and down slope (perpendicular or “normal” to the contour; called **slope\_down** in NASIS); and 2) across slope (along the horizontal contour; called **slope\_across** in NASIS). These two descriptors are commonly reported as a pair. The first term refers to up and down slope (or vertical), and the second term refers to across slope; e.g., *Linear, Convex, or LV*.

Down Slope (Vertical)	Across Slope (Horizontal)	Code
concave	concave	CC
concave	convex	CV
concave	linear	CL
convex	concave	VC
convex	convex	VV
convex	linear	VL
linear	concave	LC
linear	convex	LV
linear	linear	LL




- G) **Hillslope—Profile Position** (commonly called Hillslope Position): Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., *backslope* or *BS*. This set of terms is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS




- H) **Geomorphic Component** (Geomorphic Position in PDP): Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Other unique descriptors are available for Hills, Terraces and Stepped Landforms, Mountains, and Flat Plains; e.g. (for Hills), *nose slope* or *NS*.


1) Hills	Code
interfluve	IF
crest	CT
head slope	HS
nose slope	NS
side slope	SS
free face	FF
base slope	BS




2) Terraces and Stepped Landforms	Code
riser	RI
tread	TR



3) Mountains	Code
mountaintop	MT
mountainflank	MF
upper third – mountainflank	UT
center third – mountainflank	CT
lower third – mountainflank	LT
free face	FF
mountainbase	MB



4) Flat Plains	Code
dip	DP
rise	RI
talf	TF




- I) **Microrelief:** Small, relative differences in elevation between adjacent areas on the earth's surface; e.g., *microhigh* or *MH*.


<b>Microrelief</b>	<b>Code</b>
microhigh	MH
microlow	ML
microslope	MS

**NOTE:** See graphic p. 2-54.

- J) **Drainage Pattern:** The arrangement of drainage channels on the land surface; also called drainage network.

<b>Drainage Pattern</b>	<b>Code</b>
annular	AN
artificial	AR
centripetal	CE
dendritic	DN
deranged	DR
karst	KA
parallel	PA
pinnate	PI
radial	RA
rectangular	RE
thermokarst	TH
trellis	TR





## REFERENCES

Fenneman, N.M. 1931. Physiography of the western United States. McGraw-Hill Co., New York, NY.

Fenneman, N.M. 1938. Physiography of the eastern United States. McGraw-Hill Co., New York, NY.

Fenneman, N.M. 1946 (reprinted 1957). Physical divisions of the United States. U.S. Geological Survey. U.S. Gov. Print. Office, Washington, DC. 1 sheet; 1:7,000,000.

Ruhe, R.V. 1975. Geomorphology: Geomorphic processes and surficial geology. Houghton-Mifflin, Boston, MA.

Schoeneberger, P.J., and D.A. Wysocki. 1996. Geomorphic descriptors for landforms and geomorphic components: Effective models and weaknesses. *In* Agronomy abstracts, ASA, Madison, WI.

Soil Survey Staff. 2012. Glossary of landform and geologic terms. Part 629, National soil survey handbook (NSSH) [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Wahrhaftig, C. 1965. Physiographic divisions of Alaska. U.S. Geol. Surv. Prof. Pap. 482.

Wysocki, D.A., P.J. Schoeneberger, D. Hirmas, and H.E. LaGarry. 2012. Geomorphology of soil landscapes. *In* P.M. Huang et al. (ed.) Handbook of soil science: Properties and processes, 2nd ed. CRC Press, Taylor and Francis Group, LLC, Boca Raton, FL. ISBN: 978-1-4398-0305-9.

Wysocki, D.A., P.J. Schoeneberger, and H.E. LaGarry. 2000. Geomorphology of soil landscapes. *In* M.E. Sumner (ed.) Handbook of soil science. CRC Press, LLC, Boca Raton, FL. ISBN: 0-8493-3136-6.



# SOIL TAXONOMY

*P.J. Schoeneberger, D.A. Wysocki, and J.V. Chiaretti, NRCS, Lincoln, NE*

## INTRODUCTION

---

The purpose of this section is to expand upon and augment the abbreviated soil taxonomic contents of the "Profile/Pedon Description" section. Complete definitions are found in *Keys to Soil Taxonomy* (Soil Survey Staff, 2010a).

## HORIZON AND LAYER DESIGNATIONS

---

**NOTE:** Horizons are considered to be layers of pedogenically derived or modified material. Layers are deemed to be zones of nonpedogenically derived/modified material (e.g., geologic strata).

### MASTER AND TRANSITIONAL HORIZONS or LAYERS—

Horizon	Criteria <sup>1</sup>
<b>O</b>	Organic soil materials other than limnic materials. The mineral fraction is a small percent by volume and is <80% by weight.
<b>A</b>	Mineral soil, formed at the surface or below an O horizon, little remnant rock structure, and one or more: 1) accumulation of humified organic matter but dominated by mineral matter, and not dominated by E or B horizon properties; 2) properties resulting from cultivation, pasturing, or similar disturbance; or 3) morphology resulting from surficial processes different from the underlying B or C. Excludes recent eolian or alluvial deposits that retain stratification.
<b>AB</b> or <b>AE</b> or <b>AC</b>	Dominantly A horizon characteristics but also contains some B, E, or C horizon attributes.
<b>A/B</b> or <b>A/E</b> or <b>A/C</b>	Discrete, intermingled bodies of two horizons: A material dominates with lesser but discrete bodies of B, E, or C material.
<b>E</b>	Mineral soil with some loss of silicate clay, iron, aluminum, and/or organic matter leaving a net concentration of sand and silt; little or no remnant rock structure; typically lighter color (higher value, chroma) and coarser texture than A.

Horizon	Criteria <sup>1</sup>
<b>EA</b> or <b>EB</b> or <b>EC</b>	Dominantly E horizon characteristics but also contains some A, B, or C horizon attributes.
<b>E/A</b> or <b>E/B</b>	Discrete, intermingled bodies of two horizons: E material dominates with lesser but discrete bodies of A or B material.
<b>E and Bt</b> <b>B and E</b>	Thin, heavier textured lamellae (Bt) within a predominantly E horizon with less clay (or thin E layers within a predominantly B horizon).
<b>BA</b> or <b>BE</b> or <b>BC</b>	Dominantly B characteristics but also contains some A, E, or C horizon attributes.
<b>B/A</b> or <b>B/E</b> or <b>B/C</b>	Discrete, intermingled bodies of two horizons: B material dominates with lesser but discrete bodies of A, E, or C material.
<b>B</b>	<p>Mineral soil, typically formed below O, A, or E; little or no rock structure; and with one or more of the following:</p> <ol style="list-style-type: none"> <li>1) illuvial accumulation of silicate clay, Fe, Al, humus, carbonate, gypsum, silica, or salt more soluble than gypsum (one or more);</li> <li>2) removal, addition, or transformation of carbonates, gypsum, or more soluble salts;</li> <li>3) residual concentration of oxides, sesquioxides, and silicate clays (one or more);</li> <li>4) sesquioxide coatings;</li> <li>5) alterations that form silicate clays or liberate oxides and form pedogenic structure;</li> <li>6) Brittleness;</li> <li>7) Strong gleying in the presence of aquic conditions (or artificial drainage); layers with gleying but no other pedogenic change are not B horizons. Most B horizons are or were subsurface horizons. Some formed at the surface by accumulation of evaporites. Cemented and brittle layers that have other evidence of pedogenesis are included as B horizons.</li> </ol>

<b>Horizon</b>	<b>Criteria <sup>1</sup></b>
<b>CB</b> or <b>CA</b>	Dominantly C horizon characteristics but also contains some B or A horizon attributes.
<b>C/B</b> or <b>C/A</b>	Discrete, intermingled bodies of two horizons: C material dominates, with lesser but discrete bodies of A or B material.
<b>C</b>	Mineral soil, soft bedrock (excluding <i>Strongly Cemented</i> to <i>Indurated</i> bedrock unless highly cracked); layer little affected by pedogenesis and lacks properties of O, A, E, or B horizons. May or may not be parent material of the solum.
<b>L</b>	Limnic soil materials. Sediments deposited in a body of water (subaqueous) and dominated by organic materials (aquatic plant and animal fragments and fecal material) and lesser amounts of clay.
<b>W</b>	A layer of liquid water (W) or permanently frozen ice (Wf), within or beneath the soil ( <u>excludes</u> water / ice above soil).
<b>M</b>	Root-limiting subsoil layers of human-manufactured materials; e.g., geotextile liner.
<b>R</b>	Hard bedrock (continuous, coherent <i>Strongly Cemented</i> to <i>Indurated</i> Cementation Classes).

<sup>1</sup> Soil Survey Staff, 2010a.

**HORIZON SUFFIXES**—Historically referred to as “Horizon Subscripts,” “Subordinate Distinctions,” <sup>1</sup> “Horizon\_Designation\_Suffix” in NASIS, and “Suffix Symbols” in *Keys to Soil Taxonomy*. <sup>2</sup> (Historical nomenclature and conversions are shown in the tables on page 4-6.)

<b>Horizon Suffixes</b>	<b>Criteria <sup>2</sup></b>
<b>a</b>	Highly decomposed organic matter (OM); rubbed fiber content <17% (by vol.); used only with O (see <b>e</b> , <b>i</b> ).
<b>aa</b> <sup>3</sup>	(proposed) Accumulation of anhydrite (CaSO <sub>4</sub> ).
<b>b</b>	Buried genetic horizon (not used with organic materials or to separate organic from mineral materials).
<b>c</b>	Concretions or nodules; significant accumulation of <i>cemented</i> bodies enriched with Fe, Al, Mn, Ti (cement not specified except <i>excludes</i> a predominance of silica [see <b>q</b> ]); not used for carbonates or soluble salts (see <b>z</b> ).

<b>Horizon Suffixes</b>	<b>Criteria <sup>2</sup></b>
<b>co</b>	Coprogenous earth (used only with L); organic materials deposited under water and dominated by fecal material from aquatic animals.
<b>d</b>	<i>Physical</i> root restriction due to high bulk density (natural or human-induced conditions; e.g., lodgment till, plow pans.
<b>di</b>	Diatomaceous earth (used only with L); materials deposited under water and dominated by the siliceous diatom remains.
<b>e</b>	Moderately (intermediately) decomposed organic matter; rubbed fiber content 17 to <40% (by vol.); used only with O (see <b>a, i</b> ).
<b>f</b>	Permafrost (permanently frozen subsurface soil or ice); excludes seasonally frozen ice and surface ice.
<b>ff</b>	Dry permafrost (permanently frozen soil; not used for seasonally frozen soil; no continuous ice bodies [see <b>f</b> ]).
<b>g</b>	Strong gley (Fe reduced and pedogenically removed); typically $\leq 2$ chroma; may have other redoximorphic features (RMF); not used for geogenic gray colors.
<b>h</b>	Illuvial organic matter (OM) accumulation (with B: accumulation of illuvial, amorphous OM-sesquioxide complexes); coats sand and silt particles and may fill pores; use <i>Bhs</i> if significant accumulation of sesquioxides and moist chroma <i>and</i> value $\leq 3$ .
<b>i</b>	Slightly decomposed organic matter; rubbed fiber content $\geq 40\%$ (by vol.); used only with O (see <b>a, e</b> ).
<b>j</b>	Jarosite accumulation; e.g., acid sulfate soils.
<b>jj</b>	Evidence of cryoturbation; e.g., irregular or broken horizon boundaries, sorted rock fragments (patterned ground), or OM in lower boundary between active layer and permafrost layer.
<b>k</b>	Pedogenic carbonate accumulation (e.g., $\text{CaCO}_3$ ; <50% by vol.).
<b>kk</b>	Major pedogenic carbonate accumulation; soil fabric is plugged $\approx$ continuous ( $\geq 50\%$ by vol. estimated).
<b>m</b>	Continuous pedogenic cementation or induration (>90% cemented, even if fractured); physically root restrictive. Dominant cement type can be indicated by additional letters; e.g., <i>km</i> or <i>kkm</i> —carbonates, <i>qm</i> —silica, <i>kqm</i> —carbonates and silica, <i>sm</i> —iron, <i>yym</i> —gypsum, <i>zm</i> —salts more soluble than gypsum.

<b>Horizon Suffixes</b>	<b>Criteria <sup>2</sup></b>
<b>ma</b>	Marl (used only with L); materials deposited under water and dominated by a mixture of clay and CaCO <sub>3</sub> ; typically gray or beige.
<b>n</b>	Pedogenic exchangeable sodium accumulation.
<b>o</b>	Residual accumulation of sesquioxides.
<b>p</b>	Tillage or other disturbance of surface layer (pasture, plow, etc.). Designate <i>Op</i> for disturbed organic surface, <i>Ap</i> for mineral surface even if the layer clearly was originally an E, B, C, etc.
<b>q</b>	Accumulation of secondary (pedogenic) silica.
<b>r</b>	Used with C to indicate weathered or soft bedrock (root-restrictive saprolite or soft bedrock), such as weathered or partially consolidated sandstone, siltstone, or shale; materials are sufficiently incoherent to allow hand digging with a spade (Excavation Difficulty classes are <i>Low</i> to <i>High</i> ). Roots only penetrate along joint planes.
<b>s</b>	Significant illuvial accumulation of amorphous, dispersible sesquioxides and organic matter complexes and moist color value or chroma $\geq 4$ . Used with B horizon; used with h as <i>Bhs</i> if moist color value and chroma are $\leq 3$ .
<b>se</b>	Presence of sulfides (in mineral or organic horizons). Typically dark colors (e.g., value $\leq 4$ , chroma $\leq 2$ ); may have a sulfurous odor.
<b>ss</b>	Slickensides; e.g., oblique shear faces 20-60° off horizontal; caused by shrink-swell clay action; wedge-shaped peds and seasonal surface cracks also are commonly present.
<b>t</b>	Accumulation (by translocation or illuviation) of silicate clays (clay films, lamellae, or clay bridging in some part of the horizon).
<b>u</b>	Presence of human-manufactured materials (artifacts); e.g., asphalt, bricks, plastic, glass, metals, construction debris, garbage.
<b>v</b>	Plinthite (high Fe, low OM, reddish contents; firm or very firm moist consistence; irreversible hardening with repeated wetting and drying).
<b>w</b>	Incipient color or pedogenic structure development, minimal illuvial accumulations; used only with B horizons, excluded from use with transition horizons.

<b>Horizon Suffixes</b>	<b>Criteria <sup>2</sup></b>
<b>x</b>	Fragipan or fragic characteristics (pedogenetically developed brittleness, firmness, bleached prisms, high bulk density, root restrictive).
<b>y</b>	Accumulation of gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ); <50% by volume (estimated).
<b>yy</b>	Dominance of gypsum ( $\approx \geq 50\%$ by vol. estimated); light colored (e.g., value $\geq 7$ , chroma $\leq 4$ ); may be pedogenetically derived or inherited transformation of primary gypsum from parent material.
<b>z</b>	Pedogenic accumulation of salts more soluble than gypsum; e.g., NaCl.

<sup>1</sup> Soil Survey Division Staff, 1993.

<sup>2</sup> Soil Survey Staff, 2010a.

<sup>3</sup> Personal communication with Soil Survey Standards Staff, 2012.

### **HORIZON AND LAYER DESIGNATIONS CONVERSION**

**CHARTS**—(**NOTE:** Gray boxes indicate the year the convention was first adopted.)

<b>Master Horizons, Layers, or Combinations</b>				
<b>1951 <sup>1</sup></b>	<b>1962 <sup>2</sup>, 1975 <sup>3</sup></b>	<b>1982 <sup>4</sup></b>	<b>1998 <sup>5</sup></b>	<b>2006 <sup>6</sup>, 2010 <sup>7</sup></b>
Aoo or Ao	O	O	O	O
Aoo	O1	Oi and/or Oe	Oi and/or Oe	Oi and/or Oe
Ao	O2	Oe and/or Oa	Oe and/or Oa	Oe and/or Oa
—	—	Oi	Oi	Oi
—	—	Oe	Oe	Oe
—	—	Oa	Oa	Oa
A	A	A	A	A
A1	A1	A	A	A
A2	A2	E	E	E
A3	A3	AB or EB	AB or EB	AB or EB
AB	AB	—	—	—
A&B	A&B	A/B or E/B	A/B or E/B	A/B or E/B
AC	AC	AC	AC	AC
—	—	E and Bt	E and Bt	E and Bt

<b>Master Horizons, Layers, or Combinations</b>				
<b>1951</b> <sup>1</sup>	<b>1962</b> <sup>2</sup> , <b>1975</b> <sup>3</sup>	<b>1982</b> <sup>4</sup>	<b>1998</b> <sup>5</sup>	<b>2006</b> <sup>6</sup> , <b>2010</b> <sup>7</sup>
B	B	B	B	B
B1	B1	BA or BE	BA or BE	BA or BE
B&A	B&A	B/A or B/E	B/A or B/E	B/A or B/E
B2	B2	B or Bw	B or Bw	B or Bw
G	g <sup>8</sup>	Ag, Bg, Cg	Ag, Bg, Cg	Ag, Bg, Cg
B3	B3	BC or CB	BC or CB	BC or CB
—	—	B/C, C/B, C/A	B/C, C/B, C/A	B/C, C/B, C/A
C	C	C	C	C
Cca	Cca	<i>Bk</i>	<i>Bk</i>	<i>Bk, Bkk</i> <sup>6</sup>
Ccs	Ccs	<i>By, Cy</i>	<i>By, Cy</i>	<i>By</i> or <i>Byy</i> , <i>Cy</i> or <i>Cyy</i> <sup>7</sup>
D	—	—	—	—
Dr	R	R	R	R
—	—	—	L <sup>3, 6</sup>	L
—	—	—	—	M <sup>6</sup>
—	—	—	W	W

<sup>1</sup> Soil Survey Staff, 1951.

<sup>2</sup> Soil Survey Staff, 1962; same content used in *Soil Taxonomy* (Soil Survey Staff, 1975), except for addition of Limnic (L) horizon.<sup>3</sup>

<sup>3</sup> Soil Survey Staff, 1975. Limnic materials and limnic layer were recognized in 1975, formally dropped in 1985 (National Soil Taxonomy Handbook 615.30); master L horizon was formally adopted in 2006.<sup>6</sup>

<sup>4</sup> Guthrie and Witty, 1982.

<sup>5</sup> Soil Survey Staff, 1998.

<sup>6</sup> Soil Survey Staff, 2006.

<sup>7</sup> Soil Survey Staff, 2010a.

<sup>8</sup> Master horizon G (1951) was changed to a horizon suffix (g) that can be used with master horizon A, B, or C; e.g., *Bg*.

**HORIZON SUFFIXES**—(**NOTE:** Gray boxes indicate the year the convention was first adopted.)

<b>Horizon Suffixes (also called "Horizon Subscripts" and "Subordinate Distinctions")</b>				
<b>1951 <sup>1</sup></b>	<b>1962 <sup>2</sup> 1975 <sup>2</sup></b>	<b>1982 <sup>3</sup></b>	<b>1998 <sup>4</sup></b>	<b>2010 <sup>5</sup></b>
—	—	a	a	a
b	b	b	b	b
ca	ca	k	k	k
cn	cn	c	c	c
—	—	—	co <sup>6</sup>	co
cs	cs	y	y	y
—	—	—	d (1988) <sup>7</sup>	d
—	—	—	di <sup>6</sup>	di
—	—	e	e	e
f	f	f	f	f
—	—	—	ff	ff
g	g	g	g	g
h	h	h	h	h
ir	ir	s	s	s
—	—	i	i	i
—	—	—	j	j
—	—	—	jj	jj
ca	ca	k	k	k
—	—	—	—	kk <sup>8</sup> (2006)
m	m <sup>9</sup>	m	m	m
—	—	—	ma <sup>6</sup>	ma
—	—	n	n	n
—	—	o	o	o
p	p	p	p	p
si	si	q	q	q
r <sup>10</sup>	—	r	r	r
ir	ir	s	s	s
sa	sa	n	n	n
—	—	—	—	se <sup>9</sup> (2011)
—	si	q	q	q



<b>Horizon Suffixes (also called "Horizon Subscripts" and "Subordinate Distinctions")</b>				
<b>1951 <sup>1</sup></b>	<b>1962 <sup>2</sup>, 1975 <sup>2</sup></b>	<b>1982 <sup>3</sup></b>	<b>1998 <sup>4</sup></b>	<b>2010 <sup>5</sup></b>
—	—	—	ss (1991)	ss
t	t	t	t	t
u	—	—	—	u <sup>8</sup> (2006)
—	—	v	v	v
—	—	w	w	w
—	x	x	x	x
cs	cs	y	y	y
—	—	—	—	yy <sup>5</sup>
sa	sa	z	z	z

<sup>1</sup> Soil Survey Staff, 1951.

<sup>2</sup> Soil Survey Staff, 1962; same content also used in *Soil Taxonomy* (Soil Survey Staff, 1975).

<sup>3</sup> Guthrie and Witty, 1982.

<sup>4</sup> Soil Survey Staff, 1998.

<sup>5</sup> Soil Survey Staff, 2010a.

<sup>6</sup> Soil Survey Staff, 1999.


<sup>7</sup> Soil Survey Staff, 1988.

<sup>8</sup> Soil Survey Staff, 2006.


<sup>9</sup> The definition is changed to no longer include fragipans (which become "x").

<sup>10</sup> Definition of r (1951; dropped 1962 <sup>2</sup>) is *not* the same as used since 1981. <sup>3</sup>

**(Soil) Textural Triangle:**  
Family Particle-Size Classes (——)



<sup>1</sup> Very fine sand fraction (0.05 - 0.1 mm) is treated as silt for Soil Taxonomy family groupings; coarse fragments are considered the equivalent of coarse sand in the boundary between silty and loamy classes.



**SOIL MOISTURE REGIMES**—Refers to soil moisture or ground water presence in or on soil at tensions  $>0$  and  $\leq 1500$  kPa ( $\approx \leq 15$  bar).

Soil moisture conditions of a pedon (i.e., *Soil Water State*) can be estimated or measured at the time of observation and subsequently assigned to a Water State class (or subclass; see p. 1–15). In a broader context, the prevailing soil moisture condition of a site can be estimated or measured for “normal years” (i.e., most typical or dominant climatic conditions). Class assignment takes into account: 1) the extent of ground water influence (usually via “depth to”) and 2) the seasonal status of water held at tensions  $< 1500$  kPa ( $\approx < 15$  bar) in the moisture control section. <sup>1</sup> In soil taxonomy <sup>1</sup>, soil moisture regimes are assigned as classes (e.g., *Ustic Soil Moisture Regime*) and are used at the higher categories of the system (i.e., from Order down through Subgroup).

<b>Soil Moisture Regime</b>	<b>Criteria <sup>1</sup> (generalized, abbreviated)</b>
aquic	A reducing regime for soils that are free of dissolved oxygen and saturated (seasonal ground water fluctuations typical). Unlike other regimes, the aquic regime may occur temporarily for only a few days.
peraquic	A reducing regime for soils that are free of dissolved oxygen and permanently saturated (ground water is almost always above, at, or very close to the surface).
aridic (torric) <sup>2</sup>	The predominantly dry regime for soils of arid and semiarid climates that are unsuitable for cultivation without irrigation. Soil is dry (in all parts of soil moisture control section) >50% of all days annually when soil is >5 °C at 50 cm and moist in some part for <90 consecutive days when soil is >8 °C at 50 cm in normal years.
udic	The predominantly moist regime for soils of humid climates with well distributed rainfall. Soil is dry (in any part of soil moisture control section) for <90 cumulative days in normal years.
perudic	An extremely wet regime for soils of climates where precipitation exceeds evapotranspiration in all months in normal years. Soil is almost always moist; soil tension is rarely >100 kPa (≈ >1 bar).
ustic	The temporarily dry regime for soils of climates that are intermediate between dry (aridic) and moist (udic). Soil is intermittently moist and dry; moisture limited but usually available when climate is suitable for plant growth. Soil is moist >180 cumulative days or >90 consecutive days.
xeric	The seasonally dry regime for soils of Mediterranean climates with cool, moist winters and warm, dry summers. Soil is moist in all parts for ≥45 consecutive days in the 4 months following the winter solstice and dry in all parts for ≥45 consecutive days in the 4 months following the summer solstice. Soil is also moist in some part >50% of all days when soil is >5 °C at depth of 50 cm or moist in some part for ≥90 consecutive days when soil is >8 °C at depth of 50 cm in normal years.

<sup>1</sup> Complete criteria available in *Keys to Soil Taxonomy*, 11th ed. (Soil Survey Staff, 2010a).

<sup>2</sup> Aridic and torric are terms for the same soil moisture regime, but they are used in different categories in soil taxonomy. Limits set

for soil temperature exclude from this regime soils in very cold and dry polar regions and in areas at high elevations. Such soils are considered to have *anhydrous conditions*.

**SOIL TEMPERATURE REGIMES AND CLASSES** (per *Keys to Soil Taxonomy*<sup>1, 2</sup>)—

Soil Temperature Regimes <sup>1</sup>	Soil Temperature Classes <sup>2</sup>	Criteria: MAST <sup>3</sup> measured at 50 cm or at the upper boundary of a root-limiting layer if shallower
Gelic	(see below)	≤0 °C in Gelic suborders and Gelic great groups <i>or</i> <1 °C in Gelisols (permafrost expected)
	Hypergelic	≤ -10 °C
	Pergelic	-10 to -4 °C
	Subgelic	-4 to 1 °C
Cryic	(no family temperature class)	≥0 to <8 °C, but no permafrost, and
		<ol style="list-style-type: none"> <li>1. In mineral soils: the MSST<sup>4</sup> is:               <ol style="list-style-type: none"> <li>a. If soil is not saturated during summer and                   <ol style="list-style-type: none"> <li>(1) If no O horizon: ≥0 to 15 °C; or</li> <li>(2) If there is an O horizon: ≥0 to 8 °C; or</li> </ol> </li> <li>b. If soil is saturated during summer and                   <ol style="list-style-type: none"> <li>(1) If no O horizon: ≥0 to 13 °C; or</li> <li>(2) If there is an O horizon or a histic epipedon: ≥0 to 6 °C.</li> </ol> </li> </ol> </li> <li>2. In organic soils: ≥0 to 6 °C.</li> </ol>
<i>For soils with a difference between mean summer and mean winter soil temperature of ≥6 °C:</i>		
Frigid	Frigid	≥0 to <8 °C (but warmer than cryic in summer)
Mesic	Mesic	8 to <15 °C
Thermic	Thermic	15 to <22 °C
Hyperthermic	Hyperthermic	≥22 °C
<i>For soils with a difference between mean summer and mean winter soil temperature of &lt;6 °C:</i>		
Isofrigid	Isofrigid	<8 °C
Isomesic	Isomesic	8 to <15 °C
Isothermic	Isothermic	15 to <22 °C
Isohyperthermic	Isohyperthermic	≥22 °C

- <sup>1</sup> Soil temperature regimes are used as criteria in the suborder, great group, and subgroup categories of soil taxonomy (Soil Survey Staff, 2010a).
- <sup>2</sup> Soil temperature classes are used as differentiae in the family category of soil taxonomy, excluding cryic soils (Soil Survey Staff, 2010a).
- <sup>3</sup> MAST=Mean annual soil temperature (Soil Survey Staff, 1999).
- <sup>4</sup> MSST=Mean summer soil temperature (see Soil Survey Staff, 1999, p. 108).

## **REFERENCES**

---

- Guthrie, R.L., and J.E. Witty. 1982. New designations for soil horizons and layers and the new Soil Survey Manual. Soil Sci. Soc. Am. J. 46:443–444.
- Natural Resources Conservation Service, Soil Classification Staff. 1999. Personal communication. USDA, NRCS, National Soil Survey Center, Lincoln, NE.
- Soil Survey Division Staff. 1993. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 1951. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 1962. Identification and nomenclature of soil horizons. Supplement to Agric. Handb.18, Soil Survey Manual (replacing pages 173–188). USDA, SCS. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 1975. Soil taxonomy. USDA, SCS, Agric. Handb. 436. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 1988. Keys to soil taxonomy, 4th ed. SMSS Tech. Monogr. 6, Cornell Univ., Ithaca, NY.
- Soil Survey Staff. 1998. Keys to soil taxonomy, 8th ed. USDA, SCS. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 1999. Soil taxonomy, 2nd ed. USDA, NRCS, Agric. Handb. 436. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 2006. Keys to soil taxonomy, 10th ed. USDA, NRCS. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 2010a. Keys to soil taxonomy, 11th ed. USDA, NRCS. U.S. Gov. Print. Office, Washington, DC.
- Soil Survey Staff. 2010b. National soil survey handbook (NSSH) [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE (<http://soils.usda.gov/technical/handbook/>).

# GEOLOGY

*P.J. Schoeneberger, D.A. Wysocki, and E.C. Benham, NRCS, Lincoln, NE*

## INTRODUCTION

The purpose of this section is to expand and augment the geologic information found or needed in the "Site Description" and "Profile/Pedon Description" sections.

### BEDROCK - KIND—

This table is repeated here from the "Site Selection" section for convenience in using the following rock charts.

Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
<b>IGNEOUS—INTRUSIVE</b>			
anorthosite	ANO	peridotite	PER
diabase	DIA	pyroxenite	PYX
diorite	DIO	quartz-diorite	QZD
gabbro	GAB	quartz-monzonite	QZM
granite	GRA	syenite	SYE
granitoid <sup>2</sup>	GRT	syenodiorite	SYD
granodiorite	GRD	tonalite	TON
monzonite	MON	ultramafic rock <sup>2</sup>	UMU
<b>IGNEOUS—EXTRUSIVE</b>			
a'a lava	AAL	pahoehoe lava	PAH
andesite	AND	pillow lava	PIL
basalt	BAS	pumice (flow, coherent)	PUM
block lava	BLL	rhyolite	RHY
dacite	DAC	scoria (coherent mass)	SCO
latite	LAT	tachylite	TAC
obsidian	OBS	trachyte	TRA

Kind <sup>1</sup>	Code	Kind <sup>1</sup>	Code
<b>IGNEOUS—PYROCLASTIC</b>			
ignimbrite	IGN	tuff, welded	TFW
pyroclastics (consolidated)	PYR	tuff breccia	TBR
pyroclastic flow	PYF	volcanic breccia	VBR
pyroclastic surge	PYS	volcanic breccia, acidic	AVB
tuff	TUF	volcanic breccia, basic	BVB
tuff, acidic	ATU	volcanic sandstone	VST
tuff, basic	BTU		
<b>METAMORPHIC</b>			
amphibolite	AMP	metavolcanics	MVO
gneiss	GNE	mica schist	MSH
gneiss, biotite	BTG	migmatite	MIG
gneiss, granodiorite	GDG	mylonite	MYL
gneiss, hornblende	HBG	phyllite	PHY
gneiss, migmatitic	MMG	schist	SCH
gneiss, muscovite-biotite	MGB	schist, biotite	BTS
granofels	GRF	schist, graphitic	GRS
granulite	GRL	schist, muscovite	MVS
greenstone	GRE	schist, sericite	SCS
hornfels	HOR	serpentinite	SER
marble	MAR	siltite	SIT
meta-conglomerate	MCN	slate	SLA
metaquartzite	MQT	slate, sulfidic	SFS
metasedimentary rocks <sup>2</sup>	MSR	soapstone (talc)	SPS
metasiltstone	MSI		
<b>SEDIMENTARY—CLASTICS</b>			
arenite	ARE	mudstone	MUD
argillite	ARG	ortho-quartzite	OQT
arkose	ARK	porcellanite	POR
breccia, nonvolcanic (angular fragments)	NBR	sandstone	SST



<b>Kind <sup>1</sup></b>	<b>Code</b>	<b>Kind <sup>1</sup></b>	<b>Code</b>
breccia, nonvolcanic, acidic	ANB	sandstone, calcareous	CSS
breccia, nonvolcanic, basic	BNB	shale	SHA
claystone	CST	shale, acid	ASH
conglomerate (rounded fragments)	CON	shale, calcareous	CSH
conglomerate, calcareous	CCN	shale, clayey	YSH
fanglomerate	FCN	siltstone	SIS
glaucinitic sandstone	GLS	siltstone, calcareous	CSI
graywacke	GRY		
<b>EVAPORITES, ORGANICS, AND PRECIPITATES</b>			
bauxite	BAU	limestone, argillaceous	RLS
chalk	CHA	limestone, cherty	CLS
chert	CHE	limestone, coral	COR
coal	COA	limestone, phosphatic	PLS
diatomite	DIA	limonite	LIM
dolomite (dolostone)	DOL	novaculite	NOV
gypsum	GYP	travertine	TRV
limestone	LST	tripoli	TRP
limestone, arenaceous	ALS	tufa	TUA
<b>INTERBEDDED</b> (alternating layers of different sedimentary lithologies)			
limestone-sandstone-shale	LSS	sandstone-shale	SSH
limestone-sandstone	LSA	sandstone-siltstone	SSI
limestone-shale	LSH	shale-siltstone	SHS
limestone-siltstone	LSI		

<sup>1</sup> Definitions for kinds of bedrock are found in the "Glossary of Landform and Geologic Terms," NSSH, Part 629 (Soil Survey Staff, 2012), or in the *Glossary of Geology* (Neuendorf et al., 2005).

<sup>2</sup> Generic term; use only with regional or reconnaissance surveys (Order 3, 4, 5; see Guide to Map Scales and Minimum-Size Delineations, p. 7–21).

## **ROCK CHARTS**

---

The following rock charts (**Igneous**, **Metamorphic**, and **Sedimentary** and **Volcaniclastic**) summarize grain size, composition, or genetic differences between related rock types. **NOTE:** 1) Most, but not all, of the rocks in these tables are found in the NASIS choice lists. Those not in NASIS are uncommon in the pedosphere but are included in the charts for completeness and to aid in the use of geologic literature. 2) Most, but not all, of the rocks presented in these tables can be definitively identified in the field; some may require additional laboratory analyses (e.g., grain counts, thin section analyses).

## IGNEOUS ROCKS CHART

KEY MINERAL COMPOSITION								
CRYSTALLINE TEXTURE	Acidic (Felsic)		INTERMEDIATE				Basic (mafic)	Ultrabasic (ultramafic)
	Potassium (K) Feldspar >2/3 of Total Feldspar Content	No Quartz	Potassium (K) Feldspar and Plagioclase (Na, Ca) Feldspar in about equal proportions	Plagioclase (Na, Ca) Feldspar >2/3 of Total Feldspar Content	Sodic (Na) Plagioclase		Calcic (Ca) Plagioclase	Pyroxene and Olivine
	Quartz	No Quartz	Quartz	No Quartz	Quartz	No Quartz	Quartz	
<b>PEGMATITIC</b> <sup>1</sup>	granite pegmatite	syenite pegmatite	← monzonite pegmatite →	diorite pegmatite	gabbro pegmatite			peridotite (mostly olivine)
<b>PHANERITIC</b> <sup>2</sup>	granite	syenite	quartz monzonite	monzonite	diorite		gabbro	pyroxenite (mostly pyroxene)
<b>PORPHYRITIC</b> <sup>3</sup>	granite porphyry	syenite porphyry	quartz-monzonite porphyry	monzonite porphyry	diorite porphyry		diabase	
<b>APHANITIC</b> <sup>4</sup> <b>micro<sup>5</sup> crypto<sup>6</sup></b>	rhyolite porphyry	trachyte porphyry	quartz-latite porphyry	latite porphyry	dacite porphyry		basalt porphyry	} lava <sup>7</sup>
<b>GLASSY</b> <sup>8</sup>	Obsidian (and its varieties: perlite, pitchstone, pumice, scoria) Pyroclastics are shown on the Sedimentary and Volcaniclastic Rocks chart.							

<sup>1</sup> Pegmatitic: Very coarse, uneven-sized crystal grains; 5 to > 20 mm  
<sup>2</sup> Phaneritic: Crystals discernable by eye or 10X lens; 1-5 mm  
<sup>3</sup> Porphyritic: Larger crystals embedded within a fine-grained matrix  
<sup>4</sup> Aphanitic: Crystals not visible by eye or 10X lens; <1 mm  
<sup>5</sup> Microcrystalline crystals resolvable by optical microscope  
<sup>6</sup> Cryptocrystalline crystals resolvable by electron microscope  
<sup>7</sup> Lava: Generic name for extrusive flows of non-clastic, aphanitic rocks (rhyolite, andesite, basalt)  
<sup>8</sup> Glassy: Noncrystalline or weakly crystalline

## METAMORPHIC ROCKS CHART

[Not all rock types listed here can be definitively identified in the field (e.g., may require grain counts). Not all rock types shown here are available on Bedrock - Kind choice list. They are included here for completeness and as aids to using geologic literature.]

<b>NONFOLIATED STRUCTURE</b>		<b>CRUDE ALIGNMENT</b>		<b>FOLIATED STRUCTURE (e.g., banded)</b>		
<b>CONTACT METAMORPHISM</b>		<b>FAULT ZONE METAMORPHISM</b>		<b>REGIONAL METAMORPHISM</b>		<b>PLUTONIC METAMORPHISM</b>
Low Grade	Medium Grade	High Grade	Low Grade	Low Grade	Medium Grade	High Grade
granofels hornfels marble metaquartzite serpentine soapstone (talc)	crush breccia mylonite	slate phyllite greenstone schist amphibolite gneiss granulite migmatite*	<-----Metaconglomerate-----> <-----Metavolcanics-----> <-----Metasedimentary----->			
* Partial melting occurs.						

(Schoeneberger and Wysocki, 1998)

**SEDIMENTARY AND VOLCANICLASTIC ROCKS**

CLASTIC			NONCLASTIC		
Dominant Grain Size			Chemical	Biochemical	Organic
Very Fine	Fine	Medium	Evaporates, Precipitates	Accretionates	Reduzates
<----- (Argillaceous) -----> < 0.002 mm	0.002 - 0.06 mm	(Arenaceous) 0.06 - 2.0 mm	<b>CARBONATE ROCKS</b> Limestones (ls) (> 50% calcite)		
<----- argillite -----> (more indurated, less laminated and fissile)	<----- shale -----> (laminated, fissile)	Sandstones (ss): arenite arkose (mainly feldspar) glauconitic ss ("greensand") graywacke (dark, "dirty" ss) orthoquartzite (mainly quartz)	anhydrite (CaSO <sub>4</sub> )  gypsum (CaSO <sub>4</sub> • 2H <sub>2</sub> O)  halite (NaCl)	<b>CARBONATE ROCKS</b> Limestones (ls) accretionary types biostromal ls organic reef pelagic ls (chalk)  bio-clastic types coquina oolithic ls lithographic ls	
claystone (non-laminated, nonfissile)	siltstone (non-laminated, nonfissile)				black shale (organics and fine sediments)  bituminous ls bog iron ores  coal
<b>VOLCANICLASTICS (includes Pyroclastics)</b>			<b>OTHER NONCLASTIC ROCKS</b>		
<----- ignimbrite -----> (mainly pumice frags; consolidated pyroclastic flows)	<----- tuff -----> (consolidate volcanic ash, tephra)	agglomerate (rounded frags) volcanic breccia (angular frags)	dolomite (>50% CaMg(CO <sub>3</sub> )) phosphatic limestone		
<----- pumice -----> (specific gravity <1.0; highly vesicular)	<----- scoria -----> (specific gravity >2.0; slightly or moderately vesicular)				Siliceous rocks (Silica dominated): chert (jasper, chalcidony, opal); diatomite Rock phosphate Iron-bearing rocks (Fe-SiO <sub>2</sub> dominated): jaspilite, specular hematite, magnetite

(Schoeneberger and Wysocki, 2000)

**MASS MOVEMENT (MASS WASTING) TYPES FOR SOIL SURVEY**  
(landforms, processes, and sediments)

		<b>LANDSLIDE</b>				
<b>Movement Types:</b>	<b>FALL</b> Free fall, bouncing, or rolling	<b>TOPPLE</b> Forward rotation over a point	<b>SLIDE*</b> Net lateral displacement along a slip face <b>Rotational Slide</b> Lateral displacement along a concave slip face with backward rotation <b>Translational Slide</b> Lateral displacement along a planar slip face; no rotation <----- Compound Slide -----> Intermediate between rotational and translational; e.g., a compound rock slide	<b>SPREAD</b> A wet layer squeezes up and out and drags along intact blocks or beds; e.g., extrusion, liquefaction  (=Lateral Spread)	<b>FLOW</b> The entire mass, wet or dry, moves as a viscous liquid	<b>COMPLEX LANDSLIDE</b> Combination of multiple types of movement
<b>Dominant Material</b>	<b>Consolidated (Bedrock)</b> Bedrock masses dominant	rock topple	rotational rock slide  (e.g., Toreva block)	rock spread  block spread	rock fragment flow (e.g., rockfall avalanche =sturzstrom)	No unique subtypes are recognized here; many possible
	<b>Unconsolidated: Coarser</b> Coarse fragments dominant	debris topple	rotational debris slide  <----- debris slide ----->	debris spread	debris avalanche (drier, steep slope) debris flow (wetter) (e.g., lahar)	Option: name the main movement types (e.g., a Complex Rock Spread-Debris Flow Landslide)
	<b>Finer</b> Fine earth particles dominant	earth topple (=soil topple)	rotational earth slide	earth spread (e.g., sand boil)	earth flow (e.g., creep, loess flow, mudflow, sandflow, solifluction)	

\* Slides, especially rotational slides, are commonly and imprecisely called "slumps."

(Schoeneberger and Wysocki, 2000; developed from Cruden and Varnes, 1996)

# NORTH AMERICAN GEOLOGIC TIME SCALE <sup>1, 2</sup>

ERA	Geologic Period	Geologic Epoch	Sub-division	Oxygen Isotope Stage	Years (BP)	
<b>CENOZOIC</b>	<b>QUATERNARY</b>	<b>Holocene</b>		(1)	0 to 10-12 ka*	
		<i>Late Pleistocene</i>		Late Wisconsin	(2)	10-12 to 28 ka
				Middle Wisconsin	(3, 4)	28 to 71 ka
				Early Wisconsin Late Sangamon	(5a - 5d)	71 to 115 ka
				Sangamon	(5e)	115 to 128 ka
			<b>Pleistocene</b>		Late Middle Pleistocene (Illinoian)	(6 - 8)
		<i>Middle Pleistocene</i>		Middle Pleistocene	(9 - 15)	300 to 620 ka
				Early Middle Pleistocene	(16 - 19)	620 to 770 ka
			<i>Early Pleistocene</i>			770 ka to 2.6 Ma**
		<b>TERTIARY</b>	<b>Neo- gene</b>	<b>Pliocene</b>		
	<b>Miocene</b>			5.3 to 23.0 Ma		
	<b>Paleo- gene</b>		<b>Oligocene</b>			23.0 to 33.9 Ma
			<b>Eocene</b>			33.9 to 55.8 Ma
			<b>Paleocene</b>			55.8 to 65.5 Ma
	<b>MESOZOIC</b>	<b>CRETACEOUS</b>	<i>Late Cretaceous</i>			65.5 to 99.6 Ma
<i>Early Cretaceous</i>			99.6 to 145.5 Ma			
<b>JURASSIC</b>					145.5 to 201.6 Ma	
<b>TRIASSIC</b>				201.6 to 251.0 Ma		
<b>PALEOZOIC</b>	<b>PERMIAN</b>				251.0 to 299.0 Ma	
	<b>PENNSYLVANIAN</b>				299.0 to 318.0 Ma	
	<b>MISSISSIPPIAN</b>				318.0 to 359.0 Ma	
	<b>DEVONIAN</b>				359.0 to 416.0 Ma	
	<b>SILURIAN</b>				416.0 to 444.0 Ma	
	<b>ORDOVICIAN</b>				444.0 to 488.0 Ma	
	<b>CAMBRIAN</b>				488.0 to ≈ 542.0 Ma	
<b>PRECAMBRIAN ERA</b>					> 542.0 Ma	

\*ka = x 1,000    \*\* Ma = x 1,000,000 ( $\approx$  = approximately)

<sup>1</sup> Modified from Morrison, 1991; Sibrava et al., 1986; and Harland et al., 1990.

<sup>2</sup> Modified from Walker and Geissman, 2009.

## TILL TERMS

Genetic classification and relationships of till terms commonly used in soil survey (Schoeneberger and Wysocki, 2000; adapted from Goldthwaite and Matsch, 1988).

Location (Facies of tills grouped by position at time of deposition)	Till Types	
	Terrestrial	Waterlaid
<b>Proglacial Till</b> (at the front of or in front of glacier)	proglacial <b>flow till</b>	waterlaid <b>flow till</b>
<b>Supraglacial Till</b> (on top of or within upper part of glacier)	supraglacial <b>flow till</b> <sup>1, 3</sup> supraglacial <b>melt-out till</b> <sup>1</sup> (ablation till—NP) <sup>1</sup> (lowered till—NP) <sup>2</sup> (sublimation till—NP) <sup>2</sup>	—
<b>Subglacial Till</b> (within the lower part of or beneath glacier)	<b>lodgment till</b> <sup>1</sup> subglacial <b>melt-out till</b> subglacial <b>flow till</b> (= "squeeze till" <sup>2, 3</sup> )  (basal till—NP) <sup>1</sup> (deformation till—NP) <sup>2</sup> (gravity flow till—NP) <sup>2</sup>	waterlaid <b>melt-out till</b> waterlaid <b>flow till</b> iceberg till (= "ice-rafted")

<sup>1</sup> *Ablation till* and *basal till* are generic terms that only describe "relative position" of deposition and have been widely replaced by more specific terms that convey both relative position and process. *Ablation till* (any comparatively permeable debris deposited within or above stagnant ice) is replaced by *supraglacial melt-out till*, *supraglacial flow till*, etc. *Basal till* (any dense, nonsorted subglacial till) is replaced by *lodgment till*, *subglacial melt-out till*, *subglacial flow till*, etc.

<sup>2</sup> Additional (proposed) till terms that are outdated or have not gained wide acceptance and are considered to be *Not Preferred* and should *not* be used.

<sup>3</sup> Also called *gravity flow till* (not preferred).



## PYROCLASTIC TERMS

(Schoeneberger and Wysocki, 2002)

Pyroclasts and Pyroclastic Deposits (Unconsolidated)			
Size Scale:			
0.062 mm <sup>1</sup>	2 mm	64 mm <sup>1</sup>	
<----- <b>tephra</b> ----->			
(all ejecta)			
<----- <b>ash</b> <sup>1</sup> ----->		<--- <b>cinders</b> <sup>2</sup> --->	<--- <b>bombs</b> <sup>1</sup> --->
		(specific gravity >1.0 and <2.0)	(fluid-shaped coarse fragments)
<-----> <b>fine ash</b> <sup>1</sup>	<-----> <b>coarse ash</b> <sup>1</sup>		
		<--- <b>lapilli</b> <sup>1</sup> --->	<--- <b>blocks</b> <sup>1</sup> --->
		(specific gravity >2.0)	(angular-shaped coarse fragments)
<----- <b>scoria</b> <sup>2</sup> ----->			
(slightly or moderately vesicular; specific gravity >2.0)			
	<-----> <b>pumiceous ash</b> <sup>3</sup>	<----- <b>pumice</b> ----->	
		(highly vesicular; specific gravity <1.0)	
Associated Lithified (Consolidated) Rock Types			
<-----> <b>fine tuff</b> <sup>1</sup>	<-----> <b>coarse tuff</b> <sup>1</sup>	<- <b>lapillistone</b> <sup>1</sup> ->	<-----> <b>pyroclastic breccia</b>
		(sp. gravity >2.0)	
<----- <b>welded tuff</b> <sup>1</sup> ----->		<----- <b>agglomerate</b> <sup>1</sup> ----->	
		(rounded, volcanic coarse fragments)	
<----- <b>ignimbrite</b> ----->		<----- <b>volcanic breccia</b> <sup>1</sup> ----->	
(consolidated ash flows and nuee ardentes)		(angular, volcanic coarse fragments)	

<sup>1</sup> These size breaks are taken from geologic literature (Fisher, 2005) and based on the modified Wentworth scale. The 0.062-mm break is very close to the USDA's 0.05-mm break between *coarse silt* and *very fine sand* (Soil Survey Division Staff, 1993). The 64-mm break is relatively close to the USDA's 76-mm break between *coarse gravel* and *cobbles*. (See "Comparison of Particle Size Classes in Different Systems" in the "Profile/Pedon Description" on p. 2-45.)

<sup>2</sup> A lower size limit of 2 mm is required in soil taxonomy (Soil Survey Staff, 1994; p. 54) but is *not* required in geologic usage (Fisher, 2005).

<sup>3</sup> The descriptor for pumice particles <2 mm, as used in soil taxonomy (Soil Survey Staff, 1999). Geologic usage does *not* recognize any size restrictions for pumice.

## **HIERARCHICAL RANK OF LITHOSTRATIGRAPHIC UNITS** *1, 2, 3*

---

**Supergroup**—The broadest lithostratigraphic unit. A supergroup is an assemblage of related, superposed groups, or groups and formations. Supergroups are most useful in regional or broad scale synthesis.

**Group**—The lithostratigraphic unit next in rank below a supergroup. A group is a named assemblage of related superposed formations, which may include unnamed formations. Groups are useful for small-scale (broad) mapping and regional stratigraphic analysis.

**Formation** (called **Geologic Formation** in NASIS)—The basic lithostratigraphic unit used to describe, delimit, and interpret sedimentary, extrusive igneous, metavolcanic, and metasedimentary rock bodies (excludes metamorphic and intrusive igneous rocks) based on lithic characteristics and stratigraphic position. A formation is commonly, but not necessarily, tabular and stratified and is of sufficient extent to be mappable at the earth's surface or traceable in the subsurface at conventional map scales.

(Formations can be, but are not necessarily, combined to form higher rank units [groups and supergroups] or subdivided into lower rank units [members or beds].)

**Member**—The formal lithostratigraphic unit next in rank below a formation and always part of a formation. A formation need not be divided selectively or entirely into members. A member may extend laterally from one formation to another.

Specifically defined types of members:

**Lens** (or **Lentil**): A geographically restricted member that terminates on all sides within a formation.

**Tongue**: A wedge-shaped member that extends beyond the main formation boundary or that wedges or pinches out within another formation.

**Bed**—The smallest formal lithostratigraphic unit of sedimentary rock. A bed is a subdivision of a member based upon distinctive characteristics and/or economic value (e.g., coal bed). Members

need not be divided selectively or entirely into beds.

**Flow**—The smallest formal lithostratigraphic unit of volcanic rock. A flow is a discrete, extrusive, volcanic body distinguishable by texture, composition, superposition, and other criteria.

<sup>1</sup> Lithostratigraphic units are mappable rock or sediment bodies that conform to the Law of Superposition (Article 2, Section A).

<sup>2</sup> Separate data element (text field) in NASIS.

<sup>3</sup> Adapted from *North American Stratigraphic Code* (North American Commission on Stratigraphic Nomenclature, 1983).

## **REFERENCES**

---

Cruden, D.M., and D.J. Varnes. 1996. Landslide types and processes. In A.K. Turner and R.L. Schuster (ed.) *Landslides: Investigation and mitigation*. Spec. Rep. 247, Transportation Research Board, National Research Council. National Academy Press, Washington, DC.

Fisher, R.V. (updated by T. Frost) 2005. Pyroclastic sediments and rocks. AGI data sheet 6.3. In J.D. Walker and H.A. Cohen (ed). *The geoscience handbook*. AGI data sheets, 4th ed. Am. Geol. Inst., Alexandria, VA.

Goldthwaite, R.P., and C.L. Matsch (ed.) 1988. Genetic classification of glacial deposits: Final report of the commission on genesis and lithology of glacial Quaternary deposits of the International Union for Quaternary Research (INQUA). A.A. Balkema, Rotterdam.

Harland, W.B., R.L. Armstrong, L.E. Craig, A.G. Smith, and D.G. Smith. 1990. A geologic time scale. Press Syndicate of University of Cambridge, Cambridge, UK. 1 sheet.

Jackson, J.A. (ed.) 1997. *Glossary of geology*, 4th ed. Am. Geol. Inst., Alexandria, VA.

Morrison, R.B. (ed.) 1991. Quaternary nonglacial geology: Conterminous United States. *Geol. Soc. Am., Decade of North American Geology, Geology of North America*, vol. K-2.

Neuendorf, K., J.P. Mehl, and J.A. Jackson. 2005. *Glossary of geology*, 5th ed. Am. Geol. Inst., Alexandria, VA.

North American Commission on Stratigraphic Nomenclature. 1983. North American stratigraphic code. *Am. Assoc. Petrol. Geol. Bull.* 67:841–875.

Schoeneberger, P.J., and D.A. Wysocki. 1998. Personal communication. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Schoeneberger, P.J., and D.A. Wysocki. 2000. Personal communication. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Schoeneberger, P.J., and D.A. Wysocki. 2002. Personal communication. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Sibrava, V., D.Q. Bowen, and D.Q. Richmond (ed.) 1986. Quaternary glaciations in the Northern Hemisphere: Final report of the International Geological Correlation Programme, Project 24. Quaternary Sci. Rev., Vol. 5. Pergamon Press, Oxford.

Soil Survey Division Staff. 1993. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1994. Keys to soil taxonomy, 6th ed. USDA, SCS. Pocohantas Press, Inc., Blacksburg, VA.

Soil Survey Staff. 1999. Soil taxonomy, 2nd ed. USDA, NRCS, Agric. Handb. 436. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 2011. Soil survey laboratory information manual. Soil Surv. Invest. Rep. 45, ver. 2.0. USDA, NRCS, National Soil Survey Center, Lincoln, NE ([ftp://ftp-fc.sc.egov.usda.gov/NSSC/Lab\\_Info\\_Manual/SSIR\\_45.pdf](ftp://ftp-fc.sc.egov.usda.gov/NSSC/Lab_Info_Manual/SSIR_45.pdf))

Soil Survey Staff. 2012. Glossary of landform and geologic terms. Part 629, National soil survey handbook (NSSH) [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Tennissen, A.C. 1974. Nature of earth materials. Prentice-Hall, Inc., Englewood Cliffs, NJ.

Walker, J.D., and J.W. Geissman. 2009. Geologic time scale. Geol. Soc. Am.

# LOCATION

*D.A. Wysocki, P.J. Schoeneberger, and E.C. Benham, NRCS, Lincoln, NE*

## **GPS LOCATION**

---

**GEODETTIC DATUM (Horizontal\_datum\_name** in NASIS)—A geodetic datum must accompany latitude and longitude. A geodetic datum is a model that defines the earth's shape and size and serves as a latitude, longitude reference. Geodetic datum is a selectable GPS parameter. The preferred datum is the World Geodetic System 1984 (WGS-84).

<b>Datum Name</b>	<b>Code</b>
American Samoa 1962	
Astro Beacon "E" 1945	
Astro Tern Island (FRIG)	
Astronomical Station 1952	
Bellevue (IGN)	
Canton Astro 1966	
Chatham Island Astro 1971	
DOS 1968	
Easter Island 1967	
Geodetic Datum 1949	
Guam 1963	
Gux 1 Astro	
Johnston Island 1961	
Kusaie Astro 1961	
Luzon	
Midway Astro 1961	
North American Datum of 1927	NAD27
North American Datum of 1983 <sup>1</sup>	NAD83
Old Hawaiian	
Pitcairn Astro 1967	
Santo (DOS) 1965	
Viti Levu 1916	
Wake Island Astro 1952	
Wake-Eniwetok 1960	
<b>World Geodetic System 1984 <sup>1</sup></b>	<b>WGS84</b>

<sup>1</sup> Preferred datum method for continental U.S.

## **PUBLIC LAND SURVEY**

---

The Public Land Survey System (PLSS) is a rectangular method for describing and subdividing land in the U.S. The PLSS process first establishes two controlling survey lines for a large tract: an east-west base line and a north-south principal meridian, which intersect at an initial point. Thompson (1987; p. 82–83) shows base lines and principal meridians for the conterminous U.S. Lines parallel (standard parallels) to the base line are established at 24- or 30-mile intervals. The meridian, baseline, and standard parallels form a lattice for further subdivision. Subsequent survey divides land into townships of 36 square miles (6 miles on a side). Each township is subdivided into 36 sections 1 mile square (640 acres). Each section is further subdivided into quarter-sections (160 acres).


<b>PLSS Principal Meridians</b>	
Black Hills	New Mexico Principal
Boise	Ohio Company Purchase
Chickasaw	Ohio River
Choctaw	Principal
Cimarron	Salt Lake
Connecticut Western Reserve	San Bernardino
Copper River	Second Principal
Fairbanks	Second Scioto River
Fifth Principal	Seward
First Principal	Sixth Principal
First Scioto River	St. Helena
Fourth Principal	St. Stephens
Fourth Principal Extended	Tallahassee
Gila and Salt River	Third Principal
Great Miami River	Third Scioto River
Humboldt	Twelve-Mile Square
Huntsville	U.S. Military
Indian	Uintah
Kateel River	Umiat
Louisiana	Ute
Michigan	Washington
Mount Diablo	West of the Great Miami
Muskingum River	Willamette
Navajo	Wind River

Prior to the GPS, soil descriptions predominantly used the PLSS for location. Land survey in certain States predates the PLSS and commonly employs the State Plane Coordinate System for location description. These States include Connecticut, Delaware, Georgia,

Kentucky, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, North Carolina, Ohio (parts), Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, and West Virginia (see State Plane Coordinate System, p. 6–7).


In soil survey, the base line and principal meridian are generally not recorded. Instead, the name of the appropriate USGS topographic 7.5-minute or 15-minute quadrangle is recorded; e.g., Pleasant Dale, NE, 7.5 min. Quad.

**TOWNSHIPS and RANGES**—Each township is identified using two indexes: 1) **Township or Tier** (north-south number relative to the base line), and 2) **Range** (east-west number relative to the Principal Meridian). For example, a township is described as T2N, R4E for second township row north of the base line and fourth range row east of the prime meridian.



**SECTIONS**—Each 1-square-mile **section** is numbered sequentially starting with 1 in the northeast corner of a township proceeding in east-west rows, wrapping back and forth to fill in the township; e.g., *Section 34, T1S, R2E* (Section 34 of Township 1 South, Range 2 East).

**NOTE:** Due to the earth's curvature, survey error, or joins to other survey systems (e.g., Metes and Bounds), occasional irregularities occur in grid areas. Survey adjustments can result in nonstandard size sections and/or breaks in the usual section number sequence.



**SECTION SUBDIVISIONS**—The PLSS subdivides sections into half- and quarter-sections. The section area fraction (1/2, 1/4) is combined with the compass quadrant that the area occupies in a section; e.g., *SW 1/4, Section 34, T1S, R2E* (southwest quarter of section 34, township 1 south, range 2 east). Additional subdivisions, by halves and quarters, describe progressively smaller areas. The land description is presented consecutively beginning with the smallest subdivision; e.g., a 20-acre parcel described as *N 1/2, NW 1/4, SW 1/4, NW 1/4 of Section 34, T1S, R2E* (north half of the northwest quarter of the southwest quarter of the northwest quarter of section 34, township 1 south, range 2 east).

**NOTE: Point locations** (e.g., soil pits) using the PLSS were traditionally measured in English units with reference to a specified



section corner or quarter-corner (1/4 post); e.g., *660 feet east and 1320 feet north of the southwest corner post, Section 34, T1S, R2E.*

## **UNIVERSAL TRANSVERSE MERCATOR (UTM) RECTANGULAR COORDINATE SYSTEM**

---

The Universal Transverse Mercator coordinate system (UTM) is an international reference (military and civilian) that depicts the earth's three-dimensional surface in a relatively accurate, two-dimensional, flat plane and uses Cartesian coordinates (meters) for location. The U.S. Army began use of the UTM projection and grid system in 1947. GPS units can display UTM coordinates, which are simpler for map distance plotting and measurement than latitude and longitude.

The UTM grid spans from 80°S through 84°N latitude (the Universal Polar Stereographic [UPS] system covers polar areas). The UTM system divides the earth into 60 equally spaced, vertically arranged planes known as zones, or world zones; each zone spans 6 degrees of longitude. The zones are sequentially numbered 1 through 60 west to east. Zone numbering begins at 180 degrees longitude, the International Date Line. UTM zone 1 encompasses 180–174 degrees W longitude, zone 2 spans 174–168 W longitude, and so forth through zone 60.


The UTM grid system also divides the earth into 20 equally spaced east-west rows. Each row circles the globe and spans exactly 8 latitude degrees, except for the 12-degree-wide row between 72 and 84 degrees north latitude. The 20 UTM rows are lettered C through X from south to north beginning at 80 degrees south latitude; I and O are omitted to avoid confusion with numbers. Row C spans 80–72 degrees south latitude, row D encompasses 72–64 degrees south latitude, and so forth. The southern hemisphere contains rows C, D, E, F, G, H, J, K, L, and M, whereas the northern hemisphere contains rows N, P, Q, R, S, T, U, V, W, and X.

The central meridian of each zone is the east-west control for UTM coordinates, and other N-S grid lines are parallel to the central meridian. UTM coordinates are expressed as a distance in meters east of a zone's central meridian. This value is called an "**easting**." The central meridian by convention is given a value of 500,000 m east; this eliminates negative distance values. A location west of the central meridian has a value <500,000 m. Easting values can range from 166,000 to 834,000 m. Some protocols give the easting value a leading zero (e.g., 0166000).

The initial north-south grid line for the northern hemisphere is the Equator, which has a value of 0 m. A UTM value called a "**northing**" is expressed as distance in meters north of the Equator. For the northern hemisphere, northings range from 0 to 9,328,000 m (84 N Lat). In the southern hemisphere, the 0 m reference is the South

Pole; the northing is expressed as distance in meters north of it. The range in northings is 1,118,000 (80 S Lat) to 10,000,000 m (Equator). Points on the Equator can be described by either the north or south reference.

A complete UTM location gives in order the zone number, row letter, easting value, and northing value; for example, *16 T, 0313702 m E, 4922401 m N*. The row letters designate the hemisphere location (northern or southern).



All quadrangle maps prepared by the USGS show the UTM coordinates (Snyder, 1987). On 7.5-minute quadrangle maps (1:24,000 and 1:25,000 scale) and 15-minute quadrangle maps (1:50,000, 1:62,500, and standard edition 1:63,360 scales), the UTM grid lines are indicated at 1,000-meter intervals, either by blue ticks in the map margin or with full grid lines. The maps display shortened 1,000-meter values at the tick or grid lines. The full meter values are shown only at ticks nearest the southeast and northwest map corners.

To obtain a UTM grid location from a USGS map, use the grid lines, draw lines connecting corresponding ticks on opposite map edges, or place a transparent grid overlay on the map. Measure distance between any map point and the nearest grid line in cm. If the map scale is 1:24000, multiply the measured cm distance by 240 to

obtain meters on the ground. The northing of a point is the value of the nearest grid line south plus its distance north of that line; the easting is the value of the nearest grid line west of it plus its distance east of that line. On maps at 1:100,000 and 1:250,000 scale, a full UTM grid is shown at intervals of 10,000 meters and is numbered and used in the same way. Various overlay UTM templates that facilitate distance and coordinate measurement from topographic maps are commercially available.

## ***STATE PLANE COORDINATE SYSTEM***

---

The State Plane Coordinate System (SPCS) is designed for mapping and surveying in the U.S. It was developed in the 1930s by the U.S. Coast and Geodetic Survey. Historically, soil description locations sometimes used the SPCS system where the PLSS is nonexistent. The States that have used this system are Connecticut, Delaware, Georgia, Kentucky, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, North Carolina, Ohio (parts), Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, and West Virginia.

The SPCS divides all 50 states of the United States, Puerto Rico, and the U.S. Virgin Islands into 120 numbered zones. The zones correspond to political boundaries (State and most counties). The SPCS establishes a separate coordinate system and two Principal lines in each State: a north-south line and an east-west line. USGS 7.5-minute topographic maps indicate SPCS grids by tick marks along the neatlines (outer border). **NOTE:** Older topo maps based on NAD27 have grid units in feet. After adoption of NAD83, meters become the grid unit.

Specific location coordinates are described by distance and primary compass direction (north [northing], south [southing], east [easting], or west [westing]) relative to the Principal lines; e.g., *10,240 m easting and 1,234 m northing*.

Contact the NRCS State office or the Regional Soil Survey MLRA Office for State-specific details.

## ***REFERENCES***

---

Mozola, A.J. 1989. U.S. public land survey. *In* J.T. Dutro et al., AGI data sheets, 3rd ed. Am. Geol. Inst., United Book Press, Inc.

Snyder, J.P. 1987. Map projections—A working manual. U.S. Geol. Surv. Prof. Pap. 1395. U.S. Gov. Print. Office, Washington, DC.

Thompson, M.M. 1987. Maps for America, 3rd ed. U.S. Geol. Surv., U.S. Dep. Interior. U.S. Gov. Print. Office, Washington, DC.




# MISCELLANEOUS



*D.A. Wysocki, P.J. Schoeneberger, and E.C. Benham, NRCS, Lincoln, NE*


## **PERCENT OF AREA COVERED**

---


The following graphics are Area Percent Covered used to describe "Amount" or "Quantity." (**NOTE:** Within each large box [e.g., 2%], a quadrant contains the same total area covered but contains different object sizes and numbers.)











15%



20%




25%




30%








35%




40%




45%




50%




60%




70%



75%



80%



90%



Field  $K_{sat}$  is an important soil property and its measurement is greatly preferred over laboratory determined or mathematically predicted  $K_{sat}$  values. Field  $K_{sat}$  reflects horizon, pedon, and larger-scale macropore networks that strongly influence water flow in soils. Various methods exist for field  $K_{sat}$  measurement (Soil Survey Staff, 1982; Bouma et al., 1982; Amoozegar and Warrick, 1986; Soil Survey Staff, 2009).



Where measured  $K_{sat}$  values are unavailable, mathematical models or predictions can provide approximate estimates. Such  $K_{sat}$  estimates rely on other estimated or measured soil physical properties (e.g., texture, bulk density, porosity). Estimated  $K_{sat}$  values are assigned as a class range to compare soils and are not used as a  $K_{sat}$  value for a specific site (Soil Survey Division Staff, 1993).

A general guide for estimating  $K_{sat}$  classes (Soil Survey Staff 2012; Rawls and Brakensiek, 1983) involves a set of textural triangles that group soils by relative bulk density (low, medium, or high) and soil texture. Use the following charts and steps to predict  $K_{sat}$  by class range.



**Step 1** - Use an estimated or known bulk density and texture to select the appropriate bulk density triangle for the layer. [e.g., a clay loam (35% sand and 35% clay) with a  $1.4 \text{ g cm}^{-3}$  bulk density fits the *Medium Bulk Density* triangle].

**Step 2** - Use the appropriate *Bulk Density /  $K_{sat}$  Class Triangle* to assign a  $K_{sat}$  class. [e.g., a clay loam texture (35% clay and 35% sand) with medium bulk density assigns an estimated *Moderately Low* (0.1 – 1.0  $\mu\text{m/sec}$ )  $K_{sat}$  Class].









## **SOIL WATER REPELLENCY (DISCUSSION)**

---

Water repellency is a soil's ability to resist spontaneous wetting when water is placed on a soil surface. Water-repellent compounds arise from organic matter decomposition, including plant root exudates, fungal processes, and surface waxes from plant leaves (Mainwaring et al., 2004). Initial, irreversible drying of organic materials (Hallett et al., 2003) causes hydrophilic functional groups to strongly bond with each other and soil particles. This process results in an exposed water-repellent surface (Dekker et al., 1998). Soil water repellency is a dynamic property that varies with climate, plant community, and microbial decomposition pathways. Fire is also an important factor; heat volatilizes water-repellent organic substances that move and condense where soil is cooler (Savage, 1974). Fire may have a 3- to 5-year influence on water repellency (DeBano and Krammes, 1966). Water repellency decreases water infiltration and increases surface runoff and soil erodibility.

Organic compounds, particularly fats, waxes, and resins, form repellent coatings on mineral grains. The repellency degree depends on the quantity of particles covered (Doerr et al., 2006). Sandy soils (with low surface area) are more prone to water repellency than are loamy or clayey soils. Soil water repellency is spatially variable both laterally and with depth (Robichaud and Miller, 1999; Hubbert et al., 2006). Soil moisture content strongly influences water repellency. Soils that are more than about 10% moisture wet more readily than dry soils (Hubbert and Oriol, 2005; MacDonald and Huffman, 2004).

A common technique to assess water repellency is the Water Drop Penetration Test (WDPT). A water drop is placed on a clean soil surface and the absorption time recorded (Letey, 1969). The WDPT test measures repellency persistence (Doerr et al., 2006). Water may penetrate instantly or take hours. Various time classifications relate water repellency to absorption time (Robichaud et al., 2008). Such classifications have convenient intervals that allow relative comparison, but the times do not have intrinsic physical meaning.

## **SOIL WATER REPELLENCY**

---

Evaluate and record the relative soil water repellency determined from a Water Drop Penetration Time (WDPT) measurement.  
(**NOTE:** Soil should be in a dry state.)

### **WDPT PROCEDURE**

- 1) Prepare with a knife or trowel a clean, level horizontal 15 x 15 cm area of soil at a desired depth.
- 2) Use an eyedropper or plastic squeeze bottle to randomly place 5 drops of distilled water (each drop approximately 5 mm in diameter) from a 1-cm height onto the prepared surface.
- 3) Record the average time (in seconds) that the drops remain on the surface before absorption.

Determine the relative water repellency class according to the following table.

<b>Relative Water Repellency Class</b>	<b>Code</b>	<b>Absorption Time (seconds)</b>
Non-Water Repellent	NWR	0 to 5
Slightly Water Repellent	SWR	> 5 to 60
Moderately Water Repellent	MWR	> 60 to 180
Strongly Water Repellent	TWR	> 180

Modified from Robichaud, 2008.

# MEASUREMENT EQUIVALENTS AND CONVERSIONS

## METRIC TO ENGLISH

Known	Symbol	Multiplier	Product	Symbol
<b>LENGTH</b>				
micrometers (microns) (=10,000 Angstrom units)	$\mu m$	3.9370 $\times 10^{-5}$	inches	<i>in</i> or " <i>"</i>
millimeters	<i>mm</i>	0.03937	inches	<i>in</i> or " <i>"</i>
centimeters	<i>cm</i>	0.0328	feet	<i>ft</i> or ' <i>'</i>
centimeters	<i>cm</i>	0.3937	inches	<i>in</i> or " <i>"</i>
meters	<i>m</i>	3.2808	feet	<i>ft</i> or ' <i>'</i>
meters	<i>m</i>	1.0936	yards	<i>yd</i>
kilometers	<i>km</i>	0.6214	miles (statute)	<i>mi</i>
<b>AREA</b>				
square centimeters	$cm^2$	0.1550	square inches	$in^2$
square meters	$m^2$	10.7639	square feet	$ft^2$
square meters	$m^2$	1.1960	square yards	$yd^2$
square kilometers	$km^2$	0.3861	square miles	$mi^2$
hectares	<i>ha</i>	2.471	acres	<i>ac</i>
<b>VOLUME</b>				
cubic centimeters	$cm^3$	0.06102	cubic inches	$in^3$
cubic meters	$m^3$	35.3146	cubic feet	$ft^3$
cubic meters	$m^3$	1.3079	cubic yards	$yd^3$
cubic meters	$m^3$	0.0008107	acre-feet (=43,560 $ft^3$ )	<i>acre-ft</i>
cubic kilometers	$km^3$	0.2399	cubic miles	$mi^3$
liters (=1000 $cm^3$ )	<i>l</i>	1.0567	quarts (U.S.)	<i>qt</i>
liters	<i>l</i>	0.2642	gallons (U.S.)	<i>gal</i>
milliliter	<i>ml</i>	0.0338	fluid ounces	<i>oz</i>
1 milliliter=1 $cm^3$ =1 gm ( $H_2O$ , at 25 °C)				
<b>MASS</b>				
grams	<i>g</i>	0.03527	ounces (avdp.)	<i>oz</i>
kilograms	<i>kg</i>	2.2046	pounds (avdp.)	<i>lb</i>
megagrams (= metric tons)	<i>Mg</i>	1.1023	short tons (2000 lb)	
megagrams	<i>Mg</i>	0.9842	long tons (2240 lb)	

## ENGLISH TO METRIC

Known	Symbol	Multiplier	Product	Symbol
<b>LENGTH</b>				
inches	<i>in or "</i>	$2.54 \times 10^4$	micrometers (microns) [=10,000 Angstrom units (A)]	$\mu m$
inches	<i>in or "</i>	2.54	centimeters	<i>cm</i>
feet	<i>ft or '</i>	30.48	centimeters	<i>cm</i>
feet	<i>ft or '</i>	0.3048	meters	<i>m</i>
yards	<i>yd</i>	0.9144	meters	<i>m</i>
miles (statute)	<i>mi</i>	1.6093	kilometers	<i>km</i>
<b>AREA</b>				
square inches	<i>in<sup>2</sup></i>	6.4516	square centimeters	<i>cm<sup>2</sup></i>
square feet	<i>ft<sup>2</sup></i>	0.0929	square meters	<i>m<sup>2</sup></i>
square yards	<i>yd<sup>2</sup></i>	0.8361	square meters	<i>m<sup>2</sup></i>
square miles	<i>mi<sup>2</sup></i>	2.59	square kilometers	<i>km<sup>2</sup></i>
acres	<i>ac</i>	0.405	hectares	<i>ha</i>
<b>VOLUME</b>				
acre-feet	<i>acre-ft</i>	1233.5019	cubic meters	<i>m<sup>3</sup></i>
acre-furrow-slice $\approx 2,000,000$ lbs	<i>afs</i> ( <i>assumes b.d. = 1.3 g/cm<sup>3</sup></i> )	=6-in.-thick layer that's 1 acre in area		
cubic inches	<i>in<sup>3</sup></i>	16.3871	cubic centimeters	<i>cm<sup>3</sup></i>
cubic feet	<i>ft<sup>3</sup></i>	0.02832	cubic meters	<i>m<sup>3</sup></i>
cubic yards	<i>yd<sup>3</sup></i>	0.7646	cubic meters	<i>m<sup>3</sup></i>
cubic miles	<i>mi<sup>3</sup></i>	4.1684	cubic kilometers	<i>km<sup>3</sup></i>
gallons (U.S.) (=0.8327 Imperial gal)	<i>gal</i>	3.7854	liters	<i>l</i>
quarts (U.S.)	<i>qt</i>	0.9463	liters (=1000 cm <sup>3</sup> )	<i>l</i>
ounces	<i>oz</i>	29.57	milliliters	<i>ml</i>
1 milliliter = 1 cm <sup>3</sup> = 1 gm (H <sub>2</sub> O, at 25 °C)				
<b>MASS</b>				
ounces (avdp.)	<i>oz</i>	28.3495	grams	<i>g</i>
ounces (avdp.)	( <i>1 troy oz. = 0.083 lb</i> )			
pounds (avdp.)	<i>lb</i>	0.4536	kilograms	<i>kg</i>
short tons (2000 lb)		0.9072	megagrams (= metric tons)	<i>Mg</i>
long tons (2240 lb)		1.0160	megagrams	<i>Mg</i>

## COMMON CONVERSION FACTORS

Known	Symbol	Multiplier	Product	Symbol
acres	<i>ac</i>	0.405	hectares	<i>ha</i>
acre-feet	<i>acre-ft</i>	1233.5019	cubic meters	<i>m<sup>3</sup></i>
acre-furrow-slice ≈ 2,000,000 lbs	<i>afs</i> ( <i>assumes b.d. = 1.3 g/cm<sup>3</sup></i> )	=6-in.-thick layer that's 1 acre square		
Angstrom units	<i>A</i>	1x 10 <sup>-8</sup>	centimeters	<i>cm</i>
Angstrom units	<i>A</i>	1x 10 <sup>-4</sup>	micrometers	<i>um</i>
Atmospheres	<i>atm</i>	1.0133 x 10 <sup>6</sup>	dynes/cm <sup>2</sup>	
Atmospheres	<i>atm</i>	760	mm of mercury (Hg)	
BTU (mean)	<i>BTU</i>	777.98	foot-pounds	
centimeters	<i>cm</i>	0.0328	feet	<i>ft or '</i>
centimeters	<i>cm</i>	0.3937	inches	<i>in or "</i>
centimeters/hour	<i>cm/hr</i>	0.3937	inches/hour	<i>in/hr</i>
centimeters/second	<i>cm/s</i>	1.9685	feet/minute	<i>ft/min</i>
centimeters/second	<i>cm/s</i>	0.0224	miles/hour	<i>mph</i>
chain (U.S.)		66	feet	<i>ft</i>
chain (U.S.)		4	rods	
cubic centimeters	<i>cm<sup>3</sup></i>	0.06102	cubic inches	<i>in<sup>3</sup></i>
cubic centimeters	<i>cm<sup>3</sup></i>	2.6417 x 10 <sup>-4</sup>	gallons (U.S.)	<i>gal</i>
cubic centimeters	<i>cm<sup>3</sup></i>	0.999972	milliliters	<i>ml</i>
cubic centimeters	<i>cm<sup>3</sup></i>	0.0338	ounces (U.S.)	<i>oz</i>
cubic feet	<i>ft<sup>3</sup></i>	0.02832	cubic meters	<i>m<sup>3</sup></i>
cubic feet (H <sub>2</sub> O, 60 °F)	<i>ft<sup>3</sup></i>	62.37	pounds	<i>lbs</i>
cubic feet	<i>ft<sup>3</sup></i>	0.03704	cubic yards	<i>yd<sup>3</sup></i>
cubic inches	<i>in<sup>3</sup></i>	16.3871	cubic centimeters	<i>cm<sup>3</sup></i>
cubic kilometers	<i>km<sup>3</sup></i>	0.2399	cubic miles	<i>mi<sup>3</sup></i>
cubic meters	<i>m<sup>3</sup></i>	35.3146	cubic feet	<i>ft<sup>3</sup></i>
cubic meters	<i>m<sup>3</sup></i>	1.3079	cubic yards	<i>yd<sup>3</sup></i>
cubic meters	<i>m<sup>3</sup></i>	0.0008107	acre-feet (=43,560 ft <sup>3</sup> )	<i>acre-ft</i>
cubic miles	<i>mi<sup>3</sup></i>	4.1684	cubic kilometers	<i>km<sup>3</sup></i>
cubic yards	<i>yd<sup>3</sup></i>	0.7646	cubic meters	<i>m<sup>3</sup></i>
degrees (angle)	°	0.0028	circumfrences	
Faradays		96500	coulombs (abs)	
fathoms		6	feet	<i>ft</i>

<b>Known</b>	<b>Symbol</b>	<b>Multiplier</b>	<b>Product</b>	<b>Symbol</b>
feet	<i>ft or ' </i>	30.4801	centimeters	<i>cm</i>
feet	<i>ft or ' </i>	0.3048	meters	<i>m</i>
feet	<i>ft or ' </i>	0.0152	chains (U.S.)	
feet	<i>ft or ' </i>	0.0606	rods (U.S.)	
foot pounds		0.0012854	BTU (mean)	<i>BTU</i>
gallons (U.S.)	<i>gal</i>	3.7854	liters	<i>l</i>
gallons (U.S.)	<i>gal</i>	0.8327	Imperial gallons	
gallons (U.S.)	<i>gal</i>	0.1337	cubic feet	<i>ft<sup>3</sup></i>
gallons (U.S.)	<i>gal</i>	128	ounces (U.S.)	<i>oz</i>
grams	<i>g</i>	0.03527	ounces (avdp.)	<i>oz</i>
hectares	<i>ha</i>	2.471	acres	<i>ac</i>
horsepower		2545.08	BTU (mean)/hour	
inches	<i>in or " </i>	2.54 x 10 <sup>4</sup>	micrometers (micron) (=10,000 Angstrom units [A])	<i>μm</i>
inches	<i>in or " </i>	2.5400	centimeters	<i>cm</i>
inches/hour	<i>in/hr</i>	2.5400	centimeters/hour	<i>cm/hr</i>
inches/hour	<i>in/hr</i>	7.0572	micrometers/sec	<i>μm/sec</i>
kilograms	<i>kg</i>	2.2046	pounds (avdp.)	<i>lb</i>
kilometers	<i>km</i>	0.6214	miles (statute)	<i>mi</i>
joules	<i>J</i>	1 x 10 <sup>7</sup>	ergs	
liters	<i>l</i>	0.2642	gallons (U.S.)	<i>gal</i>
liters	<i>l</i>	33.8143	ounces	<i>oz</i>
liters (=1000 cm <sup>3</sup> )	<i>l</i>	1.0567	quarts (U.S.)	<i>qt</i>
long tons (2240 lb)		1.0160	megagrams	<i>Mg</i>
megagrams (= metric tons)	<i>Mg</i>	1.1023	short tons (2000 lb)	
megagrams	<i>Mg</i>	0.9842	long tons (2240 lb)	
meters	<i>m</i>	3.2808	feet	<i>ft or ' </i>
meters	<i>m</i>	39.37	inches	<i>in</i>
micrometers (microns)	<i>μm</i>	1.000	microns	<i>μ</i>
micrometers/second	<i>μm/sec</i>	0.1417	inches/hour	<i>in/hr</i>
micron	<i>μ</i>	1 x 10 <sup>-4</sup>	centimeters	<i>cm</i>
microns	<i>μ</i>	3.9370	inches	<i>in or " </i>

<b>Known</b>	<b>Symbol</b>	<b>Multiplier</b>	<b>Product</b>	<b>Symbol</b>
(=10,000 Angstrom units)		$\times 10^{-5}$		
micron	$\mu$	1.000	micrometer	$\mu m$
miles (statute)	<i>mi</i>	1.6093	kilometers	<i>km</i>
miles/hour	<i>mph</i>	44.7041	cent./second	<i>cm/s</i>
miles/hour	<i>mph</i>	1.4667	feet/second	<i>ft/s</i>
milliliter	<i>ml</i>	0.0338	fluid ounces	<i>oz</i>
1 milliliter $\approx 1 \text{ cm}^3 = 1 \text{ gm (H}_2\text{O, at 25}^\circ\text{C)}$				
milliliter	<i>ml</i>	1.000028	cubic centimeters	$\text{cm}^3$
millimeters	<i>mm</i>	0.03937	inches	<i>in or "</i>
ounces	<i>oz</i>	29.5729	milliliters	<i>ml</i>
1 milliliter $\approx 1 \text{ cm}^3 = 1 \text{ gm (H}_2\text{O, at 25 }^\circ\text{C)}$				
ounces (avdp.)	<i>oz</i>	28.3495	grams	<i>g</i>
ounces (avdp.) 1 troy oz.=0.083 lb				
pints (U.S.)	<i>pt</i>	473.179	cubic centimeters	$\text{cm}^3$ <i>or cc</i>
pints (U.S.)	<i>pt</i>	0.4732	liters	<i>l</i>
pounds (avdp.)	<i>lb</i>	0.4536	kilograms	<i>kg</i>
quarts (US liquid)	<i>qt</i>	0.9463	liters (=1000 $\text{cm}^3$ )	<i>l</i>
rods (U.S.)		0.25	chains (U.S.)	<i>ft</i>
rods (U.S.)		16.5	feet (U.S.)	<i>ft</i>
short tons (2000 lb)		0.9072	megagrams (= metric tons)	<i>Mg</i>
square centimeters	$\text{cm}^2$	0.1550	square inches	$\text{in}^2$
square feet	$\text{ft}^2$	0.0929	square meters	$\text{m}^2$
square inches	$\text{in}^2$	6.4516	sq. centimeters	$\text{cm}^2$
square kilometers	$\text{km}^2$	0.3861	square miles	$\text{mi}^2$
square meters	$\text{m}^2$	10.7639	square feet	$\text{ft}^2$
square meters	$\text{m}^2$	1.1960	square yards	$\text{yd}^2$
square miles	$\text{mi}^2$	2.5900	square kilometers	$\text{km}^2$
square yards	$\text{yd}^2$	0.8361	square meters	$\text{m}^2$
yards	<i>yd</i>	0.9144	meters	<i>m</i>



## GUIDE TO MAP SCALES AND MINIMUM SIZE DELINEATIONS <sup>1</sup>

Order of Soil Survey	Map Scale	Inches Per Mile	Minimum Size Delineation <sup>2</sup>	
			Acres	Hectares
Order 1	1:500	126.7	0.0025	0.001
	1:1,000	63.4	0.100	0.004
	1:2,000	31.7	0.040	0.016
	1:5,000	12.7	0.25	0.10
	1:7,920	8.0	0.62	0.25
	1:10,000	6.34	1.00	0.41
Order 2	1:12,000	5.28	1.43	0.6
	1:15,840	4.00	2.50	1.0
	1:20,000	3.17	4.00	1.6
	1:24,000 <sup>3</sup>	2.64	5.7	2.3
Order 3	1:30,000	2.11	9.0	3.6
	1:31,680	2.00	10.0	4.1
Order 4	1:60,000	1.05	36	14.5
	1:62,500 <sup>4</sup>	1.01	39	15.8
	1:63,360	1.00	40	16.2
Order 5	1:80,000	0.79	64	25.8
	1:100,000	0.63	100	40
	1:125,000	0.51	156	63
	1:250,000	0.25	623	252
	1:500,000	0.127	2,500	1,000
	1:750,000	0.084	5,600	2,270
Very General	1:1,000,000	0.063	10,000	4,000
	1:7,500,000	0.0084	560,000	227,000
	1:15,000,000	0.0042	2,240,000	907,000

<sup>1</sup> Modified from Peterson, 1981.














<sup>2</sup> Traditionally, the minimum size delineation is assumed to be a 1/4-inch square, or a circle with an area of 1/16 inch<sup>2</sup>. Cartographically, this is about the smallest area in which a conventional soil map symbol can be legibly printed. Smaller areas can be, but rarely are, delineated and the symbol "lined in" from outside the delineation.
















<sup>3</sup> Corresponds to USGS 7.5-minute topographic quadrangle maps.












<sup>4</sup> Corresponds to USGS 15-minute topographic quadrangle maps.







## COMMON SOIL MAP SYMBOLS (TRADITIONAL)

(From Soil Survey Staff, 1990.) The following symbols are common on field sheets (original aerial photograph–based soil maps) and in many soil surveys published prior to 1997. Current guidelines for map compilation symbols are in NSSH, Exhibit 627-5, Feature and Symbol Legend for Soil Survey (Soil Survey Staff, 2012).













FEATURE	SYMBOL
<b>LANDFORM FEATURES</b>	
SOIL DELINEATIONS	
ESCARPMENTS	
Bedrock	 (Points down slope)
Other than bedrock	 (Points down slope)
SHORT STEEP SLOPE	
GULLY	
DEPRESSION, closed	
SINKHOLE	
Prominent hill or peak	
<b>EXCAVATIONS</b>	
Soil sample site (Type location, etc.)	
Borrow pit	
Gravel pit	
Mine or quarry	
LANDFILL	

FEATURE	SYMBOL
<b>MISCELLANEOUS SURFACE FEATURES</b>	
Blowout	
Clay spot	
Gravelly spot	
Lava flow	
Marsh or swamp	
Rock outcrop (includes sandstone and shale)	
Saline spot	
Sandy spot	
Severely eroded spot	
Slide or slip (tips point upslope)	
Sodic spot	
Spoil area	
Stony spot	
Very stony spot	
Wet spot	

FEATURE	SYMBOL
<b>ROAD EMBLEMS</b>	
Interstate	 
Federal	 
State	 
County, farm, or ranch	
<b>CULTURAL FEATURES</b>	
RAILROAD	
POWER TRANSMISSION LINE (normally not shown)	
PIPELINE (normally not shown)	
FENCE (normally not shown)	

FEATURE	SYMBOL
<b><i>CULTURAL FEATURES (continued)</i></b>	
LEVEES	
Without road	
With road	
With railroad	
Single side slope (showing actual feature location)	
DAMS	
Medium or small	
Large	

FEATURE	SYMBOL
<b>HYDROGRAPHIC FEATURES</b>	
<b>STREAMS</b>	
Perennial, double line (large)	
Perennial, single line (small)	
Intermittent	
Drainage end or flow direction	
<b>SMALL LAKES, PONDS, AND RESERVOIRS</b>	
Perennial water	
Miscellaneous water	
Flood pool line	
Lake or pond (perennial)	
<b>MISCELLANEOUS WATER FEATURES</b>	
Spring	
Well, artesian	
Well, irrigation	

FEATURE	SYMBOL
<b>MISCELLANEOUS CULTURAL FEATURES</b>	
Airport	
Cemetery	
Farmstead, house (omit in urban areas)	
Church	
School	
Other religion (label)	 Mt. Carmel
Located object (label)	 Ranger Station
Tank (label)	 Petroleum
Lookout tower	
Oil and/or natural gas wells	
Windmill	
Lighthouse	

## REFERENCES

---

- Amoozegar, F. and A.W. Warrick. 1986. Hydraulic conductivity of saturated soils: Field methods. *In* A. Klute (ed). 1986. Methods of soil analysis part 1. Physical and mineralogical methods, 2nd ed. ASA, Agron. Mono. 9, Madison, WI, pp. 735-770.
- DeBano, L.F., and J.S. Krammes. 1966. Water repellent soils and their relation to wildfire temperatures. *Int. Assoc. Sci. Hydrol. Bull.*, XI Annals 2, pp. 14-19.
- DeBano, L.F. 1981. Water repellent soils: A state-of-the-art. Gen. Tech. Rpt. PSW-46. Berkeley, CA: USDA, Forest Service, Pacific Southwest Forest and Range Experiment Station. 21 p.
- Dekker, L.W., C.J. Ritsema, K. Oostindie, and O.H. Boersma. 1998. Effect of drying temperature on the severity of soil water repellency. *Soil Sci.* 163:780-796.
- Doerr, S.H., R.A. Shakesby, L.W. Dekker, and C.J. Ritsema. 2006. Occurrence prediction and hydrological effects of water repellency amongst major soil and land-use types in a humid temperate climate. *Eur. J. Soil Sci.* 57:741-754.
- Hallett, P.D., D.C. Gordon, and A.G. Bengough. 2003. Plant influence on rhizosphere hydraulic properties: Direct measurements using a miniaturised infiltrometer. *New Phytologist* 157:597-603.
- Hubbert, K.R., and V. Oriol. 2005. Temporal fluctuations in soil water repellency following wildfire in chaparral steeplands, southern California. *Intern. J. Wildland Fire* 14:439-447.
- Hubbert, K.R., H.K. Preisler, P.M. Wohlgemuth, R.C. Graham, and M.G. Narog. 2006. Prescribed burning effects on soil physical properties and soil water repellency in a steep chaparral watershed, southern California. *U.S.A. Geoderma.* 130:284-298.
- Mainwaring, K.A., C.P. Morley, S.H. Doerr, P. Douglas, C.T. Llewellyn, G. Llewellyn, I. Matthews, and B.K. Stein. 2004. Role of heavy polar organic compounds for water repellency of sandy soils. *Env. Chem. Letters* 2:35-39.
- Letey, J. 1969. Measurement of contact angle, water drop penetration time, and critical surface tension. *Proceed. of the Symp. on Water-Repellent Soils*, 6-10 May 1968, UC, Riverside, pp. 43-47.
- Peterson, F.F. 1981. Landforms of the Basin and Range Province defined for soil survey. *Nevada Agric. Exp. Stn. Tech. Bull.* 28. Reno, NV.



Rawls, W.J., and D.L. Brakensiek. 1983. A procedure to predict Green and Ampt infiltration parameters. *In* Advances in infiltration. Proc. of the National Conf. on Advances in Infiltration, Dec. 12-13. Chicago, IL.

Robichaud, P.R., and S.M. Miller. 1999. Spatial interpolation and simulation of post-burn duff thickness after prescribed fire. *Intern. J. Wildland Fire* 9:137-143.

Robichaud, P.R., S.A. Lewis, and L.E. Ashmun. 2008. New procedure for sampling infiltration to assess post-fire soil water repellency. USDA, Forest Service, Rocky Mountain Research Station, Research Note RMRS-RN-33.

Savage, S.M. 1974. Mechanism of fire-induced water repellency in soil. *Soil Sci. Soc. Amer. Proc.* 38:652-657.

Soil Survey Division Staff. 1993. Soil survey manual. USDA, SCS, Agric. Handb. 18. U.S. Gov. Print. Office, Washington, DC.

Soil Survey Staff. 1982. Measuring hydraulic conductivity for use in soil surveys. Soil Survey Investigations Report No. 38, USDA, SCS, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 1990. National soil survey handbook (NSSH), Title 170, Part 601. USDA, NRCS, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 2009. Soil survey field and laboratory methods manual. Soil Surv. Invest. Rep. 51, ver. 1.0. R. Burt (ed.) USDA, NRCS.

Soil Survey Staff. 2012. National soil survey handbook (NSSH) [online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE. (<http://soils.usda.gov/technical/handbook/>)



# SOIL SAMPLING

*P.J. Schoeneberger, D.A. Wysocki, and E.C. Benham, NRCS, Lincoln, NE*

## **INTRODUCTION**

---

Laboratory measurement of soil properties (e.g., particle size, organic carbon, etc.) requires sample collection. Two fundamental sampling requirements are: 1) appropriate site selection, and 2) a detailed soil description. Soils are landscape entities that reflect geomorphic, pedologic, and hydrologic processes and parent material distribution (Wysocki et al., 2011). Thus, site selection, regardless of purpose, must consider soilscape relationships. A soil profile description identifies the horizons and their thickness and provides context for data collection and interpretation. Soil property data by itself has little value without context; soil data requires an accompanying geo-referenced description. Sampling needs and strategies vary by project objectives. Various reviews and summaries (Buol et al., 2003; Soil Survey Staff, 2004; Soil Survey Staff, 2009; Robertson et al., 1999; Carter, 1993) outline sampling strategies and techniques.

Statistical design and analysis (e.g., random, randomized block, grid, transect, traverse, geostatistical) are important aspects of sample collection (Buol et al., 2003). Discussion of statistical methods and design is beyond the scope of this publication. Summary information is available in Webster and Oliver (2007) and Webster and Oliver (1990). Note, however, that statistical blocking by geomorphic context stratifies soil areas by similar geologic and pedogenic processes. Random sampling within a geomorphically stratified area allows determination of both random soil variation and systematic landscape variation (Hall and Olson, 1991).

**SOILSCAPE SEQUENCES**—Soil sampling commonly considers pedons as distinct points separate from adjoining soils. Soil water flow is generally interpreted as predominantly vertical; lateral flow receives considerably less emphasis. In many soilscaapes, however, vertical flow is important but lateral flow is more influential. A slight difference in elevation (15 cm) in nearly level landscapes produces substantial hydrologic and morphological differences (Knuteson et al., 1989). Soilscape sampling and characterization is an important strategy for increasing and organizing both spatial and soil property data (Wysocki et al., 2011). This approach evaluates landscape-scale processes that relate ecosystem dynamics to soil distribution.

## **TYPES OF SAMPLING**


---

**HORIZON SAMPLING**—Cost and time limit the number of sample collections. The most effective and efficient strategy for sampling

is by horizon. Soil horizons develop from natural processes acting over time. Variations in soil properties may occur within a horizon, but distinct differences generally occur between horizons. Consequently, soil horizons are a meaningful and comparable schema for sample collection (see graphic). Soil horizons vary in thickness and boundary (e.g., wavy, irregular, broken) within a pedon and across landscapes. Soil profile descriptions and horizon sampling techniques incorporate depth and boundary variability and can produce homogeneous samples. It is advisable to subsample soil horizons >50 cm thick. Fixed-depth sampling alone does not capture such variation and may lead to the erroneous interpretation of data.

**INCREMENTAL SAMPLING**—Project objectives (e.g., soil genesis or archeological) may require within-horizon detail. Property variation or trends within horizons require samples at specified increments (e.g., every 10 cm). Increment samples should be taken within horizons; sample depths should not cross horizon boundaries. Increment sampling provides more detail than horizon sampling but adds time and expense. This approach is generally limited to special projects.

**FIXED-DEPTH SAMPLING**—Specified objectives (e.g., surface compaction studies) may address properties by fixed depths (e.g., 0-5 cm or 5-10 cm) instead of by soil horizons. This approach, while appropriate for certain purposes, precludes data comparison by horizon. Data collected by depth is comparable within a study and to other studies employing the same depths. Fixed-depth samples may straddle horizons that contain contrasting materials (e.g., sandy over clayey strata). The resultant data represents neither horizon and is difficult to interpret. Use this approach with caution.



## **SAMPLING TECHNIQUES**

---

Excavate a fresh soil pit for sample collection. Avoid road cuts as sample sites because dust and exposure can alter soil properties. If excavation is not possible, collect samples from intact cores (e.g., Giddings tube). If an auger is the only means for sample collection, place the collected soil onto a tarp to identify and sample by horizon. For soil characterization, collect 3- to 4-kg samples. Soils with fragments up to 20 mm in size require a minimum 1 kg (dry weight) sample for a representative quantity (ASTM, 2004). If fragment size exceeds 20 mm, larger sample sizes are needed.

During sample collection:

- 1) Collect samples in a soil pit from the bottom up. This minimizes contamination by falling debris.
- 2) Collect the sample across a horizon's full depth and breadth. Avoid atypical pockets or lenses, or subsample these separately.

## **SOIL SAMPLE KINDS**

---

**CHARACTERIZATION SAMPLES**—Samples collected from a fully described soil chosen to be representative of a given soil series or soil landscape area. All horizons are bulk sampled to a depth of 200 cm or more. In addition, three fist-sized clods are collected from each horizon for bulk density measurement. Bulk samples undergo a suite of analyses (e.g., particle size, 1500 kPa water content, CEC, pH, extractable cations, organic carbon, clay mineralogy, etc.). The specified analyses vary with soil composition. A Mollisol needs a different set of analyses for characterization than does an Andisol or Spodosol. Characterization sampling provides a complete set of measured values for soil comparison or extrapolation.

**REFERENCE SAMPLES** (grab samples)—Samples collected for a single or limited set of analyses (e.g., OC, PSA, pH) to answer a specific question. For example, there may be a question of whether or not the A horizon has sufficient organic carbon for mollic epipedon criteria ( $OC \geq 0.6\%$ ). Reference samples are generally targeted to specific horizons or layers (e.g., A horizon, Bt horizon, control section) in a profile. Sample intent is to answer a question quickly with little expense.

## **FIELD EQUIPMENT CHECKLIST**

---

### **1. Digging Tools (commonly choose 1 or 2): see graphic**

Bucket Auger  
Sharp Shooter  
Montana Sharp Shooter (for rocky soils)  
Tile Spade (only for well cultivated or loose material)  
Spade (standard shovel)  
Push Probe (e.g., Backsaver®, Oakfield®)—include a clean-out tool  
Pulaski

### **Soil Description**

Knife  
Hand Lens (10X or combination lenses)  
Acid Bottle (1N HCl)  
Water Bottle  
Color Book (e.g., Munsell®, EarthColors®)  
Picture Tapes ("pit tape"—metric preferred)  
Tape Measure (metric or English and metric)  
(3) Ultra-Fine Point Permanent Marker Pens  
Pocket pH Kit or Electronic "Wand"  
Pocket Soil Thermometer  
Camera  
Sample bags (for grab samples)  
Soil Description Sheet (232 or PEDON description form)

### **Site Description**

Field Note Book  
GPS Unit  
Abney Level  
Clinometer  
Compass  
Altimeter (pocket-sized)

### **Field References**












Field Book for Describing and Sampling Soils  
Aerial Photographs  
Topographic Maps (1:24,000, 7.5 min; 1:100,000)  
Geology Maps  
Soil Surveys (county or area)  
AGI Field Sheets

### **Personal Protective Gear**

Small First Aid Kit  
Leather Gloves  
Sunglasses  
Insect Repellent  
Sunscreen  
Hat  
Drinking water


## **EXAMPLES OF COMMON SOIL-SAMPLING EQUIPMENT**

(Use of trade or company names is for informational purposes only and does not constitute an endorsement.)

<b>Digging Tools/Shovel Types</b>				
				
Pulaski	Standard shovel	Tile spade	Sharp-shooter	Montana sharp-shooter
<i>Primary use:</i>	<i>most materials</i>	<i>loose material</i>	<i>most materials</i>	<i>rocky soil</i>
<b>Soil Probes</b>			<b>Hydraulic Probes</b>	
				
Regular push-tube	Tile probe (solid steel rod)	Peat sampler (Macaulay)	Giddings tube	Bull probe
<i>Primary use:</i>	<i>locating hard contact</i>	<i>organic soils</i>	<i>(not effective in rocky materials)</i>	<i>wet sands, organics (no co. frags)</i>
<i>fine earth</i>				
				Vibracore


## Bucket Auger Types

Open




Regular auger  
(open teeth)

Closed



Closed bucket  
(open teeth)



Sand auger  
(pinched teeth)

Primary  
use:

*clays, loams*

*loams*

*moist sand*

## External Thread Augers



Dutch auger  
("mud auger")

Primary use: *organics, moist clay, muds*




Screw auger  
(external threads)

*rocky soils*



Flight auger

*rocky soils, deep holes*



Hollow stem  
auger

*undisturbed sample*

## REFERENCES

American Society for Testing and Materials (ASTM). 2004. Standard practice for description and identification of soils. D2488 Annual Book of ASTM Standards. Construction. Section 4, Soil and Rock; Dimension Stone. Vol. 04.08. Philadelphia, PA.

Buol, S.W., R.J. Southard, R.C. Graham, and P.A. McDaniel. 2003. Soil genesis and classification. 5th edition. Iowa State Press, Ames, IA. ISBN: 0-8138-2873-2.

Carter, M. (editor). 1993. Soil sampling and methods of analysis. Lewis Publishers, CRC Press, Inc., Boca Raton, FL. ISBN: 0-87371-861-5.



Hall, G.F., and C.G. Olson. 1991. Predicting variability of soils from landscape models. *In* M.J. Mausbach and L.P. Wilding (eds.) *Spatial variability of soils and landforms*. Soil Sci. Soc. Am. Spec. Pub. #28, SSSA, Inc., Madison, WI. ISBN: 0-89118-198-7.

Knuteson, J.A., J.L. Richardson, D.D. Patterson, and L.D. Prunty. 1989. Pedogenic carbonates in a Calciaquoll associated recharge wetland. *Soil Sci. Soc. Am. J.* 53:495-499.

Robertson, G.P., D.C. Coleman, C.S. Bledsoe, and P. Sollins (editors). 1999. *Standard soil methods for long-term ecological research*. Oxford University Press, Inc., New York, NY. ISBN: 0-19-512083-3.

Soil Survey Staff. 2004. *Soil survey laboratory methods manual*. Soil Survey Invest. Rep. (SSIR) 42, Version 4.0. R. Burt (ed.), USDA, Natural Resources Conservation Service.

Soil Survey Staff. 2009. *Soil survey field and laboratory methods manual*. Soil Survey Invest. Rep. No. 51, Version 1.0. R. Burt (ed.), USDA, Natural Resources Conservation Service. Available at <http://soils.usda.gov/technical/>.

Soil Survey Staff. 2012. *National soil survey handbook (NSSH)* [Online]. USDA, NRCS, National Soil Survey Center, Lincoln, NE (<http://soils.usda.gov/technical/handbook/>).

Webster R., and M.A. Oliver. 1990. *Statistical methods in soil and land resource survey*. Oxford University Press. Oxford, England.

Webster R., and M.A. Oliver. 2007 *Geostatistics for environmental scientists*. 2nd edition. John Wiley and Sons, Inc. West Sussex, England.

Wysocki, D.A., P.J. Schoeneberger, D.R. Hirmas, and H.E. LaGarry. 2011. *Geomorphology of soil landscapes*. *In* P.M. Huang, Y. Li, and M.E. Sumner (eds.) *Handbook of Soil Sciences: Properties and Processes*, 2nd ed, pp. 29-1–29-26. CRC Press, Boca Raton, FL.



# INDEX

## A

---

- Across Slope • 1-6, 3-40
- Air Temperature • 1-1
- Alpha Alpha Dipyrindyl • 2-88
  - Location • 2-88
- Anthric Saturation • 1-16
- Anthropogenic Feature • 1-4, 3-1, 3-10, 3-20
- Area Covered (%) Example • 7-1
- Argillans • 2-32
- Artifacts • 2-49
  - Cohesion • 2-51
  - Kind • 2-50
  - Penetrability • 2-51
  - Persistence • 2-51
  - Quantity • 2-50
  - Roundness • 2-50
  - Safety • 2-52
  - Shape • 2-51
- Aspect • See Slope Aspect

## B

---

- Bathymetry • 2-97
- Bedrock • 1-22
  - (Bedrock)-Fracture Interval Class • 1-24
  - Depth to Bedrock • 1-25
  - Kind • 1-22, 5-1
  - Weathering Class • 1-25
- Biological Concentrations • 2-19, 2-22
- Biological Crusts • 2-77, 2-79
- Bottom Type • 2-100
- Boundary Distinctness • 2-6 • See *also* Horizon Boundary
  - Distinctness
- Bridging (clay) • 2-32, 2-34
- Brittleness • 2-65
- Bulk Density Satiated
  - Discussion • 2-106
- Bypass Flow • 2-75

## C

---

- Calcium Carbonate Equivalent Test • 2-88
- Carbonate Stages • 2-30
- Caret (The) (Horizon Nomenclature) • 2-5
- Carpedolith • 2-80
- Cementation Classes (Rupture Resistance) • 2-63

- Cementing Agents • 2-64
- Characterization Samples • 8-3
- Chemical Agent • 2-87
- Chemical Crusts • 2-78
- Chemical Response • 2-85
- Clay Depletions • 2-11, 2-12
- Clay Films • 2-32
- Climate • 1-1
- Coastal Marine and Estuarine (Geomorphic Environments;  
Landforms) • 3-21
- Coats • 2-32
- Cohesion (Artifacts) • 2-51
- Color Contrast • See Mottle Contrast
- Color Location • 2-9
- Color, Mechanical Condition • 2-9
- Color Physical State (NASIS) • See Soil Matrix Color - Location or  
Condition
- Color, Redoximorphic Condition • 2-9
- Common Conversion Factors • 7-18
- Common Soil Map Symbols (Traditional) • 7-22
- Common Soil-Sampling Equipment (Examples) • 8-5
- Comparison of Particle Size Classes in Different Systems (table) •  
2-45
- Compositional Texture Modifiers • 2-41
- Concentrations • 2-20
  - Boundary • 2-27
  - Color • 2-23
  - Contrast • 2-23
  - Hardness • 2-27
  - Kind • 2-20
  - Location • 2-26
  - Moisture State • 2-23
  - Quantity • 2-23
  - Shape • 2-25
  - Size • 2-23
- Concentrations (Discussion) • 2-19
- Concretions • 2-10, 2-12, 2-19, 2-20, 2-21
- Consistence • 2-62
- Contrast of Soil Mottles • 2-16, 2-18
- Coordinates • 1-2, 2-98
- Coprogenous Earth • 1-29, 2-41
- Core Descriptions • 2-111
- Core Settlement • 2-111
- County FIPS Code • 1-3
- Cracks • 2-75
  - Depth • 2-77
  - Kind • 2-76
  - Relative Frequency • 2-77
- Crust-Related Cracks • 2-75, 2-76
- Crusts • See Soil Crusts
- Crystals • 2-19, 2-20, 2-21

## **D**

---

- Date • 1-1, 2-98
- Decision Flowchart For Describing Soil Colors • 2-8
- Delineation Size (Transect) • 1-3
- Dendritic Tubular Pores • 2-73, 2-74
- Densic Contact • 1-29
- Densic Materials • 1-29
- Depositional Crust • 2-78
- Depressional (Landforms) • 3-35
- Depth (To Bedrock) • 1-25
- Describer(s) Name • 1-1, 2-98
- Desert Pavement • 2-80
- Diagnostic Characteristics - Mineral Soils • 1-29
- Diagnostic Characteristics - Organic Soils • 1-29
- Diagnostic Horizons • 1-28
- Diagnostic Properties • 1-28
- Diagnostic Subsurface Horizons • 1-28
- Diatomaceous Earth • 1-29, 2-41
- Dipyridyl • See Alpha Alpha Dipyridyl
- Dissolved Oxygen • 2-102
- Down Slope • 1-6, 3-40
- Drainage • 1-11
- Drainage Class • 1-11, 2-101
- Drainage Network • See Drainage Pattern
- Drainage Pattern • 3-1, 3-44
- Durinode • 1-29, 2-21
- Duripan • 1-28, 2-44
- Duripan Fragments • 1-27, 2-46

## **E**

---

- Earth Cover - Kind • 1-16
- Effervescence • 2-87
  - Chemical Agent • 2-87
  - Class • 2-87
- Electrical Conductivity • 2-89
- Electrical Conductivity of Subaqueous Soils • 2-106
- Elevation • 1-5, 2-98, 3-39,
- Endosaturation • 1-15
- English To Metric (conversions) • 7-17
- Eolian Deposits • 1-18
- Eolian (Geomorphic Environment; Landforms) • 3-26
- Epipedons • 1-28
- Episaturation • 1-16
- Erosion • 1-25
  - Degree Class • 1-26
  - Kind • 1-25
- Erosion Accelerated Kind • See Erosion Kind
- Erosional Lag • 2-80

Erosional (Landforms) • 3–34  
Estuarine (Geomorphic Environment; Landforms) • 3–21  
Evaporites • 1–24, 5–3, 5–7  
Excavation Difficulty • 2–69  
Excavation Difficulty Class • 2–69  
Extra-Structural Cracks • 2–75

## **F**

---

Faunal Burrows • 2–80  
Fecal Pellets • 2–22  
Ferriargillans • 2–12, 2–32  
Field Equipment Checklist • 8–4  
Field Notes • 2–90  
Field pH (NASIS) • See Reaction (pH)  
Filaments • 2–25  
Films • 2–12, 2–32  
Final Core Length • 2–112  
Finely Disseminated Materials • 2–19, 2–20, 2–21  
FIPS Code • 1–3  
Fissures • 2–75  
Flat Plains (Geomorphic Components of) • 1–10, 3–43  
Flooding • 1–13  
    Duration • 1–13  
    Frequency • 1–13  
    Months • 1–13  
Fluidity • 2–65  
Fluidity Class • 2–105  
Fluvial (Geomorphic Environment; Landforms) • 3–24  
Formation (Lithostratigraphic Unit) • 1–25, 5–12  
Fragipan • 1–28  
Fragment Hardness • See Rock and Other Fragments - Hardness  
Fragment Roundness • See Rock and Other Fragments - Roundness  
Fragments • See Rock and Other Fragments  
Freeze-thaw Crust • 2–78

## **G**

---

Geodetic Datum • 1–2, 2–98, 6–1  
Geologic Formation (NASIS) • See Formation  
Geologic Time Scale • See North American Geologic Time Scale  
Geology • 5–1  
    References • 5–13  
Geomorphic Component • 1–8, 3–41,  
    Flat Plains • 1–10, 3–43, 3–2  
    Hills • 1–8, 3–42,  
    Mountains • 1–9, 3–1, 3–43  
    Terraces, Stepped Landforms • 1–8, 3–42, 3–1  
Geomorphic Description • 1–4, 3–1, 3–11  
    References • 3–46

Geomorphic Description (Outline) • 3-10  
Geomorphic Description System • 3-1  
Geomorphic Information • 1-4  
Gilgai Microfeatures • 2-54  
Glacial and Periglacial Deposits • 1-18  
Glacial (Geomorphic Environment; Landforms) • 3-26  
Glauconite • 2-19  
GPS • 2-98  
Ground Surface • 2-6  
Group (Lithostratigraphic Unit) • 5-12  
Guide To Map Scales And Minimum-Size Delineations • 7-21  
Gully (Erosion) • 1-25  
Gully (microfeature) • 3-18, 3-25, 3-34

## **H**

---

Hard Rock Fragments • See Rock Fragments  
Hierarchical Rank of Lithostratigraphic Units • 5-12  
Hills (Geomorphic Components of) • 1-8, 3-42  
Hillslope Position • See Hillslope - Profile Position  
Hillslope - Profile Position • 1-7, 3-1, 3-41  
Horizon and Layer Designations • 2-2  
Horizon Boundary • 2-6, 2-104  
    Distinctness • 2-6, 2-104  
    Topography • 2-7  
Horizon Depth • 2-6  
Horizon Feature Kind (NASIS) • See Special Features  
Horizon & Layer Designations • 2-103, 4-1  
Horizon & Layer Designations Conversion Charts • 4-6  
Horizon Modifiers (Other) • 2-5  
    Caret • 2-5  
    Numerical Prefixes • 2-5  
    Prime • 2-5  
Horizon Subscripts • See Horizon Suffixes  
Horizon Suffixes • 2-4, 2-103, 4-3, 4-8  
Horizon Suffixes for Subaqueous Soils (Discussion) • 2-103  
Horizon Thickness • 2-6  
Human Artifacts • See Artifacts  
Hydrothermal • 3-30 • See also Volcanic and Hydrothermal  
    (Geomorphic Environments; Landforms)  
Hypocoats • 2-12, 2-32, 2-34

## **I**

---

Ice Wedge Cast • 2-80  
Igneous Rocks • 1-22, 5-1  
Igneous Rocks Chart • 5-5  
In-Place Deposits • 1-19  
Insect Casts • 2-19, 2-20, 2-22  
Inside Length • 2-111

Interbedded Rocks • 1-24, 5-3  
Interstitial Pores • 2-74  
Inter-Structural Voids • 2-73  
Interval (Transect) • 1-4  
Iron Depletions • 2-10  
Irregular Pores • 2-73, 2-74

## **K**

---

Krotovinas • 2-80  
K<sub>sat</sub> • See *also* Saturated Hydraulic Conductivity  
K<sub>sat</sub> Class Estimate • 7-10

## **L**

---

Lacustrine (Geomorphic Environment; Landforms) • 3-23  
Lamellae • 1-29, 2-25, 2-80  
Lamina • 2-80  
Land Cover • 1-16  
    Kind • 1-16  
Landform • 1-4, 3-1, 3-10, 3-12  
Landscape • 1-4, 3-1, 3-10, 3-11  
Landslide • See Mass Movement (Geomorphic Environment;  
    Landforms), Mass Movement Types for Soil Survey  
Landuse • See Earth Cover - Kind  
Limnic Materials • 1-29, 2-41  
Lithic Contact • 1-29  
Lithologic Discontinuities  
    Discussion • 2-103  
Lithostratigraphic Units • 1-25, 5-12  
Local Physiographic / Geographic Name • 1-4, 3-1, 3-9  
Location • 1-2, 6-1  
    Description • 2-100  
    References • 6-7  
Loess Doll (Kindchen, Puppy) • 2-22

## **M**

---

Major Land Resource Area • 1-3  
Manganese Effervescence Agent • See Effervescence Agent  
Manganese Effervescence Class • See Effervescence Class  
Mangans • 2-12, 2-32  
Manner of Failure • 2-65  
Map Unit • 2-100  
Marl • 1-29, 2-41  
Masses • 2-10, 2-12, 2-19, 2-20, 2-21  
Mass Movement Deposits • 1-19  
Mass Movement (Geomorphic Environment; Landforms) • 3-29  
Mass Movement Types for Soil Survey • 5-8



Master Horizons • 2-2, 4-1, 4-6  
Matrix Color • See Soil Matrix Color  
Mean Sea Level • 1-5, 3-39  
Measurement Equivalents & Conversions • 7-16  
Member (Lithostratigraphic Unit) • 5-12  
Metamorphic Rocks • 1-23, 5-2  
Metamorphic Rocks chart • 5-6  
Metric To English (conversions) • 7-16  
Microbiotic Crust • 2-77  
Microfeature • 1-4, 3-1, 3-10, 3-18  
Microfeature (terms) • 3-18  
Microrelief • 1-10, 2-54, 3-1, 3-44  
Mineral Crusts • 2-78  
Minimum Data Set • 2-90  
Miscellaneous • 7-1  
Miscellaneous Field Notes • 2-90  
MLRA • 1-3  
Month / Day / Year • 1-1  
Mottles • 2-18  
    Color • 2-18  
    Contrast (also called Color Contrast) • 2-18  
    Moisture State • 2-18  
    Quantity • 2-18  
    Shape • 2-18  
    Size • 2-18  
Mountains (Geomorphic Components of) • 3-43  
Multicolored Pattern • 2-9

## **N**

---

Name • 1-1  
Nodules • 2-10, 2-12, 2-19, 2-20, 2-21  
North American Geologic Time Scale • 5-9

## **O**

---

Observation Method • 2-1, 2-101  
    Kind • 2-1, 2-101  
    Relative Size • 2-2  
Observed Soil Moisture Status (NASIS) • 1-14 • *See also* (Soil)  
    Water State  
Odor • 2-90, 2-106  
    Intensity • 2-90, 2-106  
    Kind • 2-90, 2-106  
Organic Deposits • 1-20  
Organics • 1-24, 5-3  
Ortstein • 1-28, 2-44  
Ortstein Fragments • 1-27, 2-46  
Oxidation / Reduction • 2-10

## **P**

---

- Paralithic Contact • 1-29
- Paralithic Materials • 1-29
- Pararock Fragments • 2-39
- Parent Material • 1-18
- Particle Size Classes • 2-45
- Ped and Void Surface Features • 2-32
  - Amount • 2-35
  - Color • 2-36
  - Continuity • 2-35
  - Distinctness • 2-35
  - Kind • 2-32
  - Location • 2-36
- Pedogenic Carbonate Development Stages
  - Coarse Fragment Matrix • 2-31
  - Fine Earth Matrix • 2-30
- Pedogenic Carbonate Stages (Discussion) • 2-28
- Pedon Description
  - Example • 2-95
  - Form • 2-93
- Pendant • 2-27
- Penetrability (Artifacts) • 2-51
- Penetration Orientation • 2-68
- Penetration Resistance • 2-67
- Penetration Resistance Class • 2-68
- Periglacial (Geomorphologic Environment; Landforms) • 3-28
- Permeability Classes • 2-85
- Permeability (Discussion) • 2-81
- Peroxide Color Change • 2-104
- Persistence (Artifacts) • 2-51
- Petrocalcic Horizon • 1-28, 2-44
- Petroferric • 1-29, 2-44
- Petrogypsic Horizon • 1-28, 2-44
- pH • 2-85, 2-102
  - Method • 2-86, 2-102
- Physical Crusts • 2-78
- Physiographic Division • 1-4, 3-1, 3-2
- Physiographic Location • 1-4, 3-1, 3-2
- Physiographic Province • 1-4, 3-1, 3-2
- Physiographic Section • 1-4, 3-1, 3-2
- Phytoliths (plant) • 2-22
- Pipestems • 2-25
- Piping (Erosion) • 1-25
- Placic Horizon • 1-28
- Plant Common Name • 1-18
- Plant Opal • See Phytoliths
- Plant Scientific Name • 1-18
- Plant Symbol • 1-17
- Plasticity • 2-66
- Plasticity Class • 2-66

- Plinthite • 1-29, 2-25
- Pocket Penetrometer • 2-67
- Ponding • 1-14
  - Depth • 1-14
  - Duration • 1-14
  - Frequency • 1-14
- Pores • 2-73
  - Quantity • 2-70
  - Quantity Class • 2-70
  - Shape • 2-73, 2-74
  - Size • 2-70
  - Size Classes • 2-70, 2-72
  - Vertical Continuity • 2-75
- Pores (Discussion) • 2-73
- Precipitates • 1-24, 5-3, 5-7
- Preferential Flow • 2-75
- Pressure Faces • 2-32, 2-34
- Primary Packing Voids • 2-74
- Prime (The) (Horizon Nomenclature) • 2-5, 2-104
- Profile / Pedon Description • 2-1
  - References • 2-115
- Public Land Survey • 6-2
- Pyroclastic Terms • 5-11

## **Q**

---

- quadrangle • 1-2

## **R**

---

- Raindrop Impact Crust • 2-78
- Range (location) • 6-3
- Reaction Oxidized pH • 2-105
  - Discussion • 2-105
- Reaction (pH) • 2-85
- Reaction to Alpha-dipyridil (NASIS) • See Reduced Conditions
- Reaction to H<sub>2</sub>O<sub>2</sub> (Peroxide Color Change) • 2-104
- Redox Concentrations • 2-10, 2-12
- Redox Depletions • 2-10, 2-12
- Redoximorphic Features • 2-12
  - Boundary • 2-17
  - Color • 2-17
  - Contrast • 2-15
  - Hardness • 2-17
  - Kind • 2-12
  - Location • 2-17
  - Moisture State • 2-17
  - Quantity • 2-13
  - Shape • 2-17
  - Size • 2-13

- Redoximorphic Features-RMF (Discussion) • 2-10
- Reduced Conditions • 2-88
- Reduced Matrix • 2-11, 2-12
- Reduced Monosulfide Presence • 2-104
- Reference Samples • 8-3
- Relative\_Exposure\_UOM • 2-2
- Relative Size (of Exposure) • 2-2
- Relative Slope Segment Position • 1-6, 3-39
- Restriction • 1-30
  - Kind • 1-30
- Rill (Erosion) • 1-25
- Rill (Microfeature) • 3-18, 3-33, 3-34
- Riser Length • 2-111
- RMF Shapes • 2-26
- Rock and Other Fragments • 2-46
  - Hardness • 2-49
  - Kind • 2-46
  - Roundness • 2-48
  - Size Classes and Descriptive Terms • 2-47
  - Volume Percent • 2-47
- Rock Charts • 5-4
- Rock Fragments • 2-38, 2-43, 2-46
  - Quantity • 2-38
  - Size • 2-38
- Root Channels • 2-73
- Root Pseudomorphs • 2-25
- Roots • 2-70
  - Location • 2-72
  - Quantity • 2-70
  - Quantity Class • 2-70
  - Size • 2-70
  - Size Classes • 2-70, 2-72
- Roundness (Artifacts) • 2-50
- Roundness (Rock and Other Fragments) • 2-48, 2-49
- Rupture Resistance • 2-62
  - Blocks, Peds, And Clods • 2-63
  - Surface Crusts and Plates • 2-64

## **S**

---

- Safety (Artifacts) • 2-52
- Salinity • 2-88, 2-102, 2-106
- Salinity Class • 2-89
- Sampling Techniques • 8-3
- Sampling, Types of • 8-1
  - Fixed-Depth Sampling • 8-2
  - Horizon Sampling • 8-1
  - Incremental Sampling • 8-2
- Sand Coats • 2-32
- Satiation • 1-15

Saturated Hydraulic Conductivity • 2-83  
Saturated Hydraulic Conductivity Classes • 2-84  
Saturated Hydraulic Conductivity (Discussion) • 2-81  
Saturation • 1-15  
Scale (topographic quadrangle) • 1-2  
Scientific Plant Name • 1-18  
Secondary Carbonates • 1-30  
Section (location) • 6-3  
Sedimentary and Volcaniclastic Rocks Chart • 5-7  
Sedimentary Rocks • 1-23, 5-2  
Series Name • 1-4  
Shape (Artifacts) • 2-51  
Sheet (Erosion) • 1-25  
Shell Fragments • 2-22  
Shot • 2-25  
Silt Coats • 2-32  
Site Description • 1-1, 2-98, 2-111  
Site/Pedon ID • 2-98  
Skeletans • 2-32  
Slake Test • 2-40  
Slickensides • 1-29, 2-33  
Slope Aspect • 1-5, 3-39,  
Slope Complexity • 1-5, 3-39,  
Slope Gradient • 1-5, 3-39,  
Slope (Landforms) • 3-32  
Slope Shape • 1-6, 3-40,  
Smeariness • 2-65  
Sodium Adsorption Ratio (SAR) • 2-89  
Soft, Powdery Lime • See Secondary Carbonates  
Soft Rock Fragments • See Pararock Fragments  
Soil Color • 2-8  
Soil Crusts • 2-79  
    Kind • 2-79  
Soil Crusts (Discussion) • 2-77  
Soil Drainage Class • See Drainage Class  
Soil Matrix Color • 2-9  
Soil Matrix Color - Location or Condition • 2-9  
Soil Moisture Regimes • 4-11  
Soil Moisture Status • See Soil Water State  
Soil Permeability • 2-81  
Soil Sample Kinds • 8-3  
Soil Sampling • 8-1  
    References • 8-6  
Soilscape Sequences • 8-1  
Soil Series Name • 1-4  
Soil Structure • 2-52  
    Grade • 2-55  
    Size • 2-55  
    Type • 2-52  
Soil Surface • 2-6  
Soil Survey Area Identification Number (SSID) • 1-2

- Soil Taxonomy • 4-1
- References • 4-14
- Soil Taxonomy Classification • 1-30
- Soil Temperature • 1-1
- Soil Temperature Depth • 1-1
- Soil Temperature Regimes and Classes • 4-13
- Soil Textural Triangle • See Textural Triangle
- Soil Texture • 2-36
- Soil Water Repellency • 7-15
- Soil Water Repellency (Discussion) • 7-14
- Solution (Geomorphic Environment; Landforms) • 3-25
- Special Features • 2-80
- Sphericity • 2-48
- Sponge Spicules • 2-22
- State Physiographic Area • 1-4, 3-1, 3-9
- State Plane Coordinate System • 6-7
- Stickiness • 2-66
- Stickiness Class • 2-66
- Stone Line • 2-80
- Stop Number (Transect) • 1-4
- Stress Features • 2-32, 2-34
- Stringers • 2-25
- Structure • See Soil Structure
- Structure Shape • See Soil Structure - Type
- Subaqueous Soils • 2-97
- Subaqueous Soils Profile Description
  - Example • 2-110
  - Form • 2-109
- Submerged Aquatic Vegetation • 2-101
- Subordinate Distinctions • See Horizon Suffixes
- Sulfides
  - Discussion • 2-106
- Surface Coats • 2-12
- Surface Crust and Plates Classes (Rupture Resistance) • 2-64
- Surface Crust and Plates (Rupture Resistance) • 2-62
- Surface Crusts • 2-77 • See *also* Soil Crusts
- Surface Fragments • 1-26
  - Class • 1-26
  - Kind • 1-26
  - Mean Distance Between Fragments • 1-26
  - Shape • 1-26
  - Size • 1-26
- Surface Morphometry • 1-5, 3-39,

## **T**

---

- Table Comparing Particle Size Systems • 2-45
- Tabular List for Determination of Color Contrast • 2-16
- Tectonic and Structural (Geomorphic Environments; Landforms) • 3-31

Terms Used In Lieu of Texture • 2–43  
Terraces and Stepped Landforms (Geomorphic Components of) • 1–8  
Terraces, Stepped Landforms (Geomorphic Components of) • 3–42  
Textural Triangle  
    Combined Textural Triangles: Fine Earth Texture Classes and Family Particle-Size Classes • 4–11  
    Family Particle-Size Classes • 4–10  
    Fine Earth Texture Classes • 2–38  
Texture Class • 2–37  
Texture Modifiers • 2–38, 2–39  
Texture Modifiers (Compositional) • 2–41  
Threadlike • 2–25  
Tidal Datum (Discussion) • 2–99  
Tidal Period • 2–111  
Till Terms • 5–10  
Time  
    End • 2–98  
    Start • 2–98  
Time Scale • See North American Geologic Time Scale  
Tongues • 2–80  
Topographic Quadrangle • 1–2  
Total Pipe Length • 2–111  
Townships (location) • 6–3  
Transects • 1–3  
    Delineation Size • 1–3  
    ID • 1–4  
    Kind • 1–3  
Trans-Horizon Cracks • 2–75, 2–76  
Transitional Horizons • 2–2, 4–1  
Transported Deposits • 1–21  
Tubular Pores • 2–73, 2–74  
Tunnel (Erosion) • 1–25

## **U**

---

Unconfined Compressive Strength • 2–67  
Undulation • 2–6  
Universal Transverse Mercator (UTM) Rectangular Coordinate System • 6–5

## **V**

---

Variegated (color) • 2–9  
Vegetation • 1–17  
Vegetation Cover • 1–18  
Vertical Datum  
    Discussion • 2–100  
Vesicular Crust • 2–78

Vesicular Pores • 2-74  
Vibracore Log Sheet  
    Example • 2-114  
    Form • 2-113  
Vibracore Sampling for Subaqueous Soils (Discussion) • 2-111  
Volcanic and Hydrothermal (Geomorphic Environment; Landforms)  
    • 3-30  
Volcanic Deposits • 1-20  
Volcaniclastic Rock Terms • 5-7  
Vughs • 2-73, 2-74

## **W**

---

Water Bodies (Landforms) • 3-37  
Water Column Measurements • 2-102  
Water Depth • 2-100, 2-111  
Water Laid Deposits • 1-21  
Water Repellent Layer • 2-80  
Water Status • 1-11  
Water Table • See Depth to Water Table, Seasonal High Water Table  
Water Temperature • 2-102  
Waypoint • 2-98  
Wedge Structure • 2-53, 2-54  
Wetland (Landforms) • 3-36  
Where Core Is Stored • 2-112