

FORMULARIO DE CÁLCULO DIFERENCIAL E INTEGRAL

VER.3.7
 Jesús Rubi Miranda (jesusrubi1@yahoo.com)
<http://mx.geocities.com/estadisticapapers/>
<http://mx.geocities.com/dicalculus/>

VALOR ABSOLUTO

$|a| = \begin{cases} a & \text{si } a \geq 0 \\ -a & \text{si } a < 0 \end{cases}$
 $|a| = |-a|$
 $a \leq |a|$
 $|a| \geq 0$
 $|a| \geq 0$ y $|a| = 0 \Leftrightarrow a = 0$
 $|ab| = |a||b|$ ó $\prod_{k=1}^n |a_k| = \left| \prod_{k=1}^n a_k \right|$

$|a+b| \leq |a|+|b|$ ó $\left| \sum_{k=1}^n a_k \right| \leq \sum_{k=1}^n |a_k|$

EXONENTES

$a^p \cdot a^q = a^{p+q}$
 $\frac{a^p}{a^q} = a^{p-q}$
 $(a^p)^q = a^{pq}$
 $(a \cdot b)^p = a^p \cdot b^p$

$\left(\frac{a}{b}\right)^p = \frac{a^p}{b^p}$
 $a^{p/q} = \sqrt[q]{a^p}$

LOGARITMOS

$\log_a N = x \Leftrightarrow a^x = N$
 $\log_a MN = \log_a M + \log_a N$
 $\log_a \frac{M}{N} = \log_a M - \log_a N$
 $\log_a N^r = r \log_a N$
 $\log_a N = \frac{\log_b N}{\log_b a} = \frac{\ln N}{\ln a}$
 $\log_{10} N = \log N$ y $\log_e N = \ln N$

ALGUNOS PRODUCTOS

$a \cdot (c+d) = ac+ad$
 $(a+b) \cdot (a-b) = a^2 - b^2$
 $(a+b) \cdot (a+b) = (a+b)^2 = a^2 + 2ab + b^2$
 $(a-b) \cdot (a-b) = (a-b)^2 = a^2 - 2ab + b^2$
 $(x+b) \cdot (x+d) = x^2 + (b+d)x + bd$
 $(ax+b) \cdot (cx+d) = acx^2 + (ad+bc)x + bd$
 $(a+b) \cdot (c+d) = ac+ad+bc+bd$
 $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$
 $(a-b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$
 $(a+b+c)^2 = a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$
 $(a-b) \cdot (a^2 + ab + b^2) = a^3 - b^3$
 $(a-b) \cdot (a^3 + a^2b + ab^2 + b^3) = a^4 - b^4$
 $(a-b) \cdot (a^4 + a^3b + a^2b^2 + ab^3 + b^4) = a^5 - b^5$
 $(a-b) \cdot \left(\sum_{k=1}^n a^{n-k} b^{k-1} \right) = a^n - b^n \quad \forall n \in \mathbb{N}$

$(a+b) \cdot (a^2 - ab + b^2) = a^3 + b^3$
 $(a+b) \cdot (a^3 - a^2b + ab^2 - b^3) = a^4 - b^4$
 $(a+b) \cdot (a^4 - a^3b + a^2b^2 - ab^3 + b^4) = a^5 + b^5$
 $(a+b) \cdot (a^5 - a^4b + a^3b^2 - a^2b^3 + ab^4 - b^5) = a^6 - b^6$

$(a+b) \cdot \left(\sum_{k=1}^n (-1)^{k+1} a^{n-k} b^{k-1} \right) = a^n + b^n \quad \forall n \in \mathbb{N}$ impar
 $(a+b) \cdot \left(\sum_{k=1}^n (-1)^{k+1} a^{n-k} b^{k-1} \right) = a^n - b^n \quad \forall n \in \mathbb{N}$ par

SUMAS Y PRODUCTOS

$a_1 + a_2 + \dots + a_n = \sum_{k=1}^n a_k$

$\sum_{k=1}^n c = nc$

$\sum_{k=1}^n ca_k = c \sum_{k=1}^n a_k$

$\sum_{k=1}^n (a_k + b_k) = \sum_{k=1}^n a_k + \sum_{k=1}^n b_k$

$\sum_{k=1}^n (a_k - a_{k-1}) = a_n - a_0$

$\sum_{k=1}^n [a + (k-1)d] = \frac{n}{2}[2a + (n-1)d]$

$\frac{n}{2}(a+l)$

$\sum_{k=1}^n ar^{k-1} = a \frac{1-r^n}{1-r} = \frac{a-r^n}{1-r}$

$\sum_{k=1}^n k = \frac{1}{2}(n^2 + n)$

$\sum_{k=1}^n k^2 = \frac{1}{6}(2n^3 + 3n^2 + n)$

$\sum_{k=1}^n k^3 = \frac{1}{4}(n^4 + 2n^3 + n^2)$

$\sum_{k=1}^n k^4 = \frac{1}{30}(6n^5 + 15n^4 + 10n^3 - n)$

$1+3+5+\dots+(2n-1) = n^2$

$n! = \prod_{k=1}^n k$

$\binom{n}{k} = \frac{n!}{(n-k)!k!}, \quad k \leq n$

$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$

$(x_1 + x_2 + \dots + x_n)^n = \sum \frac{n!}{n_1!n_2!\dots n_k!} x_1^{n_1} \cdot x_2^{n_2} \cdot \dots \cdot x_n^{n_n}$


CONSTANTES

$\pi = 3.14159265359\dots$
 $e = 2.71828182846\dots$

TRIGONOMETRÍA

$\sin \theta = \frac{CO}{HIP}$ $\csc \theta = \frac{1}{\sin \theta}$
 $\cos \theta = \frac{CA}{HIP}$ $\sec \theta = \frac{1}{\cos \theta}$
 $\operatorname{tg} \theta = \frac{\sin \theta}{\cos \theta} = \frac{CO}{CA}$ $\operatorname{ctg} \theta = \frac{1}{\operatorname{tg} \theta}$


π radianes = 180°




θ	sen	cos	tg	ctg	sec	csc
0°	0	1	0	∞	1	∞
30°	1/2	$\sqrt{3}/2$	1/ $\sqrt{3}$	$\sqrt{3}$	2/ $\sqrt{3}$	2
45°	1/ $\sqrt{2}$	1/ $\sqrt{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
60°	$\sqrt{3}/2$	1/2	$\sqrt{3}$	1/ $\sqrt{3}$	2	2/ $\sqrt{3}$
90°	1	0	∞	0	∞	1

$y = \angle \text{sen } x \quad y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
 $y = \angle \text{cos } x \quad y \in [0, \pi]$
 $y = \angle \operatorname{tg } x \quad y \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
 $y = \angle \operatorname{ctg } x = \angle \operatorname{tg} \frac{1}{x} \quad y \in (0, \pi)$
 $y = \angle \operatorname{sec } x = \angle \operatorname{cos} \frac{1}{x} \quad y \in [0, \pi]$
 $y = \angle \operatorname{csc } x = \angle \operatorname{sen} \frac{1}{x} \quad y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$


Gráfica 1. Las funciones trigonométricas: $\text{sen } x$, $\text{cos } x$, $\operatorname{tg } x$:




Gráfica 2. Las funciones trigonométricas $\operatorname{csc } x$, $\operatorname{sec } x$, $\operatorname{ctg } x$:



Gráfica 3. Las funciones trigonométricas inversas $\operatorname{arsen } x$, $\operatorname{arccos } x$, $\operatorname{arctg } x$:



Gráfica 4. Las funciones trigonométricas inversas $\operatorname{arctg } x$, $\operatorname{arcsec } x$, $\operatorname{arccsc } x$:



IDENTIDADES TRIGONOMÉTRICAS

$\operatorname{sen}^2 \theta + \operatorname{cos}^2 \theta = 1$
 $1 + \operatorname{ctg}^2 \theta = \operatorname{csc}^2 \theta$
 $\operatorname{tg}^2 \theta + 1 = \operatorname{sec}^2 \theta$
 $\operatorname{sen}(-\theta) = -\operatorname{sen} \theta$
 $\operatorname{cos}(-\theta) = \operatorname{cos} \theta$
 $\operatorname{tg}(-\theta) = -\operatorname{tg} \theta$
 $\operatorname{sen}(\theta + 2\pi) = \operatorname{sen} \theta$
 $\operatorname{cos}(\theta + 2\pi) = \operatorname{cos} \theta$
 $\operatorname{tg}(\theta + 2\pi) = \operatorname{tg} \theta$
 $\operatorname{sen}(\theta + \pi) = -\operatorname{sen} \theta$
 $\operatorname{cos}(\theta + \pi) = -\operatorname{cos} \theta$
 $\operatorname{tg}(\theta + \pi) = \operatorname{tg} \theta$
 $\operatorname{sen}(\theta + n\pi) = (-1)^n \operatorname{sen} \theta$
 $\operatorname{cos}(\theta + n\pi) = (-1)^n \operatorname{cos} \theta$
 $\operatorname{tg}(\theta + n\pi) = \operatorname{tg} \theta$
 $\operatorname{sen}(n\pi) = 0$
 $\operatorname{cos}(n\pi) = (-1)^n$
 $\operatorname{tg}(n\pi) = 0$

$\operatorname{sen}\left(\frac{2n+1}{2}\pi\right) = (-1)^n$
 $\operatorname{cos}\left(\frac{2n+1}{2}\pi\right) = 0$
 $\operatorname{tg}\left(\frac{2n+1}{2}\pi\right) = \infty$

$\operatorname{sen} \theta = \operatorname{cos}\left(\theta - \frac{\pi}{2}\right)$
 $\operatorname{cos} \theta = \operatorname{sen}\left(\theta + \frac{\pi}{2}\right)$
 $\operatorname{sen}(\alpha \pm \beta) = \operatorname{sen} \alpha \operatorname{cos} \beta \pm \operatorname{cos} \alpha \operatorname{sen} \beta$
 $\operatorname{cos}(\alpha \pm \beta) = \operatorname{cos} \alpha \operatorname{cos} \beta \mp \operatorname{sen} \alpha \operatorname{sen} \beta$

$\operatorname{tg}(\alpha \pm \beta) = \frac{\operatorname{tg} \alpha \pm \operatorname{tg} \beta}{1 \mp \operatorname{tg} \alpha \operatorname{tg} \beta}$
 $\operatorname{sen} 2\theta = 2 \operatorname{sen} \theta \operatorname{cos} \theta$
 $\operatorname{cos} 2\theta = \operatorname{cos}^2 \theta - \operatorname{sen}^2 \theta$
 $\operatorname{tg} 2\theta = \frac{2 \operatorname{tg} \theta}{1 - \operatorname{tg}^2 \theta}$
 $\operatorname{sen}^2 \theta = \frac{1}{2}(1 - \operatorname{cos} 2\theta)$
 $\operatorname{cos}^2 \theta = \frac{1}{2}(1 + \operatorname{cos} 2\theta)$
 $\operatorname{tg}^2 \theta = \frac{1 - \operatorname{cos} 2\theta}{1 + \operatorname{cos} 2\theta}$

$\operatorname{sen} \alpha + \operatorname{sen} \beta = 2 \operatorname{sen} \frac{1}{2}(\alpha + \beta) \cdot \operatorname{cos} \frac{1}{2}(\alpha - \beta)$
 $\operatorname{sen} \alpha - \operatorname{sen} \beta = 2 \operatorname{sen} \frac{1}{2}(\alpha - \beta) \cdot \operatorname{cos} \frac{1}{2}(\alpha + \beta)$
 $\operatorname{cos} \alpha + \operatorname{cos} \beta = 2 \operatorname{cos} \frac{1}{2}(\alpha + \beta) \cdot \operatorname{cos} \frac{1}{2}(\alpha - \beta)$
 $\operatorname{cos} \alpha - \operatorname{cos} \beta = -2 \operatorname{sen} \frac{1}{2}(\alpha + \beta) \cdot \operatorname{sen} \frac{1}{2}(\alpha - \beta)$

$\operatorname{tg} \alpha \pm \operatorname{tg} \beta = \frac{\operatorname{sen}(\alpha \pm \beta)}{\operatorname{cos} \alpha \cdot \operatorname{cos} \beta}$

$\operatorname{sen} \alpha \cdot \operatorname{cos} \beta = \frac{1}{2}[\operatorname{sen}(\alpha - \beta) + \operatorname{sen}(\alpha + \beta)]$
 $\operatorname{sen} \alpha \cdot \operatorname{sen} \beta = \frac{1}{2}[\operatorname{cos}(\alpha - \beta) - \operatorname{cos}(\alpha + \beta)]$
 $\operatorname{cos} \alpha \cdot \operatorname{cos} \beta = \frac{1}{2}[\operatorname{cos}(\alpha - \beta) + \operatorname{cos}(\alpha + \beta)]$


$\operatorname{tg} \alpha \cdot \operatorname{tg} \beta = \frac{\operatorname{tg} \alpha + \operatorname{tg} \beta}{\operatorname{ctg} \alpha + \operatorname{ctg} \beta}$

FUNCIONES HIPERBÓLICAS

$\operatorname{senh } x = \frac{e^x - e^{-x}}{2}$
 $\operatorname{cosh } x = \frac{e^x + e^{-x}}{2}$
 $\operatorname{tgh } x = \frac{\operatorname{senh } x}{\operatorname{cosh } x} = \frac{e^x - e^{-x}}{e^x + e^{-x}}$
 $\operatorname{ctgh } x = \frac{1}{\operatorname{tgh } x} = \frac{e^x + e^{-x}}{e^x - e^{-x}}$
 $\operatorname{sech } x = \frac{1}{\operatorname{cosh } x} = \frac{2}{e^x + e^{-x}}$
 $\operatorname{csch } x = \frac{1}{\operatorname{senh } x} = \frac{2}{e^x - e^{-x}}$

$\operatorname{senh} : \mathbb{R} \rightarrow \mathbb{R}$
 $\operatorname{cosh} : \mathbb{R} \rightarrow [1, \infty)$
 $\operatorname{tgh} : \mathbb{R} \rightarrow \langle -1, 1 \rangle$
 $\operatorname{ctgh} : \mathbb{R} - \{0\} \rightarrow \langle -\infty, -1 \rangle \cup \langle 1, \infty \rangle$
 $\operatorname{sech} : \mathbb{R} \rightarrow \langle 0, 1 \rangle$
 $\operatorname{csch} : \mathbb{R} - \{0\} \rightarrow \mathbb{R} - \{0\}$

Gráfica 5. Las funciones hiperbólicas $\operatorname{senh } x$, $\operatorname{cosh } x$, $\operatorname{tgh } x$:



FUNCS HIPERBÓLICAS INVERSAS

$\operatorname{senh}^{-1} x = \ln(x + \sqrt{x^2 + 1}), \quad \forall x \in \mathbb{R}$
 $\operatorname{cosh}^{-1} x = \ln(x + \sqrt{x^2 - 1}), \quad x \geq 1$
 $\operatorname{tgh}^{-1} x = \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right|, \quad |x| < 1$
 $\operatorname{ctgh}^{-1} x = \frac{1}{2} \ln \left| \frac{x+1}{x-1} \right|, \quad |x| > 1$
 $\operatorname{sech}^{-1} x = \ln \left(\frac{1 \pm \sqrt{1-x^2}}{x} \right), \quad 0 < x \leq 1$
 $\operatorname{csch}^{-1} x = \ln \left(\frac{1}{x} + \frac{\sqrt{x^2+1}}{|x|} \right), \quad x \neq 0$

IDENTIDADES DE FUNCIONES HIP

$\cosh^2 x - \sinh^2 x = 1$
 $1 - \operatorname{tgh}^2 x = \operatorname{sech}^2 x$
 $\operatorname{ctgh}^2 x - 1 = \operatorname{csch} x$
 $\sinh(-x) = -\sinh x$
 $\cosh(-x) = \cosh x$
 $\operatorname{tgh}(-x) = -\operatorname{tgh} x$
 $\sinh(x \pm y) = \sinh x \cosh y \pm \cosh x \sinh y$
 $\cosh(x \pm y) = \cosh x \cosh y \pm \sinh x \sinh y$
 $\operatorname{tgh}(x \pm y) = \frac{\operatorname{tgh} x \pm \operatorname{tgh} y}{1 \pm \operatorname{tgh} x \operatorname{tgh} y}$
 $\sinh 2x = 2 \sinh x \cosh x$
 $\cosh 2x = \cosh^2 x + \sinh^2 x$
 $\operatorname{tgh} 2x = \frac{2 \operatorname{tgh} x}{1 + \operatorname{tgh}^2 x}$

$\sinh^2 x = \frac{1}{2}(\cosh 2x - 1)$
 $\cosh^2 x = \frac{1}{2}(\cosh 2x + 1)$
 $\operatorname{tgh}^2 x = \frac{\cosh 2x - 1}{\cosh 2x + 1}$
 $\operatorname{tgh} x = \frac{\sinh 2x}{\cosh 2x + 1}$
 $e^x = \cosh x + \sinh x$
 $e^{-x} = \cosh x - \sinh x$

OTRAS

$ax^2 + bx + c = 0$
 $\Rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
 $b^2 - 4ac = \text{discriminante}$

$\exp(\alpha \pm i\beta) = e^\alpha (\cos \beta \pm i \operatorname{sen} \beta)$ si $\alpha, \beta \in \mathbb{R}$

LÍMITES

$\lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e = 2.71828...$
 $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e$
 $\lim_{x \rightarrow 0} \frac{\operatorname{sen} x}{x} = 1$
 $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0$
 $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$
 $\lim_{x \rightarrow 1} \frac{x-1}{\ln x} = 1$

DERIVADAS

$D_x f(x) = \frac{df}{dx} = \lim_{\Delta x \rightarrow 0} \frac{f(x+\Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$
 $\frac{d}{dx}(c) = 0$
 $\frac{d}{dx}(cx) = c$
 $\frac{d}{dx}(cx^n) = ncx^{n-1}$
 $\frac{d}{dx}(u \pm v \pm w \pm \dots) = \frac{du}{dx} \pm \frac{dv}{dx} \pm \frac{dw}{dx} \pm \dots$
 $\frac{d}{dx}(cu) = c \frac{du}{dx}$

$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$
 $\frac{d}{dx}(uvw) = uv \frac{dw}{dx} + uw \frac{dv}{dx} + vw \frac{du}{dx}$
 $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v(du/dx) - u(dv/dx)}{v^2}$
 $\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$
 $\frac{dF}{dx} = \frac{dF}{du} \cdot \frac{du}{dx}$ (Regla de la Cadena)
 $\frac{du}{dx} = \frac{1}{dx/du}$
 $\frac{dF}{dx} = \frac{dF/du}{dx/du}$
 $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{f_2'(t)}{f_1'(t)}$ donde $\begin{cases} x = f_1(t) \\ y = f_2(t) \end{cases}$

DERIVADA DE FUNCIONES LOG & EXP

$\frac{d}{dx}(\ln u) = \frac{du/dx}{u} = \frac{1}{u} \frac{du}{dx}$
 $\frac{d}{dx}(\log u) = \frac{\log e}{u} \cdot \frac{du}{dx}$
 $\frac{d}{dx}(\log_a u) = \frac{\log_a e}{u} \cdot \frac{du}{dx}$ $a > 0, a \neq 1$
 $\frac{d}{dx}(e^x) = e^x$
 $\frac{d}{dx}(a^x) = a^x \ln a$
 $\frac{d}{dx}(u^x) = xu^{x-1} \frac{du}{dx} + \ln u \cdot u^x \cdot \frac{dv}{dx}$

DERIVADA DE FUNCIONES TRIGO

$\frac{d}{dx}(\operatorname{sen} u) = \cos u \frac{du}{dx}$
 $\frac{d}{dx}(\cos u) = -\operatorname{sen} u \frac{du}{dx}$
 $\frac{d}{dx}(\operatorname{tgh} u) = \operatorname{sech}^2 u \frac{du}{dx}$
 $\frac{d}{dx}(\operatorname{ctg} u) = -\operatorname{csc}^2 u \frac{du}{dx}$
 $\frac{d}{dx}(\operatorname{sec} u) = \operatorname{sec} u \operatorname{tgh} u \frac{du}{dx}$
 $\frac{d}{dx}(\operatorname{csc} u) = -\operatorname{csc} u \operatorname{ctg} u \frac{du}{dx}$
 $\frac{d}{dx}(\operatorname{vers} u) = \operatorname{sen} u \frac{du}{dx}$

DERIV DE FUNCIONES TRIGO INVER

$\frac{d}{dx}(\angle \operatorname{sen} u) = \frac{1}{\sqrt{1-u^2}} \cdot \frac{du}{dx}$
 $\frac{d}{dx}(\angle \cos u) = -\frac{1}{\sqrt{1-u^2}} \cdot \frac{du}{dx}$
 $\frac{d}{dx}(\angle \operatorname{tgh} u) = \frac{1}{1+u^2} \cdot \frac{du}{dx}$
 $\frac{d}{dx}(\angle \operatorname{ctg} u) = -\frac{1}{1+u^2} \cdot \frac{du}{dx}$
 $\frac{d}{dx}(\angle \operatorname{sec} u) = \pm \frac{1}{u\sqrt{u^2-1}} \cdot \frac{du}{dx}$ $\begin{cases} + \text{ si } u > 1 \\ - \text{ si } u < -1 \end{cases}$
 $\frac{d}{dx}(\angle \operatorname{csc} u) = \mp \frac{1}{u\sqrt{u^2-1}} \cdot \frac{du}{dx}$ $\begin{cases} - \text{ si } u > 1 \\ + \text{ si } u < -1 \end{cases}$
 $\frac{d}{dx}(\angle \operatorname{vers} u) = \frac{1}{\sqrt{2u-u^2}} \cdot \frac{du}{dx}$

DERIVADA DE FUNCIONES HIPERBÓLICAS

$\frac{d}{dx} \operatorname{senh} u = \cosh u \frac{du}{dx}$
 $\frac{d}{dx} \operatorname{cosh} u = \operatorname{senh} u \frac{du}{dx}$
 $\frac{d}{dx} \operatorname{tgh} u = \operatorname{sech}^2 u \frac{du}{dx}$
 $\frac{d}{dx} \operatorname{ctgh} u = -\operatorname{csch}^2 u \frac{du}{dx}$
 $\frac{d}{dx} \operatorname{sech} u = -\operatorname{sech} u \operatorname{tgh} u \frac{du}{dx}$
 $\frac{d}{dx} \operatorname{csch} u = -\operatorname{csch} u \operatorname{ctgh} u \frac{du}{dx}$

DERIVADA DE FUNCIONES HIP INV

$\frac{d}{dx} \operatorname{senh}^{-1} u = \frac{1}{\sqrt{1+u^2}} \cdot \frac{du}{dx}$
 $\frac{d}{dx} \operatorname{cosh}^{-1} u = \frac{\pm 1}{\sqrt{u^2-1}} \cdot \frac{du}{dx}$ $\begin{cases} + \text{ si } \operatorname{cosh}^{-1} u > 0 \\ - \text{ si } \operatorname{cosh}^{-1} u < 0 \end{cases}$
 $\frac{d}{dx} \operatorname{tgh}^{-1} u = \frac{1}{1-u^2} \cdot \frac{du}{dx}$ $|u| < 1$
 $\frac{d}{dx} \operatorname{ctgh}^{-1} u = \frac{1}{1-u^2} \cdot \frac{du}{dx}$ $|u| > 1$
 $\frac{d}{dx} \operatorname{sech}^{-1} u = \frac{\mp 1}{u\sqrt{1-u^2}} \cdot \frac{du}{dx}$ $\begin{cases} - \text{ si } \operatorname{sech}^{-1} u > 0, u \in (0,1) \\ + \text{ si } \operatorname{sech}^{-1} u < 0, u \in (0,1) \end{cases}$
 $\frac{d}{dx} \operatorname{csch}^{-1} u = -\frac{1}{|u|\sqrt{1+u^2}} \cdot \frac{du}{dx}$ $u \neq 0$

INTEGRALES DEFINIDAS, PROPIEDADES

$\int_a^b \{f(x) \pm g(x)\} dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$
 $\int_a^b c f(x) dx = c \cdot \int_a^b f(x) dx$ $c \in \mathbb{R}$
 $\int_a^b f(x) dx = \int_c^b f(x) dx + \int_a^c f(x) dx$
 $\int_a^b f(x) dx = -\int_b^a f(x) dx$
 $\int_a^b f(x) dx = 0$
 $m \cdot (b-a) \leq \int_a^b f(x) dx \leq M \cdot (b-a)$
 $\Leftrightarrow m \leq f(x) \leq M \forall x \in [a, b], m, M \in \mathbb{R}$
 $\int_a^b f(x) dx \leq \int_a^b g(x) dx$
 $\Leftrightarrow f(x) \leq g(x) \forall x \in [a, b]$
 $\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$ si $a < b$

INTEGRALES

$\int adx = ax$
 $\int af(x) dx = a \int f(x) dx$
 $\int (u \pm v \pm w \pm \dots) dx = \int u dx \pm \int v dx \pm \int w dx \pm \dots$
 $\int u dv = uv - \int v du$ (Integración por partes)
 $\int u^n dx = \frac{u^{n+1}}{n+1}$ $n \neq -1$
 $\int \frac{du}{u} = \ln|u|$

INTEGRALES DE FUNCIONES LOG & EXP

$\int e^x dx = e^x$
 $\int a^x dx = \frac{a^x}{\ln a}$ $\begin{cases} a > 0 \\ a \neq 1 \end{cases}$
 $\int u a^u dx = \frac{a^u}{\ln a} \cdot \left(u - \frac{1}{\ln a}\right)$
 $\int u e^u dx = e^u (u-1)$
 $\int \ln u dx = u \ln u - u + \int \frac{1}{u} dx$
 $\int \log_a u dx = \frac{1}{\ln a} (u \ln u - u) = \frac{u}{\ln a} (\ln u - 1)$
 $\int u \log_a u dx = \frac{u^2}{4} \cdot (2 \log_a u - 1)$
 $\int u \ln u dx = \frac{u^2}{4} (2 \ln u - 1)$

INTEGRALES DE FUNCIONES TRIGO

$\int \operatorname{sen} u dx = -\cos u$
 $\int \cos u dx = \operatorname{sen} u$
 $\int \sec^2 u dx = \operatorname{tgh} u$
 $\int \operatorname{csc}^2 u dx = -\operatorname{ctg} u$
 $\int \operatorname{sec} u \operatorname{tgh} u dx = \operatorname{sec} u$
 $\int \operatorname{csc} u \operatorname{ctg} u dx = -\operatorname{csc} u$
 $\int \operatorname{tgh} u dx = -\ln|\cos u| = \ln|\sec u|$
 $\int \operatorname{ctg} u dx = \ln|\operatorname{sen} u|$
 $\int \operatorname{sec} u dx = \ln|\sec u + \operatorname{tgh} u|$
 $\int \operatorname{csc} u dx = \ln|\operatorname{csc} u - \operatorname{ctg} u|$

$\int \operatorname{sen}^2 u dx = \frac{u}{2} - \frac{1}{4} \operatorname{sen} 2u$
 $\int \cos^2 u dx = \frac{u}{2} + \frac{1}{4} \operatorname{sen} 2u$
 $\int \operatorname{tg}^2 u dx = \operatorname{tgh} u - u$
 $\int \operatorname{ctg}^2 u dx = -(\operatorname{ctg} u + u)$
 $\int u \operatorname{sen} u dx = \operatorname{sen} u - u \cos u$
 $\int u \cos u dx = \cos u + u \operatorname{sen} u$

INTEGRALES DE FUNCIONES TRIGO INV

$\int \angle \operatorname{sen} u dx = u \angle \operatorname{sen} u + \sqrt{1-u^2}$
 $\int \angle \cos u dx = u \angle \cos u - \sqrt{1-u^2}$
 $\int \angle \operatorname{tgh} u dx = u \angle \operatorname{tgh} u - \ln|\sqrt{1+u^2}|$
 $\int \angle \operatorname{ctg} u dx = u \angle \operatorname{ctg} u + \ln|\sqrt{1+u^2}|$
 $\int \angle \operatorname{sec} u dx = u \angle \operatorname{sec} u - \ln(u + \sqrt{u^2-1}) = u \angle \operatorname{sec} u - \angle \operatorname{cosh} u$
 $\int \angle \operatorname{csc} u dx = u \angle \operatorname{csc} u + \ln(u + \sqrt{u^2-1}) = u \angle \operatorname{csc} u + \angle \operatorname{cosh} u$

INTEGRALES DE FUNCIONES HIP

$\int \operatorname{senh} u dx = \cosh u$
 $\int \operatorname{cosh} u dx = \operatorname{senh} u$
 $\int \operatorname{sech}^2 u dx = \operatorname{tgh} u$
 $\int \operatorname{csch}^2 u dx = -\operatorname{ctgh} u$
 $\int \operatorname{sech} u \operatorname{tgh} u dx = -\operatorname{sech} u$
 $\int \operatorname{csch} u \operatorname{ctgh} u dx = -\operatorname{csch} u$

$\int \operatorname{tgh} u dx = \ln|\cosh u|$
 $\int \operatorname{ctgh} u dx = \ln|\operatorname{senh} u|$
 $\int \operatorname{sech} u dx = \angle \operatorname{tg}(\operatorname{senh} u)$
 $\int \operatorname{csch} u dx = -\operatorname{ctgh}^{-1}(\cosh u) = \ln \operatorname{tgh} \frac{1}{2} u$

INTEGRALES DE FRAC

$\int \frac{dx}{u^2 + a^2} = \frac{1}{a} \angle \operatorname{tg} \frac{u}{a}$
 $= -\frac{1}{a} \angle \operatorname{ctg} \frac{u}{a}$
 $\int \frac{du}{u^2 - a^2} = \frac{1}{2a} \ln \frac{u-a}{u+a}$ ($u^2 > a^2$)
 $\int \frac{du}{a^2 - u^2} = \frac{1}{2a} \ln \frac{a+u}{a-u}$ ($u^2 < a^2$)
 $\int \frac{du}{\sqrt{a^2 - u^2}} = \angle \operatorname{sen} \frac{u}{a}$
 $= -\angle \operatorname{cos} \frac{u}{a}$

INTEGRALES CON RAIZ

$\int \frac{du}{\sqrt{u^2 \pm a^2}} = \ln(u + \sqrt{u^2 \pm a^2})$
 $\int \frac{du}{u\sqrt{a^2 \pm u^2}} = \frac{1}{a} \ln \left| \frac{u}{a + \sqrt{a^2 \pm u^2}} \right|$
 $\int \frac{du}{u\sqrt{u^2 - a^2}} = \frac{1}{a} \angle \operatorname{cos} \frac{a}{u}$
 $= \frac{1}{a} \angle \operatorname{sec} \frac{u}{a}$

$\int \sqrt{a^2 - u^2} du = \frac{u}{2} \sqrt{a^2 - u^2} + \frac{a^2}{2} \angle \operatorname{sen} \frac{u}{a}$
 $\int \sqrt{u^2 \pm a^2} du = \frac{u}{2} \sqrt{u^2 \pm a^2} + \frac{a^2}{2} \ln(u + \sqrt{u^2 \pm a^2})$

MAS INTEGRALES

$\int e^{au} \operatorname{sen} bu dx = \frac{e^{au} (a \operatorname{sen} bu - b \operatorname{cos} bu)}{a^2 + b^2}$
 $\int e^{au} \operatorname{cos} bu dx = \frac{e^{au} (a \operatorname{cos} bu + b \operatorname{sen} bu)}{a^2 + b^2}$

ALGUNAS SERIES

$f(x) = f(x_0) + f'(x_0)(x-x_0) + \frac{f''(x_0)(x-x_0)^2}{2!}$
 $+ \dots + \frac{f^{(n)}(x_0)(x-x_0)^n}{n!}$: Taylor
 $f(x) = f(0) + f'(0)x + \frac{f''(0)x^2}{2!}$
 $+ \dots + \frac{f^{(n)}(0)x^n}{n!}$: Maclaurin
 $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$
 $\operatorname{sen} x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots + (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!}$
 $\operatorname{cos} x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots + (-1)^{n-1} \frac{x^{2n-2}}{(2n-2)!}$
 $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n-1} \frac{x^n}{n}$
 $\angle \operatorname{tg} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots + (-1)^{n-1} \frac{x^{2n-1}}{2n-1}$